

LESCO

10404-37 7-29-87
10-3-23 + PCNB

A GRANULAR FUNGICIDE FOR CONTROL OF CERTAIN SOIL-BORNE TURFGRASS DISEASES

ACTIVE INGREDIENT:

Pentachloronitrobenzene 12.50%

INERT INGREDIENTS:

..... 87.50%

Total 100.00%

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)* 10.00%

0.60% Ammoniacal Nitrogen

2.40% Urea Nitrogen

7.00% Coated Slow Release Urea Nitrogen

AVAILABLE PHOSPHORIC ACID (P₂O₅) 3.00%

SOLUBLE POTASH (K₂O) 23.00%

SULFUR (S) TOTAL 12.85%

4.50% Sulfur (S) (Free)

8.35% Sulfur (S) (Combined)

INGREDIENTS: Sulfur Coated Urea, Ammonium Phosphate, Sulfate of Potash.

94% of the Urea Nitrogen is derived from Lesco Sulfur Coated Urea.

ACCEPTED
JUL 29 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10404-37

KEEP OUT OF REACH OF CHILDREN
CAUTION
STATEMENT OF PRACTICAL TREATMENT
(First Aid Procedures)
If swallowed, induce vomiting. If on skin, remove contaminated clothing and wash skin with soap and warm water. Inhaled, remove to fresh air. If in eyes, flush with water for 15 minutes.
For further treatment, get medical attention.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION
Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. This material may cause skin irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt while handling product. Wash hands and face after use and prior to eating, drinking or smoking. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands. Keep out of reach of children.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
A. STORAGE INSTRUCTIONS:
Store in original container only, separate from other pesticides, fertilizer, food and feed. Do not store near open flame or heat sources.
B. PESTICIDE DISPOSAL INSTRUCTIONS:
Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
C. CONTAINER DISPOSAL:
Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LAWNS AND TURF

Brown Patch and Dollar Spot: For southern or warm season lawns such as St. Augustine, Bermuda and similar grasses. Make one application in the fall or spring at first indication of infection. Apply 6 lbs. per 1,000 sq. ft. Following application, the treated areas should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. For Dichondra: Treat as above, using 6 lbs. per 1,000 sq. ft. For Northern or cool season lawns such as Rye Grass, Fescue, Bluegrass, Bent or similar grasses. Apply 1.6 to 2.0 lbs. per 1,000 sq. ft. at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. Water treated area lightly to wash chemical down to soil. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

Helminthosporium Leaf Spot: Apply at rate of 4 to 6 lbs. per 1,000 sq. ft. Preventative applications should be made in spring or fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

Snow Mold (Typhula Incarnata, Fusarium Nivale): Apply at the rate of 4 to 6 lbs. per 1,000 sq. ft. using a granular lawn spreader or comparable equipment. Application should be made prior to first snowfall.

Do not graze treated areas or feed clippings to livestock.

CAUTION: For northern or cool season grasses, such as Rye Grass, Fescue, Bluegrass, Bent or similar grass, application is not recommended when temperatures exceed 75° F.

APPROXIMATE LESCO ROTARY SPREADER SETTINGS

# Product	#N	Effective Spreading Width	3rd Hole	Approx. Setting
1000 sq. ft.	1000 sq. ft.	#A, A	OPEN	D1
4	40	2) B	OPEN	D1
6	60	2) C	OPEN	D1

***IMPORTANT:** These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

WARRANTY
LESCO warrants that this fungicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this fungicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

E.P.A. Reg. No. 10404-37
E.P.A. Est. No. 10404-OH-02
#015942

Manufactured by:
LESCO, INC.
20005 Lake Road, P.O. Box 16915
Rocky River, Ohio 44116

Net Weight
50 Lbs.

2281000W