

Atrazine 0.76% Plus Fertilizer

Weed Control with Premium Lawn Food

50 Pounds covers up to 11,000 Sq.Ft. of Turfgrass
Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Dollarweed, Florida Pusley, Lippia, Pigweed
Over Forty-Five More Weeds as Listed
A WEED & FEED FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS LAWNS

ACTIVE INGREDIENTS:

Atrazine, 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	0.76%
Related active triazine compounds	0.02%
OTHER INGREDIENTS:	99.22%
Total:	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to by the poison control center or doctor Do not give anything by mouth to an unconscious person
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing Call a poison control center or doctor for treatment advice
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> Move person to fresh air If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible Call a poison control center or doctor for further treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Loaders, applicators, and other handlers must wear long-sleeved shirt, long pants, shoes and socks.
Do not apply granular lawn products by hand. Avoid contact with hands or skin.
Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Loaders, applicators and other handlers must wear long-sleeved shirt and long pants, shoes and socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS
Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.
This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
When applying multiple products containing atrazine, the total pounds of atrazine applied in pounds of atrazine per acre may not exceed 4 pounds of atrazine from all applications made in a year.
Turfgrass at Residential Sites (including homes, day-care facilities, schools, playgrounds, parks, recreational areas, and sports fields)
This product must be watered in immediately after application. Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing.
- This product must be watered in immediately.
- Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.
For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.
ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.atrazine.org/watershedinfo or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact LESCO for a refund.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa annua)	Cutleaf Evening Primrose	Oxalis (Common and Yellow Woodsorrel)
Annual Sowthistle	Dickcissel	Pigweed
Clover	Dollarweed	Purslane
Black Medic	Florida Beggarweed	Ragweed
Blue Toadflax	Florida Betony	Red Sorrel
Bull Paspalum	(Rattlesnake Weed)	Rescuegrass
Buckhorn Plantain	Florida Pusley	Sandspur
Bur Clover	Fragrant Cudweed	Shepherd's Purse
Carolina Geranium (Cranesbill)	Rabbit Tobacco	Sicklepod
Chickweed (Common and Mouse Ear)	Henbit	Spotted Spurge
Cocklebur	Knockweed	Spurweed
Common Dandelion	Lambsquarters	Stinging Nettle
Common Purslane	Lespedeza	Swinecress
Crabgrass	Lippia (Mintweed)	Venus Looking Glass
Crownfoot Grass	Moneywort	Wild Mustard
Cudweed	Morning Glory	Wood Sorrel
	Narrowleaf Vetch	
	Parsley-Pier	

IMPORTANT: In addition to providing turfgrasses with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from seeds."
FOR WEEDS THAT ARE PRESENT: Includes broadleaf weeds from the list such as Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.
FOR WEEDS GERMINATING FROM SEEDS: Applied before weed seeds germinate, includes grassy weeds from the list such as Crabgrass, Paspalum, and Spotted Spurge.

WHERE TO USE
FOR CONTROL of weeds in St. Augustine and Centipede grass in or on residential lawns, corporate lawns, parks, recreational areas, golf course, cemeteries, and athletic fields.
NOT FOR USE on turf being grown for sale or other commercial use as sod or for commercial seed production or for research purposes.
NOT FOR USE on Bluegrass, Fescue, Bentgrass, Bahiagrass, Ryegrass, Bermudagrass, or Dichondra lawns.

WHEN TO APPLY
Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post-emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good weed control.
For pre-emergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent parbolic wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

HOW TO APPLY
Apply 4.5 lb to 6.0 lb uniformly over 1,000 sq ft of lawn area (equivalent to the required 1.5 - 2.0 lb Atrazine per acre). Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommended rates. Do not apply more than 4 lb a.i. Atrazine (2 applications at 6.0 lb product) per acre per year. Apply to dry grass. This product must be watered in immediately after application.


SUGGESTED SPREADER SETTINGS*
(Recommended Rates in lb per 1,000 sq ft)

Spreader	4.5 lb	6.0 lb
LESCO Rotary	XX	XX
LESCO Pendulum	XX	XX
Cyclone® or Spyrke™XX	XX	XX
Scotts® Speedy Green®	XX	XX
SCOTT'S® R8-A	XX	XX
Lely®	XX	XX
Vicon®	XX	XX

*Settings are approximate. Age, condition of spreader and operator speed can cause variations. Be sure to calibrate your spreader with each application.

USE PRECAUTIONS
Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses since injury to those other grasses may result. Do not apply this product to areas where there are trees, shrubs, or other plants that are susceptible to injury from herbicide drift. Do not apply this product to areas where there are trees, shrubs, or other plants that are susceptible to injury from herbicide drift. Do not apply this product to areas where there are trees, shrubs, or other plants that are susceptible to injury from herbicide drift.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.
EMPTY CONTAINER DISPOSAL: Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.
SPILL: In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.



WITH 44% AVAILABLE NITROGEN MHA ELITE NOVEX PROFESSIONAL TURF FERTILIZER

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	XX XX%
XX XX% Ammoniacal Nitrogen	
XX XX% Urea Nitrogen*	
XX XX% Water Insoluble Nitrogen	
XX XX% Water Soluble Controlled Release Nitrogen	
AVAILABLE PHOSPHATE (P₂O₅)	XX XX%
SOLUBLE POTASH (K₂O)	XX XX%
MAGNESIUM (Mg) total	XX XX%
XX XX% Water Soluble Magnesium	
SULFUR (S) total	XX XX%
XX XX% Free Sulfur (S)	
XX XX% Combined Sulfur (S)	
COPPER (Cu) total	XX XX%
XX XX% Soluble Copper (Cu)	
IRON (Fe) total	XX XX%
XX XX% Soluble Iron (Fe)	
MANGANESE (Mn) total	XX XX%
XX XX% Water Soluble Manganese	
ZINC (Zn) total	XX XX%
XX XX% Soluble Zinc (Zn)	

2/2

NON-AGRICULTURAL REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

DERIVED FROM: XXXX XXXX XXXX

CHLORINE (Cl) Max

XX XX%

*XX% Slowly available Nitrogen from LESCO® Poly Plus® Sulfur Coated Urea

WARRANTY

LESCO Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. LESCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of LESCO and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL LESCO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

Information concerning the raw materials composing this product can be obtained by writing to LESCO Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, OH 44114-1849 referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-a1.com>.

Created: XXXXXX 1013105
Rev: XXXXXX XXX

Net Weight: XX lb (XX kg)

EPA REG. NO. 10404-39

EPA EST. NO. 10404-OH-04 (M), 10404-FL-01 (S)

First letter of lot code indicates manufacturing site

#XXXXXX

F526

Manufactured by: LESCO, Inc.

722 Webster Turn Drive

Sebring, FL 33870

ACCEPTED
7-5-05
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10404-39