

PH 23

10404-44

1041

LESCO Bentgrass

Selective Broadleaf Herbicide FOR USE ON ORNAMENTAL LAWNS AND TURF GRASSES

Contains 0.20 lbs Dicamba, 1.50 lbs MCPP and 0.30 lbs 2,4-D per gallon

ACTIVE INGREDIENTS

*Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic) acid	2.68%
Dimethylamine Salt of Related Compounds	0.67%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	6.64%
***Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	12.97%
INERT INGREDIENTS	
Total 20.00%	
100.00%	

*Equiv. to 2.19% or 0.20 lbs. of (3,6-dichloro-o-anisic) acid per gallon
 **Equiv. to 5.40% or 0.50 lbs. of 2,4-Dichlorophenoxy acetic acid per gallon
 ***Equiv. to 16.48% or 1.5 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon (Isomers Specific by AOAC 14th Edition #6321)

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido traducida completamente"

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF INHALED:** Remove victim to fresh air and apply artificial respirations if indicated. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention immediately. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

See Side/Back Panel For Additional Precautionary Statements

EPA REG NO. 10404-44 EPA EST NO. 10404-OH 2
Contents 5 Gallons

LESCO, INC.

2005 LAKE ROAD - ROCKY RIVER, OHIO 44116 U.S.A.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - May be fatal if swallowed. Corrosive; causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear face shield, goggles or safety glasses, long pants, long-sleeved shirt, shoes and chemical resistant gloves when mixing, loading or applying this product. Avoid contamination of food-stuffs. Wash hands after use and before eating or smoking.

REENTRY AND WORKER SAFETY STATEMENT
 Do not reenter or permit workers to reenter treated areas until sprays have dried.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and near-lake plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. **Groundwater Contamination** - Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling pesticide at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. When cleaning equipment, do not pour the washwater on the ground, spray or drain over a large area away from susceptible plants, wells and other water sources. **IMPORTANT** Do not use on Centipede grass or St. Augustine grass or dichondra, nor lawns or turf where desirable grasses or clovers are present. Do not apply to areas underlaid by roots of desirable trees and shrubs. See paragraph on Seeding.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. Store in a separate area for herbicides. Store above 32° F.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: (Metal) Triple rinse (or equivalent). Then offer for recycling or reconditioning or incineration and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. (Plastic) - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. Do not burn or bury out of sight.

NOTICE

LESCO INC warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

IMPORTANT USE PRECAUTIONS

Avoid drift of spray (coarse sprays are less likely to drift) onto vegetables, flowers, ornamentals, shrubs, trees, and other desirable crop plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable plants or where it may contaminate water. Use of this product on newly established bentgrass may result in undesirable injury. Use of this product on "old" bentgrass varieties may result in undesirable turf injury.

APPLICATION RATES HOW AND WHEN TO USE

Bentgrass Lawns and Turf - Apply as a spray application with a hand held sprayer or to large areas with a suitable power "boom-type" sprayer using 1 to 1.45 fluid ounces in 1 to 3 gallons of water per 1,000 sq. ft. (equiv. to 3 to 4 pints per acre). Apply as a fall spray during growing season. Spray works faster when temperatures are over 60-70° F and weeds are young and actively growing. To avoid spray particle drift adjust spray nozzles as close to turf as practical, spraying only when calm or wind is less than 5 m.p.h. If hardy weeds persist make a second application after three weeks. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid spraying during long excessively dry or hot periods unless adequate irrigation is available. If application is made when soil is quite warm, use the lower rate to reduce potential injury. Do not irrigate within 24 hours of application. Weeds that start growing in the fall such as herbicid, may be controlled more effectively if sprayed at that time. If dandelions persist as a special problem, individual spot treatment with a hand sprayer is recommended. Do not overdo Bentgrass in closely mowed Bentgrass (putting and bowling greens) apply Lesco Bentgrass Selective Broadleaf Herbicide at a maximum rate of 1 fluid ounce per 1,000 sq. ft. preferably in May or through September. Any yellowing will disappear after about one week. In addition to Bentgrass this product will control weeds in varieties of Bluegrass, Ryegrass, Fescue, Bermuda and Zoysia grass. Apply using 1.45 to 2.2 fl. oz./1,000 sq. ft. (equiv. to 4 to 6 pints per acre) Hard-to-kill summer weeds may require up to 6 pints of this product per acre however do not apply this rate to Bentgrass. Do not use this product on Centipede or St. Augustine grasses.

WEEDS CONTROLLED

Bedstraw	Knockweed	Ragweed
Black Medick	Lambquarters	Sheep Sorrel
Buckhorn	Leopards	Shepherd's Purse
Burdock	Mallow	Smartweed
Chickweed	Morning-glory	Spotted
Chicory	Onion	Spurge
Clover	Peppercorn	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Decks	Pinktop	Wild Onion
Ground Ivy	Polygon	Yarrow
Heart-Ail	Polygon Oak	(and many more broadleaf weeds)
Henbit	Purslane	

SEEDING

Do not apply to newly seeded turf until after the second or third mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rates. A good rain or deep watering should occur before seeding. (corrected vers. Acq'd 08-21-81/ This is Amend. to Bentgrass precautionary, dated 10-30-82)