

FOR INSECT CONTROL ON TURFGRASSES

RECOMMENDED APPLICATIONS

ACTIVE INGREDIENT:

1-Methylalhyl 2-[[ethoxy[[1-methylalhyl)-
amino]phosphinothioyl]oxy]benzoate* 1.5%

INERT INGREDIENTS: 98.5%
Total 100.0%

*U.S. Patent No. 3,621,062 Canadian Patent No. 825,198
OFTANOL is a Reg. TM of Bayer AG, Germany

STOP - READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Wash contaminated clothing with soap and hot water before re-use. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

SWALLOWED: Call a physician or Poison Control Center. Give 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush with plenty of water. **IF ON SKIN:** Wash with soap and hot water. **IF INHALED:** Remove to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

PHYSICIAN INFORMATION: OFTANOL 1.5% Granular Insecticide is a cholinesterase inhibitor. Atropine and 2-PAM have been shown to be effective antidotes.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

HOUSEHOLD: Do not re-use empty container. Wrap and put in trash collection.

PESTICIDE DISPOSAL (except household): Pesticide that cannot be used in accordance with label directions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL (except household): Do not use container in connection with food, feed or ornamental. Dispose of in a sanitary landfill or by incineration if permitted by State and local authorities. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. Dispose of pesticide as directed above.

IMPORTANT: Read these entire Directions for Use and Warranty before using LESCO OFTANOL 1.5% Granular insecticide.

LESCO OFTANOL 1.5% Granular is recommended for use on ornamental, commercial and recreational turf grass areas.

APPLICATION EQUIPMENT

Apply OFTANOL 1.5% Granular uniformly over the turf area with granular turf application equipment. Both drop-type and rotary-type may be used to apply OFTANOL 1.5% Granular. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate applicator equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPLICATION TECHNIQUES

OFTANOL 1.5% Granular application using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of OFTANOL 1.5% Granular over the areas and then immediately apply the remaining half at right angles or crosswise. The full recommended rate of OFTANOL 1.5% Granular must be used for maximum benefit. When applying OFTANOL 1.5% Granular, use an appropriate spreader width marking systems to assure treatment of entire turf area. Also make applications when the wind is calm to help prevent missed areas.

WARRANTY

LESCO warrants that this insecticide conforms to the chemical descriptions on its label. When used in accordance with label directions under normal conditions it is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not foreseeable. The user assumes all risks of use. (08-30-81 Fire Ant + Precaution stmt. amend.)

AREA	PEST	DOSAGE OFTANOL 1.5% G	REMARKS Be sure to Read "APPLICATION" paragraphs on this label.
Turf Grasses Non-crop areas (such as fence rows & roadsides)	SOIL INSECTS		Do not apply more than 6 lbs. of OFTANOL 1.5% Granular per 1,000 Sq. Ft. per year or an equivalent amount of 1.45 ounces of Active Ingredient per 1,000 Sq. Ft. regardless of formulation used. Minimum interval between applications must be 30 days.
	White Grub larvae such as Japanese beetle, Black Turf grass Atanidus, Chafers spp. Phyllophaga species: such as Green June beetle, Oriental beetle, Asiatic beetle, Billbug (larval Stage)	3 lbs./1,000 sq. ft. for soil insects	For White Grub and Billbug Larval Control: Water turf area thoroughly immediately after application, using sufficient water to wet soil 1 to 1 1/2" deep.
	Mole Crickets		For Mole Cricket Control: Water immediately after application using 1/2 to 1 inch of water.
	Fire Ants	3 lbs /1,000 sq. ft.	Area Fire Ant Control: Apply specified dosage uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made, only one area treatment may be made.
		0.5 oz. per 9 sq. ft.	Established Fire Ant Mounds - Drench Treatment: Wet Conditions: Distribute dosage evenly over 9 sq. ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for Wet Conditions, however to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment. Consult your local Turf Specialist, State Agricultural Extension Service Station or State Extension Service Specialists for more specific information regarding timing of application.
	SURFACE INSECTS		Apply 1/2 to 1 inch of water immediately after treatment. Reapply as necessary for pest re-infestations. Do not apply more than 6 lbs. OFTANOL 1.5% Granular per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. ft. regardless of formulation used. Minimum interval between applications is 15 days for the 1.5 rate and 30 days for the 3 lb. rate
	Sod Webworm larvae	1.5 or 3 lbs /1,000 sq. ft.	
	Chinch Bugs		
	Hyperodes Weevil (larval stage)	3 lbs /1,000 sq. ft.	For Hyperodes Weevil Control: Apply between 1 1/2 to 2 1/2 inches of Forsythia and Flowering Dogwood, usually April 15th to May 15th and water in as directed above. Consult your local Turf Specialist, State Agricultural Extension Service Station, or State Extension Specialist for more specific information regarding timing of application.

ACCEPTED

SEP 28 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is amended, for the pesticide registered under EPA Reg. No. 10404-47

BEST AVAILABLE COPY

APPROXIMATE LESCO ROTARY SPREADER SETTINGS

Pounds of product per 1,000 sq. ft.	lbs. A/L/A	Effective Spreading Width	3 rd / tote	Approximate Setting
3.0	2	7 FEET	OPEN	D 1/2

IMPORTANT: This setting is approximate. Age, condition of spreader and operator speed can cause wide variations. Be sure to calibrate your spreader with each application.

RESTRICTIONS: Do not mow treated turf areas before rainfall or irrigation has been applied as per recommended applications. Do not graze treated areas or use clippings from treated areas for feed. Keep children and pets off treated area until dry. 10404-47, pm4 08-08-92

EPA REG. NO. 10404-47
#002704

NET WEIGHT 40 POUNDS

EPA EST. NO. 10404-04-1

LESCO, INC. - 2005 Lake Road - Rocky River, Ohio 44116