



PRE-M® 60 DG HERBICIDE

A 60% Water Dispersible Granule for Use As A Preemergent Weed Control in Lawns, Turf and Ornamentals.

ACTIVE INGREDIENT:

Pendimethalin [N (1 ethylpropyl)	
3-4 dimethyl 2, 6-dinitrobenzamine	60.0%
INERT INGREDIENTS:	40.0%
TOTAL	100.0%

One Pound contains 0.6 lbs. of Pendimethalin,
PRE-M® is a Reg. Trademark of Lesco, Inc.

EPA Reg. No. 10404-52

EPA Est. No. 5905-IA 1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center immediately. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation occurs.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. May cause eye and skin irritation. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CHEMIGATION: Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Do not store above 120°F. Store product in its original labeled container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

MIXING AND APPLICATION INSTRUCTIONS

Add LESCO PRE-M® 60 DG slowly to partially filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. **MAINTAIN CONTINUOUS AGITATION WHILE ADDING LESCO PRE-M® 60 DG.** If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. Avoid application when winds may cause drift.

Treatments are most effective in controlling weeds when adequate rain fall or overhead irrigation is received within 7 days after application. Avoid contact of spray solution with driveways, stone, wood or other porous surfaces. Rinse immediately to avoid staining.

COMPATIBILITY

LESCO PRE-M® 60 DG is compatible with most commonly used herbicides. When tank mixing be sure to add LESCO PRE-M® 60 DG to the partially filled tank first, mix thoroughly, and then add combination products to the mixture as directed or as instructed on its EPA approved label. Do not apply tank mix combinations unless product is approved for such tank-mix use and previous experience indicates the mixture is effective and will not result in application problems or plant injury. Do not attempt to mix LESCO PRE-M® 60 DG directly with liquid fertilizers. Premix one part of LESCO PRE-M® 60 DG with two parts water and add this diluted mixture slowly into the tank while agitating.

GENERAL INFORMATION

LESCO PRE-M® 60 DG treatments will not control established weeds. Applications must be made prior to germination of weed seeds. Three applications per year may be made. Allow at least 2 months between applications except when otherwise indicated on this label.


Use only on well established turfgrass with a dense and uniform stand. Do not use on newly seeded or sprigged turfgrass. Do not use on winter-overseeded turfgrass. Do not use on turfgrass severely thinned by winter stress.

LESCO PRE-M® 60 DG provides effective residual weed control which may affect new turfgrass seedlings or sprigs. Do not reseed or winter overseed treated turfgrass within 4 growing months after application. Do not sprig turfgrass within 5 months after application. Do not use on Bentgrass or Dichondra maintained as desirable turf.

WARRANTY LESCO, INC. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Directions For Use continued on back panel

Licensed under one or more of the following U.S. Patents
3,920,742; 4,066,441; 4,511,395



BEST AVAILABLE COPY

PRE-M® 60 DG HERBICIDE

A 60% Wettable Powder for Use As A Preemergent Weed Control in Lawns, Turf and Ornamentals.

Directions For Use (Continued from front panel)

RATE AND APPLICATION CHART

TURF SPECIES	WEEDS CONTROLLED	PRE-M 60 DG RATE / ACRE	COMMENTS
NORTHERN GRASSES			
Fine Fescue Kentucky Bluegrass Perennial Ryegrass Tall Fescue	Barnyardgrass Crabgrass Fall Panicum Foxtail Hop Clover Prostrate Spurge Yellow Woodsorrel Cudweed	2.5 to 3.75 lbs per acre (0.9 to 1.4 oz) per 1,000 sq ft)	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs per acre or 0.9 oz. per 1,000 sq ft.
	Goosegrass Evening Primrose	5.0 lbs. per acre (1.8 oz per 1,000 sq ft)	
	Annual Bluegrass Common Chickweed Mouseear Chickweed Henbit	2.5 to 3.75 lbs (0.9 to 1.4 oz) per 1,000 sq ft)	
SOUTHERN GRASSES			
Bahia Bermudagrass Centipedegrass St. Augustinegrass Tall Fescue Zoysiagrass	Barnyardgrass Crabgrass Fall Panicum Foxtail Prostrate Spurge Yellow Woodsorrel Cudweed	2.5 to 3.75 lbs per acre (0.9 to 1.4 oz per 1,000 sq ft)	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs. per acre or 0.9 oz. per 1,000 sq ft.
	Crabgrass Goosegrass Hop Clover Evening Primrose	5.0 lbs per acre (1.8 oz per 1,000 sq ft)	
	Henbit Annual Bluegrass Common Chickweed Mouseear Chickweed	2.5 to 3.75 lbs per acre (0.9 to 1.4 oz per 1,000 sq ft)	

WEED CONTROL IN AND AROUND ORNAMENTAL PLANTINGS

PRE-M® 60 DG is recommended for use on the following established field grown and container ornamentals, plants, trees, and established ground covers. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of PRE-M® 60 DG and roots. Directed sprays where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Sprays can be safely applied over the top of those plants listed below. Plant only those ornamentals listed on this label into soil treated the previous season with PRE-M® 60 DG or injury may occur. Do not apply to seedbeds. PRE-M® 60 DG may be used for the preemergence control of weeds as listed in and around the following established ornamental plants.

ORNAMENTAL PLANTS

COMMON NAME	SCIENTIFIC NAME
Abelia, Glossy	Abelia grandiflora var. grandiflora
Ash, Red	Fraxinus pennsylvanica
Azalea	Rhododendron indicum
Barberry, Japanese	Berberis thunbergii var. atropurpurea
Birch, European Weeping	Betula pendula
Birch, River	Betula nigra
Buxwood, Common	Buxus sempervirens
Capejasmine	Gardenia jasminoides var. mystery
Cedar, Northern White	Taxa occidentalis
Cedar, Red	Juniperus virginiana
Cornelaster, Rock	Cornus alternifolia
Osage, Common	Ligustrum sinense
Dorwood, Flowering	Cornus florida
Dorwood, Silky	Cornus amomum
Eunonymus	Eunonymus alata var. compacta
Fir, Fraser	Abies fraseri
Fir, White	Abies concolor
Forsythia, Border	Forsythia intermedia
Hemlock, Eastern	Tsuga canadensis
Holly, American	Ilex Opaca var. tsuga
Holly, Chinese	Ilex cornuta
Holly, Japanese	Ilex crenata compacta
Holly, Yaupon	Ilex vomitoria var. shillingii dwarf
Juniper, Chinese	Juniperus chinensis var. pfitzeri var. pfitzeri aurea
Juniper, Shore	Juniperus oerferta
Juniper, Trailing	Juniperus horizontalis var. plumosa var. prince of wales var. youngstown
Laurel, Mountain	Kalmia latifolia
Lilac, Common	Syringa vulgaris
Locust, Honey	Gleditsia Trifacanthos var. inermis
Maple, Red	Acer rubrum
Nandina	Nandina domestica
Oak, Live	Quercus virginiana
Oak, Water	Quercus nigra
Peach	Prunus persica
Photina	Photina trazeii
Pieris, Japanese	Pieris japonica
Pine, Austrian*	Pinus nigra
Pine, Loblolly	Pinus taeda
Pine, White	Pinus strobus
Privet, California	Ligustrum ovalifolium
Privet, Japanese	Ligustrum japonicum
Robira	Ptilosporum tobira
Sourwood	Oxydendron arboreum
Spirea	Spirea vanhouttei
Spruce, Colorado Blue	Picea pungens
Waxmyrtle, Common	Myrica cerifera
Yew	Taxus Media
Yew, Japanese	Taxus cuspidata

* Not labeled for use in California

WEED SPECIES CONTROLLED

The following grass and broadleaf weeds are controlled by preemergent treatments of PRE-M 60 DG at the rates as shown under the section entitled APPLICATION RATES FOR ORNAMENTALS.

GRASSES CONTROLLED

COMMON NAME	SCIENTIFIC NAME
Barnyardgrass	Echinochloa crus-galli
Bluegrass, annual	Poa Annua
Crabgrass	Digitaria, spp
Crowfootgrass	Dactyloctenium aegyptium
Foxtail, Giant	Setaria faberi
Foxtail, Green	Setaria viridis
Foxtail, Yellow	Setaria lutescens
Goosegrass	Elymus Indica
Itchgrass	Rottboellia exaltata
Johnsongrass (1' seed)	Sorghum halepense
Junglerice	Echinochloa colonum
Lovegrass	Eragrostis spp
Panicum, Browntop	Panicum fasciculatum
Panicum, Fall	Panicum dichotomiflorum
Panicum, Texas	Panicum texanum
Sandbur, Field	Cenchrus incertus
Signalgrass	Bracharia platyphylla
Sprangletop, Mexican	Leptochloa univervia
Sprangletop, Red	Leptochloa filiformis
Witchgrass	Panicum capillare
Woolly Cupgrass	Eriochloa villosa

BROADLEAF WEEDS CONTROLLED

COMMON NAME	SCIENTIFIC NAME
Carpetweed	Mollugo verticillata
Chickweed, Common	Stellaria media
Chickweed, Mouseear	Cerastium vulgatum
Clover, Hop	Trifolium procumbens
Cudweed	Gnaphalium spp
Fiddleneck	Amsinckia intermedia
Flaree	Erodium spp
Henbit	Lamium amplexicaule
Knotweed (prostrate)	Polygonum aviculare
Kochia	Kochia scoparia
Lambsquarter	Chenopodium album
Pigweed	Amaranthus spp.
Puncturevine	Tribulus terrestris
Purslane	Portulaca oleracea

APPLICATION RATES FOR ORNAMENTALS

For preemergence control of the weed species listed in the next column, apply PRE-M® 60 DG at the following rates.

Length of Control and Amount of PRE-M 60 DG / Acre
Short Term Control (2-4 months) 3.3 lbs. per Acre
Long-Term Control (6-8 months) 6.6 lbs. per Acre

PRE-M® 60 DG will not control established weeds. The efficacy of PRE-M® 60 DG will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

FOR GROUND COVER ORNAMENTALS

PRE-M® 60 DG may be used for the preemergence control of the weed species listed in and around the following established ground cover plants.

COMMON NAME	SCIENTIFIC NAME
Capeweed	Arctotheca calendula
Daisy, Trailing African	Osteospermum fruticosum
Gazania	Gazania splendens
Rose-of-Sharon	Hypericum calycinum
Iceplant, Large Leaf	Carpobrotus edule
Ivy, English	Hedera helix
Ivy, Geranium	Pelargonium peltatum
Myoporum	Myoporum parviflorum
Pachysandra	Pachysandra terminalis
Potentilla	Potentilla jackmanii
Potentilla	Potentilla verna

PRECAUTION

Apply only to established plantings.
DO NOT apply PRE-M® 60 DG in ornamentals nurseries.

MFG. FOR: LESCO, INC. - 2005 LAKE ROAD - ROCKY RIVER, OHIO 44116

EPA Reg. No. 10404 52
EPA Est. No. 5905 IA 1

BEST AVAILABLE COPY



PRE-M® 60 DG HERBICIDE

A 60% Water Dispersible Granule for Use As A Preemergent Weed Control in Lawns, Turf and Ornamentals.

ACTIVE INGREDIENT:

Pendimethalin [N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzamine] 60.0%

INERT INGREDIENTS: 40.0%
TOTAL 100.0%

One Pound contains 0.6 lbs. of Pendimethalin,
PRE-M® is a Reg. Trademark of Lesco, Inc.

EPA Reg. No. 10404-52 EPA Est. No. 5905-IA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center immediately. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation occurs.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May cause eye and skin irritation. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENT. HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling

CHEMIGATION: Do not apply this product through any type of irrigation system

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Do not store above 120°F. Store product in its original labeled container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

MIXING AND APPLICATION INSTRUCTIONS

Add LESCO PRE-M® 60 DG slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. **MAINTAIN CONTINUOUS AGITATION WHILE ADDING LESCO PRE-M® 60 DG.** If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. Avoid application when winds may cause drift.

BEST AVAILABLE COPY

Treatments are most effective in controlling weeds when adequate rain fall or overhead irrigation is received within 7 days after application. Avoid contact of spray solution with driveways, stone, wood or other porous surfaces. Rinse immediately to avoid staining.

COMPATIBILITY

LESCO PRE-M® 60 DG is compatible with most commonly used herbicides. When tank mixing be sure to add LESCO PRE-M® 60 DG to the partially filled tank first, mix thoroughly, and then add combination products to the mixture as directed or as instructed on its EPA approved label. Do not apply tank mix combinations unless product is approved for such tank-mix use and previous experience indicates the mixture is effective and will not result in application problems or plant injury. Do not attempt to mix LESCO PRE-M® 60 DG directly with liquid fertilizers. Premix one part of LESCO PRE-M® 60 DG with two parts water and add this diluted mixture slowly into the tank while agitating

GENERAL INFORMATION

LESCO PRE-M® 60 DG treatments will not control established weeds. Applications must be made prior to germination of weed seeds. Three applications per year may be made. Allow at least 2 months between applications except when otherwise indicated on this label.

Use only on well established turfgrass with a dense and uniform stand. Do not use on newly seeded or sprigged turfgrass. Do not use on winter-overseeded turfgrass. Do not use on turfgrass severely thinned by winter stress.

LESCO PRE-M® 60 DG provides effective residual weed control which may affect new turfgrass seedlings or sprigs. Do not reseed or winter overseed treated turfgrass within 4 growing months after application. Do not sprig turfgrass within 5 months after application. Do not use on Bentgrass or Dichondra maintained as desirable turf.

WARRANTY LESCO, INC. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Directions For Use continued on back panel

Licensed under: one or more of the following U.S. Patents:
3,920,742; 4,066,441; 4,511,395.

ACCEPTED

JUN - 4 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, certified under EPA Reg. No. 10404-52

PRE-M® 60 DG HERBICIDE

A 60% Wettable Powder for Use As A Preemergent Weed Control in Lawns, Turf and Ornamentals.

Directions For Use (Continued from front panel)

RATE AND APPLICATION CHART

TURF SPECIES	WEEDS CONTROLLED	PRE-M 60 DG RATE / ACRE	COMMENTS
NORTHERN GRASSES			
Five Fescue Kentucky Bluegrass Pennisetum Pycnanthum Tall Fescue	Bermudagrass Crabgrass Fall Panicum Foxtail Hop Clover Prostrate Spurge Yellow Woodruff Cudweed	2.5 to 3.75 lbs. per acre (0.9 to 1.4 oz. per 1,000 sq. ft.)	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs per acre or 0.9 oz. per 1,000 sq. ft.
	Goosegrass Evening Primrose	5.0 lbs. per acre (1.8 oz. per 1,000 sq. ft.)	
	Annual Bluegrass Common Chickweed Mooseear Chickweed Herb	2.5 to 3.75 lbs (0.9 to 1.4 oz. per 1,000 sq. ft.)	Apply in late summer to early fall, before weeds germinate
SOUTHERN GRASSES			
Bahia Bermudagrass Centipede St. Augustine Tall Fescue Zoysia	Bermudagrass Crabgrass Fall Panicum Foxtail Prostrate Spurge Yellow Woodruff Cudweed	2.5 to 3.75 lbs. per acre (0.9 to 1.4 oz. per 1,000 sq. ft.)	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs per acre or 0.9 oz. per 1,000 sq. ft.
	Crabgrass Goosegrass Hop Clover Evening Primrose	5.0 lbs per acre (1.8 oz per 1,000 sq. ft.)	
	Herb Annual Bluegrass Common Chickweed Mooseear Chickweed	2.5 to 3.75 lbs per acre (0.9 to 1.4 oz per 1,000 sq. ft.)	Apply in late summer to early fall, before weeds germinate

WEED CONTROL IN AND AROUND ORNAMENTAL PLANTINGS

PRE-M® 60 DG is recommended for use on the following established field grown and container ornamentals, plants, trees, and established ground covers. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of PRE-M® 60 DG and roots. Directed sprays where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Sprays can be safely applied over the top of those plants listed below. Plant only those ornamentals listed on this label into soil treated the previous season with PRE-M® 60 DG or injury may occur. Do not apply to seedbeds. PRE-M® 60 DG may be used for the preemergent control of weeds as listed in and around the following established ornamental plants

ORNAMENTAL PLANTS

COMMON NAME	SCIENTIFIC NAME
Abelia, Glossy	Abelia grandiflora var. grandiflora
Ash, Red	Fraxinus pennsylvanica
Azalea	Rhododendron indicum
Barberry, Japanese	Barberis thunbergii var. atropurpurea
Birch, European Weeping	Betula pendula
Birch, River	Betula nigra
Boraxwood, Common	Bonus sempervirens
Cape Jasmine	Gardenia jasminoides var. mystery
Cedar, Northern	Thuja occidentalis
Cedar, Red	Juniperus virginiana
Cornflower, Field	Cyanus horridulus
Cornflower, Rock Garden	Lagerströmia indica
Dogwood, Flowering	Cornus florida
Dogwood, Silky	Cornus amomum
Elm, Warty	Elaeagnus alata var. compacta
Fir, Fraser	Abies frazeri
Fir, White	Abies concolor
Forsythia, Border	Forsythia intermedia
Hemlock, Eastern	Tsuga canadensis
Holly, American	Ilex Opaca var. laevis
Holly, Chinese	Ilex cornuta
Holly, Japanese	Ilex crenata compacta
Holly, Yaupon	Ilex vomitoria var. shillings dwarf
Juniper, Chinese	Juniperus chinensis var. pfitzeri var. pfitzeri aurea
Juniper, Shore	Juniperus conferta
Juniper, Trailing	Juniperus horizontalis var. plumosa var. prince of wales var. youngblom
Laurel, Mountain	Kalmia latifolia
Lilac, Common	Syringa vulgaris
Locust, Honey	Gleditsia triacanthos var. inermis
Maple, Red	Acer rubrum
Mandarin	Mandarin domestica
Oak, Live	Quercus virginiana
Oak, Water	Quercus nigra
Peach	Prunus persica
Pine	Pinus strobus
Pine, Japanese	Pinus japonica
Pine, Austrian*	Pinus nigra
Pine, Loblolly	Pinus taeda
Pine, White	Pinus strobus
Privet, California	Ligustrum ovalifolium
Privet, Japanese	Ligustrum japonicum
Rubus	Pteridium foliosum
Sourwood	Oxydendron arboreum
Spiraea	Spiraea verticillata
Spruce, Colorado Blue	Picea pungens
Wormyrtle, Common	Myrica caribaea
Yew	Taxus Media
Yew, Japanese	Taxus canadensis

WEED SPECIES CONTROLLED

The following grass and broadleaf weeds are controlled by preemergent treatments of PRE-M 60 DG at the rates as shown under the section entitled APPLICATION RATES FOR ORNAMENTALS.

GRASSES CONTROLLED

COMMON NAME	SCIENTIFIC NAME
Bermudagrass	Echinochloa crus-galli
Bluegrass, annual	Poa Annua
Crabgrass	Digitaria spp
Crowfootgrass	Dactyloctenium aegyptium
Foxtail, Giant	Setaria faberi
Foxtail, Green	Setaria viridis
Foxtail, Yellow	Setaria faberiana
Goosegrass	Elymus indica
Itchgrass	Rottboellia exaltata
Johnsongrass (seed)	Sorghum halepense
Jungerling	Echinochloa colorem
Lrwgrass	Eragrostis spp
Panicum, Browntop	Panicum fasciculatum
Panicum, Fall	Panicum dichotomiflorum
Panicum, Texas	Panicum texanum
Sandbar, Field	Cenchrus incertus
Signalgrass	Brachiaria platyphylla
Sprangletop, Mexican	Leptochloa univiridis
Sprangletop, Red	Leptochloa illinoensis
Witchgrass	Panicum capillare
Woolly Cupgrass	Eriochloa villosa

BROADLEAF WEEDS CONTROLLED

COMMON NAME	SCIENTIFIC NAME
Carpenterweed	Molligo verticillata
Chickweed, Common	Stellaria media
Chickweed, Mooseear	Cerastium vulgatum
Clover, Hop	Trifolium procumbens
Cudweed	Gratiola spp
Fiddleneck	Ammannia intermedia
Flare	Erodium spp
Herb	Lamium amplexicaule
Knotweed (prostrate)	Polygonum aviculare
Kochia	Kochia scoparia
Lawnsparter	Chenopodium album
Pigweed	Amaranthus spp
Puncturevine	Tribulus terrestris
Purslane	Portulaca olerace

APPLICATION RATES FOR ORNAMENTALS

For preemergence control of the weed species listed in the next column, apply PRE-M® 60 DG at the following rates.

Length of Control and Amount of PRE-M 60 DG / Acre
Short Term Control (2-4 months) 3.3 lbs. per Acre
Long-Term Control (6-8 months) 6.6 lbs. per Acre

PRE-M® 60 DG will not control established weeds. The efficiency of PRE-M® 60 DG will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

FOR GROUND COVER ORNAMENTALS

PRE-M® 60 DG may be used for the preemergence control of the weed species listed in and around the following established ground cover plants.

COMMON NAME	SCIENTIFIC NAME
Capeweed	Arctotheca calendula
Ulex, Trailing Arctium	Oxyspermum frutescens
Gazania	Gazania splendens
Rose-of-Sharon	Hypericum calycinum
Impatiens, Large Leaf	Carpobrotus edulis
Ivy, English	Hedera helix
Ivy, Geranium	Pelargonium peltatum
Mycoporum	Mycoporum parviflorum
Pachysandra	Pachysandra terminalis
Potentilla	Potentilla jackmanii
Potentilla	Potentilla verna

PRECAUTION
Apply only to established plantings
DO NOT apply PRE-M® 60 DG in ornamental nurseries

EPA Reg. No. 10404-52
EPA Est. No. 5905-M-1

MFG. FOR: LESCO, INC. - 2005 LAKE ROAD - ROCKY RIVER, OHIO 44116

BEST AVAILABLE COPY