

# A 60% Water Dispersible Granule for Use As A Pre-emergent Weed Control in Lawns, Turf & Ornamentals.

**BEST AVAILABLE COPY**

**ACTIVE INGREDIENT:**  
Pendimethalin (N-(1-ethylpropyl)-  
3,4-dimethyl-2,6-dinitrobenzamine) . . . . . 60.0%  
40.0%

**INERT INGREDIENTS:** . . . . . TOTAL 100.0%

One Pound contains 0.8 lbs. Pendimethalin. PRE-M® is a Reg. Trademark of Lesco, Inc.  
EPA Reg. No. 10404-52 EPA Est. No. 33967-NJ-1 or Est. No. 5905-IA-1

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENTS OF PRACTICAL TREATMENT

**IF SWALLOWED:** Do not induce vomiting. Call a physician or Poison Control Center immediately.  
**IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation occurs.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May cause eye and skin irritation. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **CHEMIGATION:** Do not apply this product through any type of irrigation system.

### STORAGE AND DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Do not store above 120°F. Store product in its original labeled container in a cool, dry, locked place out of reach of children.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### MIXING AND APPLICATION INSTRUCTIONS

Add LESCO PRE-M® 60 DG slowly to a partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. Maintain continuous agitation while adding the LESCO PRE-M 60 DG. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed. Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. Avoid application when winds may cause spray drift.

Treatments are most effective in controlling weeds when adequate rain fall or head irrigation is received within 7 days after application. Avoid contact of spray solution with driveways, stone, wood or other porous surfaces. Rinse immediately to avoid staining.

### COMPATIBILITY

LESCO PRE-M® 60 DG is compatible with most commonly used herbicides. When tank mixing be sure to add LESCO PRE-M® 60 DG to the partially filled tank first, mix thoroughly, and then add combination products to the mixture as directed or as instructed on its EPA approved label. Do not apply tank mix combinations unless product is approved for such tank-mix use and previous experience indicates the mixture is effective and will not result in application problems or plant injury. Do not attempt to mix LESCO PRE-M® 60 DG directly with liquid fertilizers. Pre-mix one part of LESCO PRE-M 60 DG with two parts water and add this diluted mixture slowly into the tank while agitating.

### GENERAL INFORMATION

LESCO PRE-M® 60 DG treatments will not control established weeds. Applications must be made prior to germination of weed seeds. Three applications per year may be made. Allow at least 2 months between applications except when otherwise indicated on this label. Use only on well established turfgrass with a dense and uniform stand. Do not use on newly seeded or sprigged turfgrass. Do not use on winter-overseeded turfgrass. Do not use on turfgrass severely thinned by winter stress. LESCO PRE-M® 60 DG provides effective residual weed control which may affect new turfgrass seedlings or sprigs. Do not reseed or winter over-seed treated turfgrass within 4 growing months after application. Do not sprig turfgrass within 5 months after application. Do not use on Bentgrass or Dichondra maintained as desirable turf.

### APPLICATION RATES FOR ORNAMENTALS

For preemergence control of the weed species listed in the next column, apply PRE-M® 60 DG at the following rates:

**Length of Control and Amount of PRE-M 60 DG/ Acre**  
Short Term Control (2-4 months) 3.3 lbs. per Acre  
Long Term Control (6-9 months) 6.6 lbs. per Acre

PRE-M® 60 DG will not control established weeds.

The efficacy of PRE-M® 60 DG will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

**FOR SPRINKLER COVERED ORNAMENTALS**

### RATE AND APPLICATION CHART FOR LAWNS & TURF

| TURF SPECIES   | WEEDS CONTROLLED  | PRE-M 60 DG RATE /ACRE  | COMMENTS   |
|--|---|---|--|
| <b>NORTHERN GRASSES</b>  |   |   |  |
| Fine Fescue<br>Kentucky Bluegrass<br>Perennial Ryegrass<br>Tall Fescue                     | Barnyardgrass<br>Crabgrass<br>Fall Panicum<br>Foxtail<br>Hop Clover<br>Prostrate Spurge<br>Yellow Woodsorrel<br>Cudweed<br>Goosegrass<br>Evening Primrose | 2.5 to 3.75 lbs. per acre<br>(0.9 to 1.4 oz. per 1,000 sq ft) | Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs per acre or 0.9 oz./1,000 sq ft.  |
|  | Annual Bluegrass<br>Common Chickweed<br>Henbit<br>Mouse ear Chickweed   | 5.0 lbs per acre<br>(1.8 oz per 1,000 sq ft)                  | Apply in late summer to early fall, before weeds germinate.  |
| <b>SOUTHERN GRASSES</b>  |   |   |  |
| Bahua<br>Bermudagrass<br>Centipedegrass<br>St. Augustinegrass<br>Tall Fescue<br>Zoysagrass | Barnyardgrass<br>Crabgrass<br>Fall Panicum<br>Foxtail<br>Prostrate Spurge<br>Yellow Woodsorrel<br>Cudweed   | 2.5 to 3.75 lbs per acre<br>(0.9 to 1.4 oz per 1,000 sq ft)   | Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs per acre or 0.9 oz / 1,000 sq ft. |
|  | Crabgrass<br>Goosegrass<br>Hop Clover<br>Evening Primrose   | 5.0 lbs per acre<br>(1.8 oz per 1,000 sq ft)                  |  |
|  | Annual Bluegrass<br>Common Chickweed<br>Henbit<br>Mouse-ear Chickweed   | 2.5 to 3.75 lbs per acre<br>(0.9 to 1.4 oz per 1,000 sq ft)   | Apply in late summer to early fall, before weeds germinate.  |

### WEED CONTROL IN AND AROUND ORNAMENTAL PLANTINGS

PRE-M® 60 DG is recommended for use on the following established field grown and container ornamentals, plants, trees, and established ground covers. Care must be taken that soil or planting mixes have settled firmly following rains, watering and that there are no cracks that would allow direct contact of PRE-M® 60 DG and roots. Directed sprays where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Sprays can be safely applied over-the-top of those plants listed below. Plant only those ornamentals listed on this label into soil treated the previous season with PRE-M® 60 DG or injury may occur. Do not apply to seedbeds. PRE-M® 60 DG may be used for the preemergence control of weeds as listed in and around the following established ornamental plants.

#### ORNAMENTAL PLANTS

| Common Name           | Scientific Name        |
|-----------------------|------------------------|
| Abeba, Glossy         | Abeba grandiflora      |
| Ash, Red              | Fraxinus grandiflora   |
| Azalea                | Fraxinus pennsylvanica |
| Barberry              | Rhododendron indicum   |
|                       | Berberis thunbergii    |
|                       | var. atropurpurea      |
| Birch, Weeping        | Betula pendula         |
| Birch, River          | Betula nigra           |
| Borwood               | Berula sempervirens    |
| Cape Jasmine          | Gardenia jasminoides   |
|                       | var. myrsinifolia      |
| Cedar, Northern white | Thuja occidentalis     |
| Cedar, Red            | Juniperus virginiana   |
| Colonaster            | Crataegus horizontalis |
| Crape Myrtle, common  | Lagerstroemia indica   |
| Dogwood, Flowering    | Cornus florida         |
| Dogwood, Sticky       | Cornus amomum          |
| Euonymus              | Euonymus alata         |
|                       | var. compacta          |
| Fig, Fraser           | Ficus benjamina        |
| Fig, White            | Ficus colorata         |
| Forsythia, Border     | Forsythia intermedia   |
| Hemlock, Eastern      | Tsuga canadensis       |
| Holly, American       | Ilex opaca             |
| Holly, Chinese        | Ilex cornuta           |
| Holly, Japanese       | Ilex crenata compacta  |
| Holly, Yaupon         | Ilex vomitoria         |
|                       | var. shillingii dwarf  |
| Juniper, Chinese      | Juniperus chinensis    |
|                       | var. pfitzeri          |
|                       | var. pfitzeri aurea    |
| Juniper, Shore        | Juniperus conferta     |
| Juniper, Trailing     | Juniperus horizontalis |
|                       | var. phumosa           |
|                       | var. prince of wales   |
|                       | var. youngstown        |
| Laurel, Mountain      | Kalmia latifolia       |
| Lilac, Common         | Syringa vulgaris       |
| Locust, Honey         | Gleditsia triacanthos  |
|                       | var. inermis           |
| Maple, Red            | Acer rubrum            |
| Mandarin              | Nandina domestica      |
| Oil, Live             | Quercus virginiana     |
| Oil, Water            | Quercus nigra          |
| Peach                 | Prunus persica         |
| Pine                  |                        |
| Pine, Bristle         |                        |

#### WEED SPECIES CONTROLLED

The following grasses and broadleaf weeds are controlled by preemergence treatments of PRE-M® 60 DG at the rates as shown under the section entitled APPLICATION RATES FOR ORNAMENTALS.

#### GRASSES CONTROLLED

| Common Name          | Scientific Name          |
|----------------------|--------------------------|
| Barnyardgrass        | Echinochloa crusgalli    |
| Bluegrass, Annual    | Poa annua                |
| Crabgrass            | Digitaria, spp           |
| Crowgrass            | Dactyloctenium aegyptium |
| Field Bindweed       | Sida lutea               |
| Field Green          | Sida verticillata        |
| Field Yellow         | Sida leucostoma          |
| Goosegrass           | Elymus indica            |
| Johnsongrass         | Sorghum halepense        |
| Johnsongrass (Proso) | Echinochloa colonum      |
| Lovegrass            | Eragrostis spp.          |
| Parrotgrass          | Panicum leucostachyum    |
| Parrotgrass, Fall    | Panicum dichotomiflorum  |
| Parrotgrass, Green   | Panicum tanarum          |
| Parrotgrass, Field   | Cynchura latifolia       |
| Signalgrass          | Brachiaria platyphylla   |
| Sprangletop, Mexican | Leptochloa univervis     |
| Sprangletop, Red     | Leptochloa billardieri   |
| Witchgrass           | Panicum capillare        |
| Woolly Cupgrass      | Cladonia villosa         |

#### BROADLEAF WEEDS CONTROLLED

| Common Name          | Scientific Name         |
|----------------------|-------------------------|
| Carpetweed           | Mollugo verticillata    |
| Chickweed, Common    | Stellaria media         |
| Chickweed, Mouse-ear | Corostem vulgatum       |
| Clover, Horse        | Trifolium pratense      |
| Cudweed              | Gratiola spp            |
| Fiddleneck           | Amaranthus retrofractus |
| Fleece               | Erodium spp             |
| Henbit               | Lamium amplexicaule     |
| Knotted              | Polygonum aviculare     |
| Kochia               | Kochia scoparia         |
| Lambquarters         | Chenopodium album       |
| Pigweed              | Amaranthus spp          |
| Puncturewort         | Tribulus terrestris     |

thoroughly after handling and before eating or drinking. Do not apply directly to crops. This product is toxic to fish. Do not apply directly to water. Drift may favor drift from target areas. Do not use in ornamentals.

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-684-53

### DIRECTION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. CHEMIGATION, Do not apply this product through any type of irrigation system.

## STORAGE AND DISPOSAL

**STORAGE** Do not contaminate water, food or feed by storage or disposal. Do not store above 120°F. Store product in its original labeled container in a cool, dry, locked place out of reach of children.  
**PESTICIDE DISPOSAL** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL** Completely empty bag into application equipment. Dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

### MIXING AND APPLICATION INSTRUCTIONS

Add LESCO PRE-M 60 DG slowly to a partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. Maintain continuous agitation while adding the LESCO PRE-M 60 DG. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed. Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. Avoid application when winds may cause spray drift.

Treatments are most effective in controlling weeds when adequate rain fall or overhead irrigation is received within 7 days after application. Avoid contact of spray solution with driveways, stone, wood or other porous surfaces. Rinse immediately to avoid staining.

### COMPATIBILITY

LESCO PRE-M 60 DG is compatible with most commonly used herbicides. When tank mixing be sure to add LESCO PRE-M 60 DG to the partially filled tank first, mix thoroughly, and then add combination products to the mixture as directed or as instructed on its EPA approved label. Do not apply tank mix combinations unless product is approved for such tank mix use and previous experience indicates the mixture is effective and will not result in application problems or plant injury. Do not attempt to mix LESCO PRE-M 60 DG directly with liquid fertilizers. Pre-mix one part of LESCO PRE-M 60 DG with two parts water and add this diluted mixture slowly into the tank while agitating.

### GENERAL INFORMATION

LESCO PRE-M 60 DG treatments will not control established weeds. Applications must be made prior to germination of weed seeds. Three applications per year may be made. Allow at least 2 months between applications except when otherwise indicated on this label. Use only on well established turfgrass with a dense and uniform stand. Do not use on newly seeded or sprigged turfgrass. Do not use on winter-overseeded turfgrass. Do not use on turfgrass severely thinned by winter stress. LESCO PRE-M 60 DG provides effective residual weed control which may affect new turfgrass seedlings or sprigs. Do not reseed or winter over-seed treated turfgrass within 4 growing months after application. Do not sprig turfgrass within 5 months after application. Do not use on Bentgrass or *Dichondra* maintained as desirable turf.

## APPLICATION RATES FOR ORNAMENTALS

For preemergence control of the weed species listed in the next column, apply PRE-M 60 DG at the following rates.

| Length of Control and Amount of PRE-M 60 DG/ Acre |                   |
|---|-------------------|
| Short Term Control (2-4 months)                   | 3.3 lbs. per Acre |
| Long Term Control (6-8 months)                    | 6.6 lbs. per Acre |

PRE-M 60 DG will not control established weeds.

The efficacy of PRE-M 60 DG will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

## FOR GROUND COVER ORNAMENTALS

PRE-M 60 DG may be used for the preemergence control of the weed species listed in and around the following established ground cover plants.

| Common Name             | Scientific Name               |
|-------------------------|-------------------------------|
| Capeweed                | <i>Arctotheca calendula</i>   |
| Daisy, Trailing African | <i>Osteospermum</i>           |
| Gazania                 | <i>Tricacosum Gazania</i>     |
| Rose-of-Sharon          | <i>splendens</i>              |
| Iceland, Large Leaf     | <i>Hypericum corymbosum</i>   |
| Ny. English             | <i>Carpobrotus edulis</i>     |
| Ny. Geranium            | <i>Hedera</i>                 |
| Myoporum                | <i>Polygonum portatum</i>     |
| Pachysandra             | <i>Myoporum parviflorum</i>   |
| Potentilla spp          | <i>Pachysandra terminalis</i> |
| Potentilla spp          | <i>Potentilla franchetii</i>  |

## PRECAUTIONS

Apply only to established plantings  
DO NOT apply PRE-M 60 DG in ornamentals nurseries

#019010

Net Weight 10 Pounds

LESCO, INC. - 2005 LAKE ROAD - ROCKY RIVER, OHIO 44116 U.S.A.

## SOUTHERN GRASSES

| Grass  | Application Rate  | Application Timing   |
|--|---|--|
| Bahia<br>Bermudagrass<br>Centipedegrass<br>St. Augustinegrass<br>Tall Fescue<br>Zoysiangrass   | 2.5 to 3.75 lbs. per acre<br>(0.9 to 1.4 oz per 1,000 sq ft)  | Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks using 2.5 lbs. per acre or 0.9 oz. / 1,000 sq ft. |
| Crabgrass<br>Fall Panicum<br>Foxtail<br>Prostrate Spurge<br>Yellow Woodsorum<br>Cuscuta<br>Crabgrass<br>Goosegrass<br>Hop Clover<br>Evening Primrose | 5.0 lbs. per acre<br>(1.8 oz per 1,000 sq ft)                 |  |
| Annual Bluegrass<br>Common Chickweed<br>Herb<br>Mouse-ear Chickweed  | 2.5 to 3.75 lbs. per acre<br>(0.9 to 1.4 oz. per 1,000 sq ft) | Apply in late summer to early fall, before weeds germinate   |

## WEED CONTROL IN AND AROUND ORNAMENTAL PLANTINGS

PRE-M 60 DG is recommended for use on the following established field grown and container ornamentals, plants, trees, and established ground covers. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of PRE-M 60 DG and roots. Directed sprays where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Sprays can be safely applied over the top of those plants listed below. Plant only those ornamentals listed on this label into soil treated the previous season with PRE-M 60 DG or injury may occur. Do not apply to seedbeds. PRE-M 60 DG may be used for the preemergence control of weeds as listed in and around the following established ornamental plants.

### ORNAMENTAL PLANTS

| Common Name           | Scientific Name  |
|-----------------------|--|
| Abeles, Glossy        | <i>Abelia grandiflora</i><br>var <i>grandiflora</i>  |
| Ash, Red              | <i>Fraxinus pennsylvanica</i>  |
| Azalea                | <i>Rhododendron indicum</i>  |
| Barberry              | <i>Berberis thunbergii</i><br>var <i>atropurpurea</i>  |
| Birch, Weeping        | <i>Betula pendula</i>  |
| Birch, River          | <i>Betula nigra</i>  |
| Bonwood               | <i>Berus sempervirens</i>  |
| Cape Jasmine          | <i>Gardenia jasminoides</i><br>var <i>myrsin</i>   |
| Cedar, Northern white | <i>Thuja occidentalis</i>  |
| Cedar, Red            | <i>Juniperus virginiana</i>  |
| Colonaster            | <i>Colonaster horizontalis</i>   |
| Crape myrtle, common  | <i>Lagerstroemia indica</i>  |
| Dogwood, Flowering    | <i>Cornus florida</i>  |
| Dogwood, Silky        | <i>Cornus amomum</i>   |
| Eucalyptus            | <i>Eucalyptus nitida</i><br>var <i>compacta</i>  |
| Fir, Fraser           | <i>Abies fraseri</i>   |
| Fir, White            | <i>Abies concolor</i>  |
| Forsythia, Border     | <i>Forsythia intermedia</i>  |
| Hemlock, Eastern      | <i>Tsuga canadensis</i>  |
| Holly, American       | <i>Ilex opaca</i> var <i>huxia</i>   |
| Holly, Chinese        | <i>Ilex cornuta</i>  |
| Holly, Japanese       | <i>Ilex crenata compacta</i>   |
| Holly, Yaupon         | <i>Ilex vomitoria</i><br>var <i>shillingii dwarf</i>   |
| Juniper, Chinese      | <i>Juniperus chinensis</i><br>var <i>plutzer</i>   |
| Juniper, Shore        | <i>Juniperus sibirica</i>  |
| Juniper, Trailing     | <i>Juniperus conferta</i>  |
|                       | <i>Juniperus horizontalis</i><br>var <i>plumosa</i><br>var <i>prince of wales</i><br>var <i>youngstown</i> |
| Laurel, Mountain      | <i>Kalmia latifolia</i>  |
| Lilac, Common         | <i>Syringa vulgaris</i>  |
| Locust, Honey         | <i>Gleditsia triacanthos</i><br>var <i>inermis</i>   |
|                       | <i>Acer rubrum</i><br><i>Nandina domestica</i><br><i>Quercus virginiana</i>                                |
| Maple, Red            | <i>Quercus nigra</i>   |
| Nandina               | <i>Prunus persica</i>  |
| Oak, Live             | <i>Philadelphus inxertus</i>   |
| Oak, Water            | <i>Philadelphus japonica</i>   |
| Peach                 | <i>Pinus nigra</i>   |
| Phlox                 | <i>Pinus laevis</i>  |
| Pieris, Japanese      | <i>Pinus strobus</i>   |
| Pine, Austrian        | <i>Ligustrum ovalatum</i>  |
| Pine, Loblolly        | <i>Photosporum tobbii</i>  |
| Pine, White           | <i>Oryedendron arboreum</i>  |
| Privet, California    | <i>Spirea vanhouttei</i>   |
| Privet, Japanese      | <i>Picea pungens glauca</i>  |
| Robin                 | <i>Myrica carolina</i>   |
| Sourwood              | <i>Taxus cuspidata</i>   |
| Spiraea               |  |
| Spruce, Colorado Blue |  |
| W. Myrtle, Common     |  |
| Yew                   |  |
| Yew, Japanese         |  |

### WEED SPECIES CONTROLLED

The following grasses and broadleaf weeds are controlled by preemergence treatments of PRE-M 60 DG at the rates as shown under the section entitled APPLICATION RATES FOR ORNAMENTALS.

### GRASSES CONTROLLED

| Common Name       | Scientific Name                  |
|-------------------|----------------------------------|
| Barnyardgrass     | <i>Echinochloa crusgalli</i>     |
| Bluegrass, Annual | <i>Joe-Annus</i>                 |
| Crabgrass         | <i>Digitaria, spp</i>            |
| Crabgrass, Fall   | <i>Jacynctocentrum aegyptium</i> |
| Crabgrass, Green  | <i>Setaria faberii</i>           |
| Crabgrass, Yellow | <i>Setaria viridis</i>           |
| Crabgrass         | <i>Setaria leucocera</i>         |
| Crabgrass         | <i>Echinochloa indica</i>        |
| Crabgrass         | <i>Rotibolus exaltata</i>        |
| Crabgrass         | <i>Sorghum halepense</i>         |
| Crabgrass         | <i>Echinochloa coliformis</i>    |
| Crabgrass         | <i>Elymus spp.</i>               |
| Crabgrass         | <i>Panicum fasciculatum</i>      |
| Crabgrass         | <i>Panicum dichotomiflorum</i>   |
| Crabgrass         | <i>Panicum teres</i>             |
| Crabgrass         | <i>Panicum lanatum</i>           |
| Crabgrass         | <i>Cynchus incertus</i>          |
| Crabgrass         | <i>Bracharia platyphylla</i>     |
| Crabgrass         | <i>Elychloa uncinata</i>         |
| Crabgrass         | <i>Lepochloa filiformis</i>      |
| Crabgrass         | <i>Panicum capillare</i>         |
| Crabgrass         | <i>Crabgrass villosa</i>         |

### BROADLEAF WEEDS CONTROLLED

| Common Name          | Scientific Name                 |
|----------------------|---------------------------------|
| Carpenterweed        | <i>Mollugo verticillata</i>     |
| Chickweed, Common    | <i>Stellaria media</i>          |
| Chickweed, Mouse-ear | <i>Cerastium vulgatum</i>       |
| Clover, Hop          | <i>Trifolium procumbens</i>     |
| Cudweed              | <i>Gnaphalium spp</i>           |
| Fiddleneck           | <i>Amsinckia intermedia</i>     |
| Filaree              | <i>Erodium spp</i>              |
| Henbit               | <i>Lamium amplexicaule</i>      |
| Knotted (prostrate)  | <i>Polygonumavicaria</i>        |
| Kochia               | <i>Kochia scoparia</i>          |
| Lambsquarters        | <i>Chenopodium album</i>        |
| Pigweed              | <i>Amaranthus spp.</i>          |
| Puncturevine         | <i>Tribulus terrestris</i>      |
| Purslane             | <i>Portulaca oleraceae</i>      |
| Purslane, Florida    | <i>Richardia scabra</i>         |
| Rocket, London       | <i>Sisymbrium spp</i>           |
| Shepherdspurse       | <i>Capsella bursa-pastoris</i>  |
| Smartweed, Penn'     | <i>Polygonum pennsylvanicum</i> |
| Spurge, Annual       | <i>Euphorbia, spp.</i>          |
| Spurge, prostrate    | <i>Euphorbia maculata</i>       |
| Wood-sorrel, yellow  | <i>Oxalis stricta</i>           |
| Velvetleaf           | <i>Abutilon theophrasti</i>     |

**WARRANTY.** LESCO, INC. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

**BEST AVAILABLE COPY**