

10404-56 10/1 Pm ac OK

# LESCO 24-3-7



## Turf Fertilizer with 1.15%

A granular fertilizer plus selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

- Bluegrass (Perennial) Fescue Bermudagrass Centipedegrass Bentgrass Fairways
- Ryegrass (Perennial) Bahiagrass St. Augustinegrass Zoysiagrass

Active ingredients  
 N-ethyl-N-ethyl d.a.e. 2,6-dinitro-p-toluene 0.71%  
 Methyl d.a.e. sulfone 2,6-dinitro-N-ethyl-p-toluene 0.38%  
 inert ingredients 98.91%

Contains 0.575 pounds active ingredients per 50 pound bag

Team™ and the diagonal color bar are trademarks for Lesco Products Company and its subsidiaries.

### GUARANTEED ANALYSIS

Total Nitrogen (N)	24.00%
0.34% Ammoniacal Nitrogen	
18.81% Urea Nitrogen	
5.25% Coated Slow Release Urea Nitrogen	
A-grade Phosphoric Acid (P2O5)	3.00%
Soluble Potash (K2O)	3.00%
Sulfur (S) - Free	2.90%

INGREDIENTS: Sulfur Coated Urea, Urea, Ammoniacal Phosphate, Muriate of Potash

2. Late spring, early summer germinating annual grasses - goosegrass  
 Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks after  
 than crabgrass, barnyard and other summer annual grass species. Application should therefore be  
 made accordingly.

A. Northern (cool season) turfgrasses  
 For preemergence control, apply 174 pounds (2 pounds active ingredients) of Lesco 24-3-7  
 with Team per acre in the late spring or early fall and again in late winter or spring one to two  
 weeks prior to the expected germination of goosegrass.

B. Southern (warm season) turfgrasses  
 For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredients) of Lesco  
 24-3-7 with Team per acre in the late spring or early summer and to two weeks prior to the  
 expected germination of goosegrass.

3. Northern and southern turfgrasses  
 In areas of the United States where goosegrass control is both early annual grasses (crabgrass,  
 foxtails and barnyard grass) and the later germinating goosegrass is essential, two applications of the  
 recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of  
 these annual grass species.

4. Poa annua (annual bluegrass)  
 In areas of heavy infestation of annual bluegrass, its elimination will normally result in the  
 turfgrass areas. Proper fertilization, irrigation and soil incorporation (including soil aeration)  
 testing the site turfgrasses and reseeded seeded turfgrasses to 1/8 to 1/4 inch then a thin  
 previously occupied by annual bluegrass.

A. Northern (cool season) turfgrasses  
 For preemergence control, apply 174 pounds (2 pounds active ingredients) of Lesco 24-3-7  
 with Team per acre in the late summer or early fall one to two weeks prior to the expected germination  
 of annual bluegrass. In southern areas of the United States where germination of annual  
 bluegrass commences for extended periods of time, a second application at 174-261 pounds per  
 acre can be made 10 to 12 weeks after the initial treatment.

For application to smaller areas, apply the following amounts of Lesco 24-3-7 with Team

APPROXIMATE LESCO ROTARY SPREADER SETTINGS

Per Acre	Approx. Lesco Spreader Setting	Per 1000 sq. ft.	Lbs. N/1000 sq. ft.
131 pounds	6	3.0 pounds	0.72
174 pounds	6 1/2	4.0 pounds	0.96
261 pounds	7 1/2	6.0 pounds	1.44

\*IMPORTANT: These settings are only approximate. Age, condition of spreader and operator speed  
 can cause wide variation. Be sure to calibrate your spreader with each application.  
 Preseedling  
 Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass  
 seedlings. Preseedling should be delayed for at least 6 weeks after applying Lesco 24-3-7 with Team.  
 When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and  
 fertilization be followed. For satisfactory reseeding results after using Lesco 24-3-7 with Team the  
 amount of seed used should be increased and equipment designed to place seed in contact with the  
 soil should be employed. When using at the highest recommended rate, reseeding should be delayed 12  
 to 18 weeks after application.

Application Equipment  
 Apply Lesco 24-3-7 with Team evenly over the turfgrass area without gaps or overlap. A more uniform  
 application can be made by spreading just the required amount over the area and then applying the  
 remaining half at right angles. Both disc type and rotary type spreaders may be used. Avoid the use of  
 spreaders which would apply the material in narrow concentrated bands. For best results use  
 equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use  
 according to the manufacturer's directions. Check frequently to be sure equipment is working properly  
 and distributing granules uniformly.

Helpful Application Techniques For Applying Lesco 24-3-7 with Team  
 1. When using a bag type spreader a splash board mounted under the hopper will help dispense  
 granules more evenly.  
 A chain fastened to the side of your spreader will have a marked or scuffed felt tire to keep and  
 help prevent missed areas.  
 2. When using a disc mounted or pull-type rotary applicator it is suggested that a seed application  
 ring and a cone should be used to control distribution of granules.  
 Early morning applications when the wind is calm and dew is on the ground are ideal. When tracks  
 are dry to follow and granules can be spread uniformly.  
 3. Small area application is best using a small rotary applicator. To insure proper coverage and avoid  
 missed areas, early morning treatment when wheel tracks are evenest in the dew and the wind is  
 calm are ideal.

Precautions  
 Apply Lesco 24-3-7 with Team to the turf prior to the germination of weed grasses since it will not  
 control established weeds. It may also establish 1/2 to 1/4 inch (Poa annua) and  
 broadleaf weeds at rates above 131 pounds per acre (1.5 pounds active ingredients). This should not be  
 applied in the spring to turfgrass planted in previous fall or bermudagrass where they have been  
 overseeded with winter grasses. A spring application may kill the over seeded grasses. Because of  
 varying cultural practices, do not use on bermudagrass, St. Augustinegrass or Centipedegrass until  
 these turfs are well established.

WARRANTY  
 LESCO warrants that the herbicide conforms to the label directions. It does not warrant that it will be  
 successful with every application under normal conditions. This herbicide is a pre-emergent and is  
 intended to prevent. Since timing, method of application, weather, soil pH, soil moisture, soil fertility  
 and other factors affect the use of this product, the user should refer to the label for complete  
 directions and other factors affecting the use of this product. The user assumes all risks of use of this  
 product which are not stated to be covered by the user's warranty.

### WARNING: Keep Out of Reach of Children

**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**  
**WARNING:** Causes substantial but temporary eye injury. Do not get in eyes or on  
 clothing, face, goggles, face shield or safety glasses. Wash thoroughly with soap and water  
 after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust  
 and contact with skin. Harmful if swallowed, inhaled or absorbed through the skin.  
**STATEMENT OF PRACTICAL TREATMENT**  
 If in eyes: Call a physician or Poison Control Center. Flush eyelids open and flush with a  
 steady gentle stream of water for 15 minutes.  
 If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity  
 of milk, egg white, gelatin solution, or if these are not available, large quantities of water.  
 Avoid alcohol.  
 If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.  
 If inhaled: If breathing discomfort occurs, move individual to fresh air. Get medical atten-  
 tion if breathing difficulty occurs.  
**Environmental Hazards**  
 Do not contaminate any body of water by direct application, cleaning of equipment, or disposal  
 of wastes.

**Directions for Use - Read all Directions Carefully Before Applying**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Lesco  
 24-3-7 with Team is a preemergence herbicide which is applied to established turfgrasses for the  
 control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass (hairy crabgrass  
 a crested barnyardgrass (travertgrass) and green and yellow foxtail. It controls these annual weeds  
 by killing their seeds as they germinate. Apply this product before these annual weeds begin to  
 germinate since it does not control established weeds. This product will not harm most nearby  
 established ornamental trees and shrubs.

**STORAGE AND DISPOSAL**  
**Storage:** Store in original container only. In case of spill, contain material and dispose of  
 wastes.  
**Pesticide Disposal:** Do not contaminate water, food, or feed by storage and disposal.  
 Wastes resulting from the use of this product may be disposed of on site or at an approved  
 waste disposal facility.  
**Container Disposal:** Completely empty bag into application equipment. Then dispose  
 of empty bag in a sanitary landfill or by incineration, or if allowed by State and local  
 authorities, by burning. If burned, stay out of smoke.

**APPLICATION DIRECTIONS**  
**Rates, Frequency and Timing of Application**  
 Successful preemergence control of the annual grass weeds listed on the label requires proper timing  
 of application.  
 Apply Lesco 24-3-7 Turf Fertilizer with Team approximately one to two weeks prior to the germination  
 of annual grass weeds. All summer annual grasses do not germinate at the same time. If application  
 timing does not coincide with the normal germination period of any of the annual grass weeds listed on  
 the label, weed control results may be erratic or poor. Optimum weed control performance will be  
 obtained if treated areas are irrigated soon after application.  
 Any culture practices that disturb the soil, such as aerification or verticillating, should be done prior to  
 application.

1. Spring germinating summer annual grasses - crabgrasses, foxtails and  
 barnyardgrasses.
  - A. Northern (cool season) turfgrasses  
 For preemergence control, apply 131 to 174 pounds per acre (1.5 to 2 pounds) active  
 ingredients of Lesco 24-3-7 with Team per acre in the spring one to two weeks prior to the  
 onset of continuous favorable for annual weed grass seed germination. This will apply 0.72 to  
 0.96 pounds of nitrogen per 1000 square feet.
  - B. Southern (warm season) turfgrasses  
 For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredients) of Lesco  
 24-3-7 Turf Fertilizer with Team per acre in the late winter or early spring and to two weeks  
 prior to the onset of continuous favorable for annual weed grass seed germination.  
 In areas of the United States where germination of the annual weed grass seed species continues for  
 extended periods of time, a second application at the recommended rate may be made 8 to 10 weeks  
 after initial treatment for continued control.

10404-56

EPA Reg No 10404-53  
EPA Est No 10404 NJ 01

#XXXXXX

Manufactured by  
LESCO, INC.  
30005 Lake Road, P.O. Box 16915  
Rocky River, Ohio 44116

Net Wgt 50 lbs

2460810W