

PM 25 10404-56
1/2

LESCO 24-3-7

Turf Fertilizer with 1.15%



A granular fertilizer plus selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

- Bluegrass (Perennial)
- Fescue
- Bermudagrass
- Centipedegrass
- Bentgrass Fairways
- Ryegrass (Perennial)
- Bahia grass
- St. Augustinegrass
- Zoysiagrass

Active Ingredients:
 N-butyl-N-dimethyl-3,4,5-trifluorophthalimide 0.77%
 Inert Ingredients: 99.23%
 Contains 8.575 pounds active ingredient per 56 pound bag

"Team" and the diagonal color bar are trademarks for Lesco Products herein plus Infratran

GUARANTEED ANALYSIS

Total Nitrogen (N) 24.60%
 0.63% Ammoniacal Nitrogen
 23.97% Urea Nitrogen
 Available Phosphoric Acid (P₂O₅) 3.80%
 Soluble Potash (K₂O) 7.50%
 INGREDIENTS: Sulfur Coated Urea, Urea, Ammonium Phosphate, Muriate of Potash

Keep Out of Reach of Children
WARNING

Precautionary Statements Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust and contact with skin. Harmful if swallowed, inhaled or absorbed through the skin.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Call a physician or Poison Control Center. Hold eyelids open and flush with a steady gentle stream of water for 15 minutes.
 If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.
 If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
 If inhaled: If breathing discomfort occurs, move individual to fresh air. Get medical attention if breathing difficulty occurs.

Environmental Hazards

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes.

Directions for Use - Read all Directions Carefully Before Applying

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Lesco 24-3-7 with Team is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds. This product will not harm most nearby established ornamental trees and shrubs.
 Caution: Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of spill, contain material and dispose as wastes.
Pesticide Disposal: Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application
 Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply Lesco 24-3-7 Turf Fertilizer with Team approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.
 Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrasses, foxtails and barnyardgrass.
 - A. Northern (cool season) turfgrasses.
 For preemergence control, apply 174 pounds per acre (1.5 to 2 pounds active ingredient) of Lesco 24-3-7 with Team per acre in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination. This will apply 0.45 to 0.86 pounds of nitrogen per 1000 square feet.
 - B. Southern (warm season) turfgrasses.
 For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredient) of Lesco 24-3-7 Turf Fertilizer with Team per acre in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass seed species continues for extended periods of time, a second application at the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

2. Late spring, early summer germinating annual grasses - goosegrass.
 Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should therefore, be timed accordingly.
 - A. Northern (cool season) turfgrasses.
 For preemergence control, apply 174 pounds (2 pounds active ingredient) of Lesco 24-3-7 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.
 - B. Southern (warm season) turfgrasses.
 For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredient) of Lesco 24-3-7 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and southern turfgrasses.
 In areas of the United States where preemergence control of both early annual grasses (crabgrass, foxtails and barnyardgrass) and the later germinating goosegrass is desired, two applications at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4. Poa annua (annual bluegrass).
 In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.
 - A. Northern (cool season) turfgrasses.
 For preemergence control, apply 174 pounds (2 pounds active ingredient) of Lesco 24-3-7 with Team per acre in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.
 - B. Southern (warm season) turfgrasses.
 For preemergence control, apply 261 pounds (3 pounds active ingredient) of Lesco 24-3-7 with Team per acre in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at 174-261 pounds per acre can be made 10 to 12 weeks after the initial treatment.

For application to smaller areas, apply the following amounts of Lesco 24-3-7 with Team.

Per Acre	Approx. Lesco Spreader Setting	Per 1000 sq ft	Lbs. N/1000 sq ft
131 pounds	G	3.0 pounds	0.72
174 pounds	G 1/4	4.0 pounds	0.96
261 pounds	H 1/4	6.0 pounds	1.44

IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Lesco 24-3-7 with Team. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Lesco 24-3-7 with Team, the amount of seed used should be increased and equipment designed to place seed in contact with the soil should be employed. When using at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

Application Equipment

Apply Lesco 24-3-7 with Team evenly over the turfgrass area without stops or misses. A more uniform application can be made by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. For best results use equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

Helpful Application Techniques For Applying Lesco 24-3-7 with Team

1. When using a drop type spreader, a splash board mounted under the hopper will help dispense granules more evenly.
 A chain fastened to the side of your spreader will leave a marked or scuffed turf area to follow and help prevent missed areas.
2. When using tractor mounted or pull-type rotary applicators it is suggested that a seed application ring and a line should be used to control distribution of granules.
 Early morning applications when the wind is calm and dew is on the ground are ideal. Wheel tracks are easy to follow and granules can be spread uniformly.
3. Small area application is easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm are ideal.

Precautions

Apply Lesco 24-3-7 with Team to the turf prior to the germination of weed grasses since it will not control established weeds. It may then establish annual bluegrass (Poa annua) turfs for fine-leaved species of rates above 131 pounds per acre (1.5 pounds active ingredient). This should not be applied in the spring to lawns as planned the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the over-seeded grasses. Because of varying cultural practices, do not use on bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

WARRANTY

LESCO warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

EPA Reg. No. 10404-56
 EPA Est. No. 10404-04-02

#015376

Manufactured by
LESCO, INC.
 20005 Lake Road, P.O. Box 16915
 Rocky River, Ohio 44116

Net Wgt 50 Lbs.
 3670108W

Rev. 12/87

ACCEPTED

SEP - 1 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 10404-56

2

20/2