

LESCO 26-3-7

10/1
PM 25
10/25/77

Team

Turf Fertilizer with 1.25%

A granular fertilizer plus selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

- Bluegrass (Perennial) Fescue Bermudagrass Centipedegrass Bantgrass Fairways
- Ryegrass (Perennial) Bahiagrass St. Augustinegrass Zoysiagrass

Active ingredients

Methyl N-ethyl-2,4,6-trinitro-p-sulfonate	88.4%
Influracil N, 2-(4-chlorophenyl)-6-dimethyl-p-sulfonate	0.11%
Other ingredients	98.75%

Contains 0.625 pounds active ingredient per 50 pound bag
"Team" and the diagonal color bar are trademarks for Lesco Products beneath plus Influracil

GUARANTEED ANALYSIS

Total Nitrogen (N)	26.00%
0.64% Ammoniacal Nitrogen	
19.36% Urea Nitrogen	
8.00% Coated Slow Release Urea Nitrogen	
Avail. Phosphoric Acid (P ₂ O ₅)	3.00%
Soluble Potash (K ₂ O)	7.00%
Sulfur (S) (Free)	3.30%
Sulfur (S) (Combined)	2.50%

INGREDIENTS: Sulfur Coated Urea, Urea, Ammonium Phosphate, Sulfate of Potash

2. Late spring, early summer germinating annual grasses - goosegrass. Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass. In late spring and early summer annual grass species. Applications should therefore be timed accordingly.

A. Northern cool season turfgrasses
For preemergence control, apply 174 pounds (2 pounds active ingredient) of Lesco 26-3-7 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

B. Southern warm season turfgrasses
For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredient) of Lesco 26-3-7 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and southern turfgrasses
In areas of the United States where preemergence control of both early annual grasses (crabgrass, furbish and bermudagrass) and the later germinating goosegrass is desired, two applications at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4. Poa annua (annual bluegrass)
In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in the turfgrass areas. Proper fertilization, irrigation and soil incorporated seeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A. Northern cool season turfgrasses
For preemergence control, apply 174 pounds (2 pounds active ingredient) of Lesco 26-3-7 with Team per acre in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass.

B. Southern warm season turfgrasses
For preemergence control, apply 261 pounds (3 pounds active ingredient) of Lesco 26-3-7 with Team per acre in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at 174-261 pounds per acre can be made 10 to 12 weeks after the initial treatment.
For application to smaller areas, apply the following amounts of Lesco 26-3-7 with Team:

APPROXIMATE LESCO ROTARY SPREADER SETTINGS

Per Acre	Approx. Lesco Spreader Setting	Per 1000 sq. ft.	Lbs. Per 1000 sq. ft.
120 pounds	7 1/2	275 pounds	0.72
160 pounds	9 1/2	367 pounds	0.96
240 pounds	14 1/2	550 pounds	1.43

*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

Reseeding
Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Lesco 26-3-7 with Team. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Lesco 26-3-7 with Team, the amount of seed used should be increased and equipment designed to place seed in contact with the soil should be employed. When using at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

Application Equipment
Apply Lesco 26-3-7 with Team to the turf prior to the germination of weed grasses since it will not control established weeds. It may then be applied to established annual bluegrass (Poa annua) turfs and treated lawns at rates above 131 pounds per acre (1.5 pounds active ingredient). This should not be applied in the spring to turfgrasses planted the previous fall in bermudagrass areas that have been overseeded with winter grasses, a spring application may then be made to control grasses. Because of varying cultural practices, do not use on bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

Helpful Application Techniques For Applying Lesco 26-3-7 with Team
1. When using a drop type spreader, a splash board mounted under the hopper will help disperse granules more evenly.
A chain fastened to the side of your spreader will leave a marked or scuffed turf area to follow and help prevent missed areas.

2. When using tractor mounted or pull-type rotary applicators it is suggested that a seed application ring and a time wheel be used to control distribution of granules.
Early morning applications, when the wind is calm and dew is on the ground are ideal. Wheel tracks are easy to follow and granules can be spread uniformly.

3. Small area applications are easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm are ideal.

Precautions
Apply Lesco 26-3-7 with Team to the turf prior to the germination of weed grasses since it will not control established weeds. It may then be applied to established annual bluegrass (Poa annua) turfs and treated lawns at rates above 150 pounds per acre (1.5 pounds active ingredient). This should not be applied in the spring to turfgrasses planted the previous fall in bermudagrass areas that have been overseeded with winter grasses, a spring application may then be made to control grasses. Because of varying cultural practices, do not use on bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

WARRANTY
LESCO warrants that the herbicide conforms to the Chemical Literature on its label. When used in accordance with label directions under normal conditions, the herbicide is warranted to be its intended purpose. Since timing, method of application, soil factors, plant and soil conditions, and other factors, which may affect the use of this product are beyond our control, we warrant a given quantity of this product will control weeds under conditions which are normal and reasonably foreseeable. It is not a guarantee of results of any kind.

WARNING: Keep Out of Reach of Children

Precautionary Statements Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust and contact with skin. Harmful if swallowed, inhaled or absorbed through the skin.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Call a physician or Poison Control Center. Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes.
If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gastric solution, or if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled: If breathing discomfort occurs, move individual to fresh air. Get medical attention if breathing difficulty occurs.

Environmental Hazards
Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes.

Directions for Use - Read all Directions Carefully Before Applying

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Lesco 26-3-7 with Team is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (Poa annua), smooth and hairy crabgrass, goosegrass (Lolium crabgrass), crowfoot, hairytopgrass (watergrass) and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds. This product will not harm most nearby established ornamental trees and shrubs.

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, if allowed by State and local authorities. By burning, if burned, stay out of smoke.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application
Successful preemergence control of the annual grass weeds listed on the label requires proper timing of application.

Apply Lesco 26-3-7 Turf Fertilizer with Team approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticuring, should be done prior to application.

1. Spring germinating summer annual grasses - crabgrass, foxtails and bermudagrass.

A. Northern cool season turfgrasses
For preemergence control, apply 150 to 200 pounds per acre (1.5 to 2 pounds active ingredient) of Lesco 26-3-7 with Team per acre in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination. This will apply 0.72 to 0.96 pounds of nitrogen per 1000 square feet.

B. Southern warm season turfgrasses
For preemergence control, apply 174 to 261 pounds (2 to 3 pounds active ingredient) of Lesco 26-3-7 Turf Fertilizer with Team per acre in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass seed species continues for extended periods of time, a second application at the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

ACCEPTED
DEC 23 1986
EPA Reg. No. 10404-57

EPA Reg No 10404-53
EPA Est No 10404 NJ-01
#XXXXXX

Manufactured by
LESCO, INC.
20005 Lake Road, P.O. Box 16915
Rocky River, Ohio 44116

Net Wt 50 lbs

12/0906W