

PN 25 10404-27

# LESCO 30-6-13

## Turf Fertilizer with 1.25% Team

ACCEPTE  
SEP - 1 1988

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 10404-57

A granular fertilizer plus selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bluegrass (Perennial)	Fescue	Bermudagrass	Centipedegrass	Bentgrass Fairways
Ryegrass (Perennial)	Bahiagrass	St. Augustinegrass	Zoysiagrass	

Active ingredients:	
N-butyl-N-ethyl-4,4'-dinitro-p-toluene	0.04%
Inhalin: 4,4'-thiobis-2,6-dinitro-N,N-diethyl-p-toluene	0.41%
Inert ingredients:	99.75%

Contains 0.625 pounds active ingredient per 50 pound bag.

"Team" and the diagonal cut are trademarks for Elanco Products benefit plus Inhalin.

### GUARANTEED ANALYSIS

Total Nitrogen (%)	30.00%
12.75% Ammonium Nitrogen	
24.50% Urea Nitrogen	
4.37% Water Insoluble Nitrogen (WIN)	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	5.00%
Soluble Potash (K <sub>2</sub> O)	13.00%

INGREDIENTS: Urea, Ammonium Phosphate, Monocal Phosphate, Monocal Potash, Urea Formaldehyde.

Benefit!

### Keep Out of Reach of Children

### WARNING

#### Precautionary Statements Hazards to Humans and Domestic Animals

**WARNING:** Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust and contact with skin. Harmful if swallowed. Inhalation absorbed through the skin.

#### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Call a physician or Poison Control Center. Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes.

If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, protein solution, or, if these are not available, large quantities of water. Avoid alcohol.

If skin irritation: Wash with plenty of soap and water. Get medical attention if irritation persists.

If inhaled: If breathing discomfort occurs, move individual to fresh air. Get medical attention if breathing difficulty occurs.

#### Environmental Hazards

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes.

**Directions for Use - Read all Directions Carefully Before Applying**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Lesco 30-6-13 with Team is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (*Eleusine crusgalli* or *crautford*), bermuda grass (*Agrostis*) and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate. Apply the product before these annual weeds begin to germinate since it does not control established weeds. This product will not harm most nearby established ornamental trees and shrubs.  
Chlorothalonil: Do not apply this product through any type of irrigation system.

#### STORAGE AND DISPOSAL

**Storage:** Store in original container only. In case of spill, contain material and dispose as wastes.  
**Pesticide Disposal:** Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### APPLICATION DIRECTIONS

**Rate, Frequency and Timing of Application**  
Selective preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply Lesco 30-6-13 Turf Fertilizer with Team approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application.

**1. Spring germinating summer annual grasses - crabgrasses, foxtails and bermudagrass.**

A. Northern (cool season) turfgrasses

For preemergence control, apply 125 to 160 pounds per acre (1.5 to 2 pounds active ingredient) of Lesco 30-6-13 with Team per acre in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B. Southern (warm season) turfgrasses

For preemergence control, apply 160 to 240 pounds (2 to 3 pounds active ingredient) of Lesco 30-6-13 Turf

Fertilizer with Team per acre in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass seed species continues for extended periods of time, a second application of the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

**2. Late spring, early summer germinating annual grasses - goosegrass**

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails and other summer annual grass species. Applications should therefore be timed accordingly.

A. Northern (cool season) turfgrasses

For preemergence control, apply 160 pounds (2 pounds active ingredient) of Lesco 30-6-13 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

B. Southern (warm season) turfgrasses

For preemergence control, apply 160 to 240 pounds (2 to 3 pounds active ingredient) of Lesco 30-6-13 with Team per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

**3. Northern and southern bermudagrasses**

In areas of the United States where preemergence control of both early annual grasses (crabgrass, foxtails and bermudagrass) and the later germinating goosegrass is desired, two applications at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

**4. Poor germination (annual bluegrass)**

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporation reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A. Northern (cool season) turfgrasses

For preemergence control, apply 160 pounds (2 pounds active ingredient) of Lesco 30-6-13 with Team per acre in late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

B. Southern (warm season) turfgrasses

For preemergence control, apply 240 pounds (3 pounds active ingredient) of Lesco 30-6-13 with Team per acre in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at 160-240 pounds per acre can be made 10 to 12 weeks after the initial treatment.

#### Reseeding

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying Lesco 30-6-13 with Team. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Lesco 30-6-13 with Team, the amount of seed used should be increased and equipment designed to place seed in contact with the soil should be employed. When using at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

#### Applicator Equipment

Apply Lesco 30-6-13 with Team evenly over the turfgrass area without skips or misses. A more uniform application can be made by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. For best results use equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

#### Helpful Application Techniques For Applying Lesco 30-6-13 with Team

1. When using a drop type spreader, a splash board mounted under the hopper will help disperse granules more evenly.

A chain fastened to the side of your spreader will leave a marked or scuffed turf area to follow and help prevent missed areas.

2. When using tractor mounted or pull-type rotary applicators it is suggested that a seed application ring and a bungee should be used to control distribution of granules.

Early morning applications when the wind is calm and dew is on the ground are ideal. Wheel tracks are less likely to follow and granules can be spread uniformly.

3. Small area application is easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm are ideal.

#### Precautions

Apply Lesco 30-6-13 with Team to the turf prior to the germination of weeds; since it will not control established weeds. It may thin established annual bluegrass (*Poa annua*) tufts and fine-leaved fescues at rates above 120 pounds per acre (1.5 pounds active ingredient). This should not be applied in the spring in bermudagrass areas that have been overseeded with annual grasses. A spring application may thin the over-seeded grasses. Because of varying cultural practices, do not use on bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established.

#### WARRANTY

LESCO warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since growing, method of application, weather, plant and soil conditions, measure with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty of given conditions, except the use of the product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

E.P.A. Reg. No. 10404-87  
E.P.A. Ext. No. 10404-OH-02

#016653

Net Wt. 50 Lbs.  
3880203W

1061