

10404-57

9-7-2001

1/2

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 10404-57	2. EPA Product Manager Deborah McCall	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) LESCO Turf Fertilizer with 1.25% Team	PM#	
5. Name and Address of Applicant (Include ZIP Code) LESCO, Inc. 15885 Sprague Rd. Strongsville, Ohio 44136-1799 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

NOTIFICATION

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	SEP 7 - 2001
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Combination of Redundant Labeling Statements per PR Notice 98-10.
This notification is consistent with the provision of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Bernadine Javorek	Title Manager, Regulatory Affairs	Telephone No. (Include Area Code) 440-783-9250
2. Signature <i>Bernadine Javorek</i>		8. Date Application Received (Stamped)
3. Title Manager, Regulatory Affairs		
4. Typed Name Bernadine Javorek	5. Date 8/29/01	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		

Total 100.00%
Contains 0.625 pounds active ingredient per 50 pound bag.

X-X-X

GUARANTEED ANALYSIS

Total Nitrogen.....xx.xx%
 xx.xx% Ammoniacal Nitrogen
 xx.xx% Urea Nitrogen
 xx.xx% Water Insoluble Nitrogen (WIN)
 Available Phosphate (P₂O₅).....xx.xx%
 Soluble Potash (K₂O).....xx.xx%
 DERIVED FROM: Urea, Ammonium Phosphate, Muriate of Potash, Urea Formaldehyde.
 Chlorine (Cl) Max.xx.xx%
 F1406 Made in U.S.A.
 Wash or brush off walks and painted surfaces following application to avoid discoloration.

KEEP OUT OF REACH OF CHILDREN WARNING

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Do not induce vomiting. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Call a physician or Poison Control Center. Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes. This pesticide is extremely toxic to freshwater marine and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING.

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of spill, contain material and dispose as waste.
Pesticide Disposal: Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INSTRUCTIONS AND INFORMATION

CHEMIGATION: Do not apply this product through any type of irrigation system. LESCO Fertilizer with Team™ is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass, (silver crabgrass or crowfoot), barnyardgrass (watergrass), and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds. This product will not harm most nearby established ornamental trees and shrubs.

For application to smaller areas, apply the following amounts:

APPROXIMATE ROTARY SPREADER SETTINGS for LESCO Rotary Spreader*		LESCO Calibration	
Pounds Product	Pounds Product	#A/Acre	Course
4,000 sq ft			
2.8	120	1.6	#13
3.7	160	2.0	#16
5.5	240	3.0	#17

*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

Reseeding: Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applying. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation, and fertilization be followed. For satisfactory reseeding results after using product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil should be employed. When using at the highest recommended rate, reseeding should be delayed 12 to 16 weeks after application.

- When using tractor mounted or pull-type rotary applicators, it is suggested that a seed application rig and a lime shield be used to control distribution of granules. Early morning applications when the wind is calm and dew is on the ground are ideal. Wheel tracks are easy to follow and granules can be spread uniformly.
- Small area application is easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm is ideal.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter (except those involved in the watering) until the watering-in is complete and the surface is dry.

PRECAUTIONS: Apply to the turf prior to the germination of weed grasses since it will not control established weeds. It may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rate above 120 pounds per acre (1.5 pounds active ingredient). This should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application may thin the over-seeded grasses. Because of varying-cultural practices, do not use on bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established.

APPLICATION DIRECTIONS

Rate, Frequency, and Timing of Application - Successful, preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with any of the normal germination periods of any of the annual grass weeds listed on this label, weed control results may be erratic, or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

1. Spring germinating summer annual grasses - crabgrass, foxtails, and barnyardgrass

a. Northern (cool season) turfgrasses

For preemergence control, apply 120 to 160 pounds per acre (1.5 to 2 pounds active ingredient) in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination. This will apply 0.30 to 0.40 pounds of Nitrogen per 1,000 sq ft

b. Southern (warm season) turfgrasses

For preemergence control, apply 160 to 240 pounds (2 to 3 pounds active ingredient) per acre in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass seed species continues for extended periods of time, a second application at the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

2. Late spring, early summer germinating annual grasses - goosegrass

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than crabgrass, foxtails, and other summer annual grass species. Applications should, therefore, be timed accordingly.

a. Northern (cool season) turfgrasses

For preemergence control, apply 160 pounds (2 pounds active ingredient) per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

b. Southern (warm season) turfgrasses

For preemergence control, apply 160 to 240 pounds (2 to 3 pounds active ingredient) per acre in the late spring or early summer one to two weeks prior to the expected germination of goosegrass.

3. Northern and Southern Turfgrasses

In areas of the U.S. where preemergence control of both early annual grasses (crabgrass, foxtails, and barnyardgrass) and later germinating goosegrass is desired, two applications at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

4. *Poa annua* (annual bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seeding turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

a. Northern (cool season) turfgrasses

For preemergence control, apply 160 pounds (2 pounds active ingredient) per acre in the late summer or early fall and again in late winter or spring one to two weeks prior to the expected germination of annual bluegrass.

b. Southern (warm season) turfgrasses

For preemergence control, apply 240 pounds (3 pounds active ingredient) per acre in the late summer or early fall one to two weeks prior to the expected germination of annual bluegrass. In southern areas of the U.S. where germination of annual bluegrass continues for extended periods of time, a second application at 174-261 pounds per acre can be made 10 to 12 weeks after the initial treatment.

WARRANTY

LESCO warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label direction under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label direction or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

*Team is a trademark of DowAgrosciences L.L.C.

LESCO is a registered trademark of LESCO, Inc.

G:\REGUL\WP\STRONGSV\MstrLbl\Team1.25.doc Rev. 08/27/01 WPB (090188) (93-3 080393) (93-7 040194)

Net Weight: 50 lb (22.7 kg)

NOTIFICATION

SEP 7 - 2001

EPA Reg. No. 10404-57
EPA Est. No. 10404-OH-04

F1406
Manufactured by: LESCO, Inc.
15885 Sprague Rd
Strongsville, OH 44136-1799

758073xxxxx

#XXXXXX