# LESCO Granular Turf Fungicide

**Contains 1.0% Bayleton® Fungicide - For Systemic Control of Certain Diseases on Turf**

## Systemic Fungicide Granules

**FOR CONTROL OF LAWN DISEASE ON BLUEGRASS, FESCUE, RYEGRASS, BERMUDA GRASS, ZOYSIA-GRASS, ST. AUGUSTINE GRASS AND BENTGRASS**

**FOR THE PREVENTION OR CURE OF DOLLAR SPOT, COPPER SPOT, POWDERY MILDEW, RED THREAD, RUSTS, BROWN PATCH, ANTHRACNOSE, POA ANNUA DECLINE, STRIPED SMUT, FUSARIUM BLIGHT, GRAY SNOW MOLD, PINK SNOW MOLD, FUSARIUM PATCH AND SOUTHERN BLIGHT**

ACTIVE INGREDIENT:
1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone..... 1.0%
INERT INGREDIENTS:..... 99.0%
Total 100.0%

Contains BAYLETON® Turf Fungicide a Registered Trade Mark of Mobay Corp.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

**STATEMENTS OF PRACTICAL TREATMENT**

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Give water until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush eyes with plenty of water. Get medical attention.
IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention. Wash clothing before re-use.
IF INHALED: Remove victim to fresh air. Administer artificial respiration if indicated.
SEE ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 10404-UEM EPA EST NO. 10404-OH-1

NET WEIGHT 40 POUNDS

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor, or dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use.

### ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. Do not use on crops grown for food or forage. Do not apply directly to water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from target area. Do not use clippings or grass for animal feed.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store product in original bag in a cool, dry, locked place out of reach of children.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.
HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

## APPLICATION

Bayleton® is absorbed rapidly and moves systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore in order to achieve maximum control, turf should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Apply Preventive Rates prior to appearance of disease and/or when damage from disease is anticipated. Apply Curative Rate when disease is first observed.
NOTE: Apply the specified amount of LESCO Granular Turf Fungicide with a standard granular applicator calibrated to uniformly apply recommended amount of product per 1,000 Sq.Ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of 1/4 inch of water has been applied to the treated area.
WARRANTY. LESCO warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, it is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, & other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not foreseeable. The user assumes all risks of use.

## LESCO Granular Turf Fungicide
## Recommended Applications - Turfgrass Disease Control

| DISEASE | HOW MUCH TO USE PER 1,000 SQ.FT. PREVENTIVE | CURATIVE | REMARKS |
|---|---|---|---|
| DOLLAR SPOT | 1 1/2 lbs. | - | PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of product may extend for as long as 60 days dependent upon environmental conditions. |
| | 3/4 lb | - | PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of product may extend for as long as 30 days dependent upon environmental conditions. |
| | - | 3 lbs. | CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate. |
| COPPER SPOT<br>POWDERY MILDEW<br>RED THREAD (Laetisaria fuciformis)<br>RUSTS<br>BROWN PATCH (Rhizoctonia solani) (suppression) | 1 1/2 lbs. | 3 lbs | PREVENTIVE RATE: Apply at 18 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.<br>CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate. |
| ANTHRACNOSE and POA ANNUA DECLINE | 3 lbs. | - | PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 40 days. |
| | 1 1/2 lbs. | - | PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days. |
| | - | 3 lbs. | CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on preventive schedule and rate. |
| STRIPE SMUT | PREVENTIVE RATE 3 lbs. | | Make a single application in spring as grass growth begins. |
| FUSARIUM BLIGHT (Fusarium culmorum) (Fusarium poae) | (Except California) 3 lbs. | | Apply 1st applications in mid-June or 30 days prior to time this blight normally becomes evident. Apply at 30 day intervals. For California Only apply at 3 lbs. rate on 15 day intervals or at the 6 lb. rate on 30 day intervals. |
| SUMMER PATCH (Phialophora graminicola) | (California Only) 3 to 6 lbs. | | |
| GRAY SNOW MOLD (Typhula incarnata) (Except California) | 6 lbs. | | Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant. |
| PINK SNOW MOLD<br>FUSARIUM PATCH (Fusarium nivale)<br>SOUTHERN BLIGHT (Sclerotium rolfsii) (Except California) | 3 to 6 lbs. | | Apply before conditions favorable for infection occur. Repeat as needed at 60 to 90 day intervals. Do not apply over snow cover or when turf is dormant. Use high rate in turf with a history of severe yearly disease damage. |
| GRAY SNOW MOLD<br>PINK SNOW MOLD (California Only) | SINGLE APPLICATION 7.5-12 lbs/1,000sq ft. | | Apply in late September or early October. Do not apply over snow. Use the high rate in areas with a history of severe snowmold damage. |

### LESCO ROTARY SPREADER SETTINGS

| Product/1/1000 Sq.Ft. | Effective Spreader Width | 3rd Hole | Approx. Setting |
|---|---|---|---|
| .5 lbs. | 6 Feet | OPEN | C 1/4 |
| 3 lbs. | 6 Feet | OPEN | F |
| 6 lbs. | 6 Feet | OPEN | G 1/2 |

Spreader settings are only approximate. Age and conditions of spreader may allow wide variations. Calibrate your spreader by placing 1.5 lbs to cover [illegible] and [illegible] "F". If spreader contains granules at end of test; [illegible] before the end of the test; close or reduce the setting [illegible] spreader Manufacturer's Operational Manual. For [illegible] use setting "F". For 12 lbs. use setting "J". 04-05-88.

LESCO, INC.
20005 LAKE ROAD - ROCKY RIVER, OHIO 441[illegible]

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10404-58

BEST AVAILABLE COPY