UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

OCT 3 0 1990

H.R. McClane 7210 Red Road Suite 212 Miami, FL 33143

Dear Mr. McLane:

Subject: LESCO TFC DISPERSIBLE GRANULE TURF HERBICIDE

EPA Reg. No. 10404-59

Re: Amendments

Your application dated August 29, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, Rodenticide Act, as amended, is acceptable subject to comments listed below. A stamped copy is enclosed for your records. Please submit five (5) of your final printed labels before distributing your product.

- "Environmental Hazard" statement must be revised to read 1. "Do not contaminate water when disposing of equipment washwaters". Delete old statement, "Do not contaminate water by cleaning of equipment or when disposing of equipment washwaters".
- The "Storage and Disposal" block must be placed in the 2. "Directions for Use" section. It should be placed immediately before specific directions for use or at the end of "Directions for use".

Sincerely,

Robert J. Taylor Product Manager (25) Fungicide-Herbicide Branch Registration Division (H7505C)

enclosure

CONCURRENCES		
SYMBOL H7505C		
SURNAME COLE		
DATE 10/30/90		
EPA Farm 1320-1 (12-70)	*U.S.GPO:1989-624-485/19186	OFFICIAL FILE COPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANNALS

CAUTION: MAY IRRITATE EYEB, NOSE, THROAT, AND SIGN. Avoid breathing dust or spray mist. Avoid contact with sign, eyes, and clothing. In case of contact with eyes immediately flush with plenty of water. Get medical attention if inflation preside. Wash thoroughly after handling. Remove and wash contaminated clothing before rause.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or when disposing of equipment washwaters.

MPORTANT

Do not use on food or feed crops. Injury to or lose of desirable trees or other plants may result from failure to observe the following: Do not apply (except as secommended), or dealer or flush equipment on or near desirable trees or other plants, or on assess where their notes may extend, or in locations where the chemical may be washed or moved into contact with their soots. Do not apply where sunoff water may flow onto agricultural land as Injury to crops may result. Do not contaminate any body of water, including intgestion water. Do not use on water, driveways, tennis courts, or similar areas. Prevent drift of sprey to creptain and other desirable plants. Do not use on dichondra lasms or where violats or ryagrass are the desired species. Keep from contact with fertilizers, insecticides, fungicides and seeds. Following application, the sprey tank should not be used for other than non-crop application. This is extremely important, as low rates of LESCO TFC can titll or severely injure most crops. See SPRAYER CLEAMUP section.

GENERAL INFORMATION

LESCO TFC Herbicide is a dispersible granule to be mixed in water and applied as a spot or broadcast apray treatment to certain species of established grasses. It is noncorresive, nonfaminable, nonvolatile and does not freeze. LESCO TFC spidity inhibits growth of susceptible weeds. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 3 weeks after application depending on growing conditions and weed susceptibility. Warm moist conditions following treatment promote the activity of LESCO TFC while cold, diy conditions delay the activity of LESCO TFC. Weeds hardened off by cold weather or drought stress may not be fully controlled. Degree of control and duration of effect depend on; a) rate used; b) weed spectrum; c) weed size; d) growing conditions at and following time of treatment; e) sol prit; f) precipitation, and g) soil organic metter content.

STORAGE AND DISPOSAL

cutaide of the United States.

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed. Do not contaminate water, food of feed by storage or disposal.

PESTICIDE DISPOSAL: Waster resulting from the use of this product may be disposed of on sits or at an approved waste disposed facility. CONTAINER DISPOSAL: Completely emply beg into application equipment. Then dispose of of emply container in assentiary tandfill or by incineration, or , if allowed by State and local authorities, by burning, if burned stay out of smolet.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LESCO TFC Herbicide should be used only in accordance with the recommendations of this label. The manufacturer will not be responsible for losses or demages resulting from the use of this product in any manner not specifically recommended. User assumes all risks associated with such non-recommended uses.

LESCOTFC Herbicide is to be applied as a postamergence treatment with a hand helder a boom sprayer to control perennial "bunch or clump" grasses such as tall feacue or ryegrains and meny other weeds in actively growing established Kentucky Blaegama, Fine Feacue, Bernasse (when moved over 1/2 inch in height), Bahlagnes, and Bernasse mans. For best results, apply LESCO TFC when the destrable tair is not under any sleep (a.l., feets or drought). Do not apply to destrable Ryegrass tari. Adequate moisture by rainfall or irrigation is needed to activate the herbicide. Do not apply to soil assumated with moisture or during pariods of intense heat. Use a non-fonio-sentential such as LESCO Spreader Sticter at a rate of 1/4 to 1/3 62. per gallon of spray solution.

Chemigation

Do not apply this product through any type of inigation system.

LESCO TFC

DISPERSIBLE GRANULE TURF HERBICIDE

FOR SELECTIVE CONTROL OF TALL FESCUE
AND OTHER WEEDS IN
ESTABLISHED KENTUCKY BLUEGRASS,
FINE FESCUE, BENTGRASS, BAHIAGRASS,
AND BERMUDAGRASS TURF

NET WEIGHT UNITS - 25 GRAMS per BLISTER-PAK

ACTIVE INGREDIENT:

Chlorsuffuron: 2-Chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl] benzenesulfonamide 75% NERT INGREDIENTS: 25%

- <u>2376</u> Total 100%

U.S. Patent No. 4,127,405 EPA Reg. No. 10404-59 EPA Est. No. 43152-PA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately, induce vomiting by giving victim 1 or 2 glasses of water and touching back of threat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. If IN EYES: Immediately flush eyes with plenty of water. Get medical attention if intration pensists. If ON SKIN: Wash affected area with soap and water.

See Side Fane For Aphitional Precautionary Statements

FOR PROFESSIONAL USE ON Omamental Lawns and Turfgrasses

LESCO, INC.

20005 Lake Road - Rocky River, OH 44116

RATES

LESCOTFC is to be applied as a spote soid farms as a broadcast apray. We application may result in temporary suppress desirable grass growth and process of the second and pro

Rate selection is based on weed special tablished plants and on the fine tests textured solts. For low volume or contra application rates per acre. A repeat as Apply LESCO TFC at a rate of 1.0 to 5 blister-pak (2.5 grams) to cover 725 to rate on soits with pH of 7.0 and above.

Weeds Controlled Difference Chickeneed Shartenstructure Harchit Flatt Furnigner Process Controlled at 1 to Pursana Common Special Wild carrot Dandelon's Dandelon's Soldswed Solds

To increase activity and spectrum of with 2,4-D, MCPP, DICAMBA, MCPA o Follow all label precautions for use on Spreader Sticker at a rate of 1/4 to 1/3 A second application 60 days after the condition of the second application 60 days after the condition of the second application 60 days after the condition of the second application 60 days after the condition of the second application 60 days after the condition of the second application 60 days after 60 days 60 day

Weeds Controlled at 2.76 to

Tab Feacue + */ Wild Violet +

SPRAY PREPARATION - Add LESCO TI disperse the water dispersible granules, spray within 72 hours if the tank pH is be Avoid application in low pH conditions.

APPLIC

Apply LESCOTFC with a sprayer until it spray to drift into adjacent plants or tre prevent drift.

SPRAY VOLUME: For hand held or bac to spray 2 to 3 gallons per 1,000 eq.ft. (a feacuse clumps). For larger tank and it LESCO TFC in a minimum of 20 gallons t and above.

AGITATION - Thoroughly mix and agitate excessive foaming.

ive foaming.

LESCO TFC is a very active compound tank, drain and flush with water a minimum. Fill tank half full of water and add 1 to 2g of water. Fill tank and circulate for 15 min Clean nozzles and screens separately, ornamental plants after use with TFC.

LESCO, INC. warrants that this herbicide used in accordance with label directions for its intended purposes. Since timing, in mbdure with other chemicals, and other control, no warranty is given concerning under conditions which are abnormal or of any such use. Amdmt. Version dated:

OCT 3 0 1990
Under u.e Federal Insecticida Finnicida, and Rodeniicide Act, as amended, for the pesticide recipients and recipients and recipients.

সার্ভারি কুলি সাধার্ত্তি তেনে কিছেল ক্রান্ত্রিকালে ব্যবস্থা ভারতা ক্রিক