

ACCEPTED

NOV 26 1990

10404-60

10404-60

11/26/1990

1/3

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
DANGER

Corrosive: Causes irreversible eye damage. Wear goggles or eye shield when handling this product. Do not get in eyes or on clothing.

Harmful or fatal if swallowed: Avoid breathing spray mist.

May be a potential skin sensitizer: Avoid contact with skin or clothing. Remove contaminated clothing and wash before reuse.

First Aid:

If in eyes: Flush with plenty of water, call a physician.
If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

If on skin: Wash exposed areas with soap and warm water after handling or using.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage. Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not contaminate water when disposing of equipment wash-waters. Do not apply when weather conditions favor drift from treated areas. Do not graze treated areas. Do not feed clippings to livestock.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having an allergic reaction should contact a physician.

See Replacement

See Replacement

LESCO

TwoSome™

Flowable Fungicide

Contains a combination of chlorothalonil and fenarimol fungicides

For commercial use only on:
Golf Courses, Parks, Athletic Fields, Commercial and Residential Turfgrass Areas.

Active Ingredients

Chlorothalonil: Tetrachloro-sophthalonitrile 41.5%
Fenarimol: A-(2-chlorophenyl)- α -(4-chlorophenyl)-5-pyrimidinemethanol 2.5%

Inert ingredients 56.0%

Total 100%

Contains 4.17 pounds chlorothalonil plus .25 pounds fenarimol per gallon

EPA Reg. No. 10404-60

EPA Est. No. 37429-GA-01

Keep Out of Reach of Children

DANGER

Read All Directions Before Use.

1 Gallon

Manufactured for:

LESCO, INC., 20005 Lake Road, P.O. Box 16915, Rocky River, Ohio 44116

#011092-1 gallon

#018980-4x1 gallon case

Storage and Disposal
 Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.
 Storage: Store in a cool place. Protect from excessive heat. Avoid freezing.
 Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or mixture is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
 Container Disposal: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Directions for Use:
 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
 Read All Directions Carefully Before Applying
 Not For Use in California

WARRANTY
 LESCO warrants that this fungicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this fungicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

General Information:
 TwoSome is a combination of chlorothalonal, a contact fungicide, and fenarimol, a locally systemic fungicide, in a proven 16:1 ratio for the protection against fungal turfgrass diseases. Two different active ingredients team up for highly effective control of a broader spectrum of turfgrass plant diseases when used according to the directions on this product label.

Chemigation:
 Do not apply this product through any type of irrigation system.

Mixing Instructions:
 Spray tank container several times to assure uniform mixture. TwoSome should be added to the spray tank during the last half of the filling operation. Add the required amount of TwoSome to the spray tank while keeping the agitator running. Agitate thoroughly during the spraying operation to insure a uniform spray mixture. Use only a properly calibrated sprayer with nozzle screens and in-line strainers no finer than 50 mesh. Apply mixture in sufficient water volume for uniform coverage of turfgrass plant surfaces. Do not combine TwoSome in the spray tank with pesticides, surfactants, or fertilizer unless your prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use. Do not mow or water after treatment until spray deposit on turfgrass is thoroughly dry.

Rate Combinations Chart
 TwoSome Fungicide for Turfgrass

Quantity Per 1000 Sq. Ft.	Quantity Per Acre
3.8 fl. oz.	1.00 Gal.
4.4 fl. oz.	1.50 Gal.
5.1 fl. oz.	1.75 Gal.
5.8 fl. oz.	2.00 Gal.
7.8 fl. oz.	2.50 Gal.
9.8 fl. oz.	3.00 Gal.
16.3 fl. oz.	5.00 Gal.
17.8 fl. oz.	5.00 Gal.

LESCO TwoSome FLOWABLE FUNGICIDE

EARLY SPRING PREVENTIVE PROGRAM

Disease	Rate (pounds per Acre)	Spray Interval (Days)	Comments (See label)
Leaf Spot (Helminthosporium, Drechslera spp., Bipolaris spp.)	1.0-1.75	14-21	1
Patch Diseases: Necrotic Ring Spot (Lophophora termis) Summer Patch (Magnaporthe poae) Fusarium blight (Fusarium culmorum) (Fusarium poae)	1.5-3.0	14	7, 8, 9

SUMMER PREVENTIVE PROGRAM

Disease	Rate (pounds per Acre)	Spray Interval (Days)	Comments (See label)
Dollar Spot (Sclerotinia homoeocarpa, or Monodelphoides and Lanzia spp.) Anthracnose (Colletotrichum graminicola) Rhizoctonia Blight/ Large Brown Patch (Rhizoctonia solani)	1.0-1.5	14-21	1
Other Diseases: Pyricularia grisea Curvularia Corticium Red Thread (Laetisaria trichostema) Copper Spot (Stenocarpus sp.) Powdery Mildew (Erysiphe graminis) Stem Rust of Bluegrass (Puccinia spp.)	2.0-2.86	14-21	1
	10-2.0	7-10	1

OTHER DISEASES OF TURFGRASS

Disease	Rate (pounds per Acre)	Spray Interval (Days)	Comments (See label)
Gray Leaf Spot (Pyricularia grisea) Curvularia Corticium Red Thread (Laetisaria trichostema) Copper Spot (Stenocarpus sp.)	1.0-3.0	7-10	1.2
Powdery Mildew (Erysiphe graminis) Stem Rust of Bluegrass (Puccinia spp.)	2.0-3.0	7-14	1.2

LESCO TwoSome FLOWABLE FUNGICIDE

Other Diseases of Turfgrass (cont'd)

Disease	Rate (pounds per Acre)	Spray Interval (Days)	Comments (See label)
Spring Dead Spot (Lophophora termis)	6.00	Single	4, 5
Take-All-Patch (Biarctomyces graminis) Applications	6.00	Two	5, 6

WINTER DISEASES

Disease	Rate (pounds per Acre)	Spray Interval (Days)	Comments (See label)
Fusarium Patch/ Pink Snow Mold (Microdochium nivale) Typhula Blight/ Gray Snow Mold (Typhula incarnata)	5.33	1 or more	5, 6

- 1) Use the high rate and shorter interval during conditions favoring heavy disease pressure.
- 2) Begin applications when conditions favor disease development and repeat applications as long as these conditions persist.
- 3) Applications to areas with visible disease symptoms should be made at the highest rate listed for that disease and continued at the shortest spray interval for that rate until the disease is brought under control. At this time, the lower rate, longer spray interval recommendations can begin.
- 4) Single application in September.
- 5) Application should be made within a 30-day interval prior to snow cover or frozen soil.
- 6) If snow cover is intermittent or lasting repeat applications every 3-4 weeks until disease conditions no longer prevail.
- 7) Make four (4) to five (5) applications at 14-day intervals.
- 8) Applications for necrotic ring spot, summer patch, fusarium blight, take-all patch, and spring dead spot should be applied to the turfgrass and thoroughly irrigated with 1/2 to 1 inch of water.
- 9) Use the higher rate listed when the area to be treated has a previous history of disease infestation.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. May be fatal if inhaled. Do not breath dust vapor or spray mist. Wear goggles, face shield mask or pesticide respirator approved by MSHA or OSHA. Harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

- IF IN EYES:** Hold eyes open and flush with steady gentle stream of water for 15 minutes.
- IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.
- IF INHALED:** Remove victim to fresh air, if not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate gastric lavage. Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

NOTE TO USER: This product may produce temporary allergic side effects characterized by redness of the eyes or mild bronchial irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Persons having an allergic reaction should contact a physician.

OK
on
chlorbutol
kebed