

LESCO

TwoSomeTM

ACCEPTED
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10404-60

Page 1 of 2

Flowable Fungicide

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. May be fatal if inhaled. Do not breathe dust, vapor or spray mist. Wear goggles, face shield mask or pesticide respirator approved by MSHA or OSHA. Harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If In Eyes: Hold eyes open and flush with steady, gentle stream of water for 15 minutes. **If Swallowed:** Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

If Inhaled: Remove victim to fresh air, if not breathing give artificial respiration., preferably mouth-to-mouth. Get medical attention. **If On Skin:** Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids. **NOTE TO USER:** This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Persons having an allergic reaction should contact a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Do not graze treated areas. Do not feed clippings to livestock.

Contains a combination of Chlorothalonil and Fenarimol fungicides

For Commercial Use Only On: Golf Courses, Parks, Athletic Fields, Commercial and Residential Turfgrass Areas.

ACTIVE INGREDIENTS:

Chlorothalonil: Tetrachloroisophthalonitrile	- - -	40.00%
Fenarimol: a-(2-chlorophenyl)-a-(4-chlorophenyl)-5-pyrimidinemethanol	- - -	2.40%

INERT INGREDIENTS:	- - - - -	57.60%
		Total 100.00%

Contains 4.17 lbs. Chlorothalonil plus 0.25 lbs. Fenarimol per gallon.

**KEEP OUT OF REACH OF CHILDREN
DANGER**
Read All Directions and Precautionary Statements Before Use

EPA Reg. No. 10404-60

EPA Est. No. 37429-GA-1

#011092 1 Gallon

#018980 4 x 1 Gallon Case

Manufactured For
LESCO, INC.

20005 Lake Road - P.O. Box 16915
Rocky River, OH 44116

Product Use Directions and Additional Precautionary Statements continued on Page 2.

This is 04-22-91 EPA corrected version of 11-30-90 & 03-06-91 submitted copies.

10404-60
07/18/1991
1/2

LESCO

TwoSome^{T.M.}

Flowable Fungicide

Page 2 of 2

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. STORAGE: Store in a cool place. Protect from excessive heat. Avoid freezing. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions before using. Not for Use in California.

GENERAL INFORMATION

TwoSome is a combination of chlorothalonil, a contact fungicide and Fenarimol a locally systemic fungicide, in a proven 16 to 1 ratio for the protection against fungal turfgrass diseases. Two different active ingredients team up for highly effective control of a broader spectrum of turfgrass plant diseases when used according to the directions on this product label.

CHEMIGATION: Do not apply this product through any type of irrigation system.

MIXING INSTRUCTIONS

Slowly invert container several times to assure uniform mixture. TwoSome should be added to the spray tank during the last half of the filling operation. Add the required amount of TwoSome to the spray tank while keeping the agitator running. Agitate thoroughly during the spraying operation to insure a uniform spray mixture. Use only a properly calibrated sprayer with nozzle screens and in-line strainers no finer than 50 mesh.

Apply mixture in sufficient water volume for uniform coverage of turfgrass plant surfaces. Do not combine TwoSome in the spray tank with pesticides, surfactants, or fertilizers unless your prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use. Do not mow or water after treatment until spray deposit on turfgrass is thoroughly dry.

NOTICE: LESCO INC. warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Comments for Chart; 1 thru 9

- 1) Use the high rate and shorter interval; during conditions favoring heavy disease pressure.
- 2) Begin applications when conditions favor disease development and repeat applications as long as these conditions persist.
- 3) Applications to areas with visible disease symptoms should be made at the highest rate listed for that disease and continued at the shortest spray interval for that rate until the disease is brought under control. At this time the lower rate and the longer spray interval recommendations can begin.
- 4) Single application in September.
- 5) Application should be made within a 30 day interval prior to snow cover or frozen soil.
- 6) If snow cover is intermittent or lacking, repeat applications every 3 to 4 weeks until disease conditions no longer prevail.
- 7) Make 4 to 5 applications at 14 day intervals.
- 8) Applications for necrotic ring spot, summer patch, fusarium blight, take-all-patch and spring dead spot should be applied to turfgrass and thoroughly irrigated with 1/2 to 1 inch of water.
- 9) Use the higher rate listed when the area to be treated has a previous history of disease infestation. (Format Amend. to add Fl.Oz/1m sq.ft. rates to chart) (03 04 91)

Disease	Recommended Rates Fl. Oz. / 1,000 sq.ft.	Recommended Rates Gals./Acre	Spray Interval (Days)	Comments (See Left Panel)
---------	--	------------------------------	-----------------------	---------------------------

EARLY SPRING PREVENTATIVE PROGRAM

Leaf Spot (<i>Helminthosporium Dreschlera spp.</i> <i>Bipolaris spp.</i>)	3.0 to 5.25	1.0 to 1.75	14 to 21	1
PATCH DISEASES Necrotic Ring Spot (<i>Leptosphaeria korrae</i>) Summer Patch (<i>Magnaporthe poae</i>) Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>)	4.5 to 9.0	1.5 to 3.0	14	7,8 & 9

SUMMER PREVENTATIVE PROGRAM

Dollar Spot (<i>Sclerotinia homeocarpa</i> , or <i>Moellerodiscus</i> and <i>Lantia spp.</i>)	3.0 to 4.5	1.0 to 1.5	14 to 21	1
Anthracnose (<i>Colletotrichum graminicola</i>)	6.0 to 8.0	2.0 to 2.66	14 to 21	1
Rhizoctonia Blight/ Large Brown Patch (<i>Rhizoctonia solani</i>)	3.0 to 6.0	1.0 to 2.0	7 to 10	1

OTHER DISEASES OF TURFGRASS

Gray Leaf Spot (<i>Pyricularia grisea</i>)	3.0 to 9.0	1.0 to 3.0	7 to 10	1 & 2
Curvularia	3.0 to 9.0	1.0 to 3.0	7 to 10	1 & 2
Corticum Red Thread (<i>Laetisaria luciformis</i>)	3.0 to 9.0	1.0 to 3.0	7 to 10	1 & 2
Copper Spot (<i>Gloeocercospora sorghi</i>)	3.0 to 9.0	1.0 to 2.0	10 to 21	1 & 3
Powder Mildew (<i>Erysiphe graminis</i>)	8.0 to 16.0	2.66 to 5.33	30	1 & 3
Stem Rust of Bluegrass (<i>Puccinia spp.</i>)	6.0 to 9.0	2.0 to 3.0	7 to 14	1 & 2
Spring Dead Spot (<i>Leptosphaeria korrae</i>)	18.0	6.00	Single Application	4 & 8
Take-All Patch (<i>Gaeumannomyces graminis</i>)	18.0	6.00	Two Applications	5 & 8

WINTER DISEASES

Fusarium Patch/Pink Snow Mold (<i>Microdochium nivale</i>)	16.0	5.33	One or More Applications	5 & 6
Typhula Blight/Gray Snow Mold (<i>Typhula incarnata</i> & <i>Typhula tibrikawensis</i>)	16.0	5.33	One or More Applications	5 & 6

e/c