

10404-63

06/07/2002

1/2

NOT FOR USE on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

SUGGESTED SPREADER SETTINGS*

Recommended Rates in lb per 1,000 sq ft

Table with columns: Spreader, Ground Speed, XX lb, XX lb, XX lb, XX lb. Rows include aaa, bbb, ccc, ddd, fff, ggg, hhh, iii.

*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application using the LESCO Calibrating Gauge #009952.

ACTIVE INGREDIENT: Oxadiazon; 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) 1,3,4-oxadiazolin-5-one 0.95%
INERT INGREDIENTS: 99.05%
Total: 100.00%

KEEP OUT OF REACH OF CHILDREN WARNING

FIRST AID table with categories: If inhaled, If on skin or clothing, If in eyes, If swallowed. Includes HOT LINE NUMBER section.

WHEN TO APPLY

Table for application timing: Appropriate Times to Apply This Pre-emergent Herbicide. Columns: Jan, Feb, Mar, Apr, May, June, July, Aug, Sep, Oct, Nov, Dec. Includes application instructions for Crabgrass, Goosegrass, and Poa annua.

PRECAUTIONS FOR USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipedegrass since damage will occur. Do not use on putting greens or "tees" due to varying cultural practices used.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes, or clothing. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Storage: Store product in its original bag, in a cool, dry, locked place out of the reach of children.

FOR LAWN AND ORNAMENTAL TURF

When applied according to directions, this product can be used safely on the following established turfgrass varieties: Bermudagrass, Perennial Bluegrass, Perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysiagrass lawns and ornamental turf.

CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS

Annual Bluegrass (Poa annua), Barmyardgrass, Crabgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed.

HOW TO APPLY


Measure the area to be treated and uniformly apply amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader application on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate".

HOW MUCH TO USE

For Crabgrass: Apply 50 lb/10,400 sq ft (4.8 lb/1,000) when turf has been in a yearly maintenance program (equal to 2 lb A.I. herbicide per acre). Apply 50 lb to treat 6,900 sq ft (7.25 lb/1,000) of lawn if it has been heavily infested with crabgrass (equal to 3 lb A.I. herbicide per acre).

Goosegrass and Prostrate Spurge: These weeds do not usually germinate until late spring or early summer. Apply 50 lb to treat 5,200 sq ft of lawn (9.8 lb/1,000 sq ft; 4 lb A.I./acre) and apply 1 to 2 weeks prior to seed germination.

For Annual Bluegrass (Poa annua): Apply 50 lb per 5,200 to 6,050 sq ft of lawn (8.25-9.8 lb/1,000 sq ft; equal to 3.5 to 4 lb A.I. herbicide per acre). Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination.



(OPTIONAL LANGUAGE: MINI, ELITE, NOVEX, PROFESSIONAL TURF FERTILIZER) GUARANTEED ANALYSIS

Table of Guaranteed Analysis: TOTAL NITROGEN (N) XX.XX%, AVAILABLE PHOSPHATE (P2O5) XX.XX%, SOLUBLE POTASH (K2O) XX.XX%, MAGNESIUM (Mg) total XX.XX%, SULFUR (S) total XX.XX%, COPPER (Cu) total XX.XX%, IRON (Fe) total XX.XX%, MANGANESE (Mn) total XX.XX%, ZINC (Zn) total XX.XX%, DERIVED FROM: XXXX, XXXX, XXXX, ..., CHLORINE (Cl) MAX. XX.XX%

WARRANTY

LESCO warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 15885 Sprague Rd., Strongsville, OH 44136-1799, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-1.com.

ACCEPTED

JUN - 7 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10404-63

Net Weight: XX lb (XX kg)

EPA Reg. No. 10404-63
EPA Est. No. 10404-OH-4 (M),
10404-FL-1 (S), 59197-CA-1 (C)
First letter of lot code indicates manufacturing site.

758073XXXXX

FXXXX
Manufactured by: **LESCO, Inc.**
15885 Sprague Road
Strongsville, Ohio 44136-1799

#XXXXXXXX

