
 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p>	EPA Reg. Number:	Date of Issuance:
	10404-63	NOV 8 2007
Term of Issuance:		
Name of Pesticide Product: Ronstar 0.95% Plus Fertilizer		
<p>NOTICE OF PESTICIDE: <u> </u> Registration <u> X </u> Reregistration</p> <p>(under FIFRA, as amended)</p>		
<p>Name and Address of Registrant (include ZIP Code):</p> <p>LESCO, Inc. 1301 East 9th Street Cleveland, OH 44114-1849</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:</p> <ol style="list-style-type: none"> 1. Change the Signal word to "Caution". 2. Delete the Hazards to Humans and Domestic Animals section from the label. 3. To the Non-Agricultural Use Requirements box replace "Thoroughly wash face...clothing before reuse." with "Do not enter or allow others to enter the treated area until dusts have settled." 4. To the Environmental Hazards section add "or rinseate" after "washwaters". 		

5. To the Storage and Disposal section change "Storage" to "Pesticide Storage".

<p>Signature of Approving Official:</p>  <p>Kathryn V. Montague, Acting Product Manager (23) Herbicide Branch, Registration Division (7505P)</p>	<p>Date:</p> <p>NOV 8 2007</p>
---	--------------------------------

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

3 8 7

LESCO Ronstar 0.95% Plus Fertilizer

EPA Reg. No:10404-63

Draft: Revised 2/15/06

Page 1 of 5

RONSTAR® 0.95% Plus Fertilizer

For Sale To and Use By Professional Applicators Only. Not for Sale To or Use By Homeowners/Consumers

ACTIVE INGREDIENT:

Oxadiazon; 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)

Δ1,3,4-oxadiazolin-5-one 0.95%

INERT INGREDIENTS: 99.05%

Total 100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If inhaled	<ul style="list-style-type: none"> ◆ Move person to fresh air. ◆ If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. ◆ Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> ◆ Take off contaminated clothing. ◆ Rinse skin immediately with plenty of water for 15-20 minutes. ◆ Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> ◆ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ◆ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ◆ Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> ◆ Call poison control center or doctor immediately for treatment advice. ◆ Have person sip a glass of water if able to swallow. ◆ Do not induce vomiting unless told to do so by the poison control center or doctor. ◆ Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going to treatment. You may also contact 1-(800) 424-9300 for emergency treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield, or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing dust, contact with skin, eyes, or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED

NOV 8 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticides
registered under EPA Reg. No.

10404-63

USER SAFETY RECOMMENDATIONS

Users should; wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Do not apply this product through any type of irrigation system. Do not graze livestock in treated areas. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

General Precautions and Restrictions

The total maximum rate of application may not exceed 632 lbs of product per acre per year (equivalent to 6 lbs ai/A/year), except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 842 lbs of product per acre per year (equivalent to 8 lbs ai/A/year).

A total maximum single application rate may not exceed 421 lbs of product per acre (equivalent to 4 lbs ai/A).

Frequency of application ranges from 1 to 3 applications per season. Can be applied at a minimum single application rate of 2 lbs ai/A up to a maximum single application rate of 4 lbs. ai/A to turf and ornamentals. Annual maximum application rate is 8 lbs ai/A/year. Maximum number of 3 applications per year.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls, shoes plus socks, and chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USES REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forests, nurseries, sod farms or greenhouses. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Do not store or use in or around the home or home garden.

STORAGE: Store product in its original bag, in a cool, dry, locked place out of the reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows:

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

X-X-X

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	xx.xx%
xx.xx% Ammoniacal Nitrogen	
xx.xx% Urea Nitrogen*	
AVAILABLE PHOSPHATE (P₂O₅)	xx.xx%
SOLUBLE POTASH (K₂O)	xx.xx%
SULFUR (S) Total	xx.xx%
xx.xx% Free Sulfur (S)	
xx.xx% Combined Sulfur (S)	
IRON (Fe) Total	xx.xx%
xx.xx% Water Soluble Iron	
MANGANESE (Mn) Total	xx.xx%
xx.xx% Water Soluble Manganese	
DERIVED FROM:	
CHLORINE (Cl) MAX.	XX.XX%
*XX.XX% Slowly available Nitrogen from LESCO Poly Plus® Sulfur Coated Urea.	

USE SITES

(NOT FOR USE ON HOME LAWNS)

For commercial use on residential turf (i.e. apartment/condominium complexes, parks, athletic fields, playgrounds, and cemeteries) and on golf course. In addition, is used on sod farms and on conifer nurseries and landscapes (i.e. industrial sites, ornamental, roadside plantings, woody, ornamental shrubs, vines and trees, and herbaceous ornamentals). Use sites are classified as non-food sites (i.e. primarily golf course fairways), residential outdoor use, roadsides and nurseries. When applied according to directions, this product can be used safely on the following established turfgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, TALL FESCUE, & ZOYSIAGRASS lawns and ornamental turf.

CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS

Annual Bluegrass (*Poa annua*), Barnyardgrass, Crabgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed, and others.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader applications on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After application, if rain is not expected within 24 hours, sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control. For best results, water-in the product as soon as practical after application.

RATES FOR WEED CONTROL IN TURF GRASSES

FOR CRABGRASS: Apply 50 lb per 10,400 sq ft when turf has been in a yearly maintenance program (equal to 2 lb A.I. herbicide per acre). Apply 50 lb to treat 6,900 sq ft of lawn if it has been heavily infested with crabgrass (equal to 3 lb A.I. herbicide per acre). Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When to Apply" chart.

GOOSEGRASS AND PROSTRATE SPURGE: These weeds do not usually germinate until late spring or early summer. Apply 50 lb to treat 5,200 sq ft (4 lb A.I./acre) and apply 1 to 2 weeks prior to seed germination. See chart for "When to Apply". Do not exceed maximum rate for St. Augustinegrass which is 50 lb per 6,900 sq ft.

FOR ANNUAL BLUEGRASS (*Poa annua*): Apply 50 lb per 5,200 to 6,050 sq ft of lawn (equal to 3 to 4 lb active herbicide per acre). Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control Annual Bluegrass present at time of application; therefore, only germinating seeds. Refer to chart for proper timing in your locale. Annual Barnyardgrass

and Prostrate Spurge may also be controlled by this same rate. Do not exceed maximum rate for St. Augustinegrass, which is 3 lb active ingredient / acre.

Poa annua "ANNUAL BLUEGRASS" CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY: Apply 50 lb to treat 10,400 sq ft (2 lb A.I./acre). Apply in late summer to early fall prior to *Poa annua* spp. *annua* seed germination. Established plants will not be controlled. This product should be applied no closer than 60 days prior to overseeding. NOTE: Do not use where *Poa trivialis* is present in overseed mix. IMPORTANT: In many areas Annual Bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype, "*Poa annua* spp. *annua*" which germinates from seed every year is controlled, while the biotype, "*Poa annua* spp. *raptans*" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning you locale.

FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE, AND CARPETGRASS: Where these weeds are a problem, apply 50 lb to treat 5,200 to 6,050 sq ft of lawn (equal to 3.5 to 4 lb active herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control emerged and growing weeds; only effective against germinating weed seeds. Observe maximum rate for St. Augustinegrass of 3 lb active ingredient / acre.

SUGGESTED SPREADER SETTINGS

Recommended Rates in lb per 1,000 sq ft

Spreader	Ground Speed	4.8 lb	7.25 lb	8.25 lb	9.6 lb
LESCO	3 mph	XX	XX*	XX*	XX*
Cyclone® or Spyker®	3 mph	XX	XX*	XX*	XX*
Lely®	4½ mph	XX	XX*	XX*	XX*
Vicon®	4½ mph	XX	XX*	XX*	XX*

* 2 passes

settings are approximate. The age and condition of spreader and operator's walking speed can cause variations. Be sure spreader is properly calibrated. Check spreader over 500 sq ft using ½ the "sq ft rate" then adjust spreader to apply exact amount.

WHEN TO APPLY

Appropriate Times to Apply This Pre-emergent Herbicide:

Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
1	2	3	3a
4	2a	4

1. For CRABGRASS and other Grasses and Broadleaf Weeds germinating in the Spring in Northern Turf.
2. For GOOSEGRASS and other Late Germinating Annual Grasses and Broadleaf Weeds such as PROSTRATE SPURGE.
- 2a. GOOSEGRASS in Southern Turf. See PRECAUTIONS IN USING.
3. For *Poa annua* in Northern Turf. 3a. For *Poa annua* in Southern Turf.
4. For Fall-Winter-Spring control of annual weeds in Southern Turf.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipedegrass since damage will occur. Do not use on putting greens or "tees" due to varying cultural practices used. Use only the rate of 50 lb to treat 6,900 sq ft on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended in the Ornamental Plants section for this label. For newly seeded lawns, do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

RATES FOR WEED CONTROL IN ORNAMENTAL PLANTS

If treatment is desired in non-turf areas, where weeds are presently growing, first remove existing weeds. Rake and smooth the soil by "breaking-up" clumps. Apply at rate as directed for specific weeds (50 lb to treat either 6,900 or 10,400 sq ft). This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control. For best results, water-in product as soon as practical after application.

WEED CONTROL IN CONTAINER GROWN PLANTS

This product is recommended on a wide variety of Container and Field Grown Woody Ornamental Shrubs, Vines and Trees. It may be used on the following list of Container Grown Plants. Use at the rate of 50 lb per 5,200 sq ft of pot soil surface area. Plants in 14-inch containers set out in 12-foot wide rows x 100 feet long will require 1/5 lb of this product per row of 10 pots (equivalent to approx. 4 lb A.I. per acre): Apply with a "shoulder strap held" grass-seeder or granule applicator that will direct the granules evenly to each pot soil surface for Agapanthes, Andromeda, Arborvita, Archtostaphylos, Azalea, Barberry, Boxwood, Common-Japanese, Calistemon, Camellia, Ceanothus, Cedar, Cotoneaster, Cypress, Dogwood, Douglas Fir, Eucalyptus, Euonymus-evergreen, Forsythia, Gardenia, Ginko, Heath, Heather, Hedera, Holly, Honeysuckle, Jasmine, Juniper-varieties, Justicia, Leucothoe, Magnolia, Mahonia, Myrtle,

Osmanthus, Periwinkle, Photina, Pine-varieties, Pittosporum, Podocarpus, Privet, Pyracantha, Quercus, Raphiolepus, Rhododendron, Rose, Star Jasmine, Syzygium, Viburnum, Xylosma.

FOR FIELD GROWN ORNAMENTALS

***Apply at the rate of 50 lb per 6,900 or 10,400 sq ft of area.**

This product may be used around the following Established Plants.

Arborvita, Ash, Azalea, Barberry-Japanese, Birch, Camellia, Cotoneaster, Crabapple, Dogwood, Euonymus, Forsythia, Holly, Honeysuckle, Juniper, Lilac, Maple, Oak, Periwinkle, Pine, Pittosporum, Privet, Rhododendron, Rose, Russian Olive, Spruce, Taxus, Viburnum, Xylosma. Ground Covers: Carpobrotus, Osteospermum, Sedum, Algerian Ivy, Vinca Major, English Ivy, Pachysandra, Ajuga, Euonymus.

***Use higher rate (50 lb /6,900 sq ft) for hard to kill weeds.**

May be used around the following Newly Transplanted Ornamentals.

Camellia, Dogwood, Forsythia, Hedera, Holly, Juniper, Oak, Periwinkle, Pine, Privet, Rhododendron, Russian Olive, Spruce, Taxus, Tsuga.

To obtain continued season-long weed control on the following list of plants, additional applications may be made 60 days after the previous application in the following situations.

APPLY 2 REPEAT TREATMENTS

50 lb/ 6,900 SQ FT ON FOLLOWING PLANTS

Established - Container Grown Plants — Abelia, Common Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron.

Newly Transplanted - Container Grown Plants – Common Boxwood, English Ivy, Juniper, Pine, Rhododendron.

Forest Nurseries - For Pines – Loblolly, Slash, Eastern White.

WARRANTY

LESCO, Inc. warrants that this product conforms to the chemical description on the label. When used in accordance with label directions under normal conditions, this product is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a1.com>

Contains RONSTAR Herbicide, a registered trademark of Bayer. LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Spyker and Cyclone are registered trademarks of Spyker Spreaders LLC. Lely is a registered trademark of C Van Der Lely N.V. Vicon is a registered trademark of Ransomes Cushman-Ryan Company.

Created: xxxxxx

(8/16/02)

Rev.xxxxxx BS

G:\REGULWP\STRONGSV\Labels in Progress\069173 - temp.doc

Net Weight: 50 lb (22.7 kg)



EPA Reg. No. 10404-63	
EPA Est. No. 82757-OH-001 (M), 82757-FL-001 (S) <i>First letter of lot code indicates manufacturing site.</i>	<p style="text-align: center;">F1560 Distributed by: LESCO, Inc. 1301 East 9th Street Cleveland, OH 44114-1849</p>

#XXXXXX