

-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (H-7505C)

BEST AVAILABLE COPY

BEST AVAILABLE

LESCO

1% DURSBAN® GRANULAR INSECT CONTROL Granules Bait

For Effective Long Lasting Control of Chinch Bugs, Asian Roaches, Ants, Mole Crickets and Other Outdoor Insects

FOR MOUND TREATMENT CONTROL OF IMPORTED FIRE ANTS
IN LAWNS AND RECREATIONAL AREAS
1 LBS TREATS UP TO 400 SQ.FT.

Controls Ants (Including Carpenter, Argentine, & Foraging), Armyworms (Beet, Fall, Yellow-striped), Centipedes, Chiggers(1), Chinch Bugs(2), Clover Mites, Cutworms, Crickets, Earwigs, Fiery Skipper, Fleas, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Mole Crickets(4), Roaches (Asian), Sod Webworms (Lawn Moth)(3), Sow Bugs, Springtails, Ticks (Brown Dog Tick) (1), Turfgrass Weevil (Hyperodes), European Crane Fly (5), White Grubs (Black Turfgrass Ateenius, European Chafer, Southern Chafer, Japanese Beetle, and June Beetle larvae) (5), Bluegrass Billbugs(6) Imported Fire Ants (7).
(See numbered paragraphs for insect control)

USE FOR CRAWLING INSECT CONTROL OUTSIDE OF HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS

ACTIVE INGREDIENT:

Chlorpyrifos: [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate], 1.00%

INERT INGREDIENTS: 99.00%

DURSBAN® Brand Insecticide is a Registered Trademark of DowElanco Total 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. **IF ON SKIN:** In case of contact, remove contaminated clothing and wash affected area with soap and water. **NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and water supplies. Keep children and pets off treated area until this material is washed into the soil and the grass is dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to contaminate water when disposing of equipment washwaters. Use only as directed.

SEE SIDE/PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

EPA REG. NO. 10404-AT(67)

EPA EST. NO. 10404-ON-1

NET WEIGHT (1 lb., 5 lb., 25 lb., 40 lb.) POUND(S)

LESCO, INC.

20005 LAKE ROAD - ROCKY RIVER, OHIO 44116

APR 12 1991

For the Federal Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10404-67

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty container. Wrap container and put in trash.

SPECIFIC DIRECTIONS

The following numbered paragraphs correspond to numbers in parentheses (p) on the front panel following the insect names. Insects not numbered may be controlled by following the GENERAL DIRECTIONS. Repeat treatments as needed for control.

(1) CHIGGERS, BROWN DOG TICKS, & FLEAS

Use 1 lb./400 Sq.Ft. (2.5 lb./1,000 Sq.Ft.) for area control of these insects, infesting non-crop land areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or a possible public health problem. DO NOT allow public use of treated area during applications. For Fleas (out-of-doors, only) treat sleeping quarters and other areas frequented by pets such as beneath porches and along fencing.

(2) CHINCH BUGS

Apply 1 lb. / 300 to 400 Sq Ft. (2.5 to 3.5 lbs./1,000 Sq.Ft. or 110 to 150 lb/A) when insects appear and/or their damage is imminent. To prevent damage apply as soon as daily temperatures reach 60°F and repeat every 2 months. Use lower rate as preventive treatment or for light infestations; the higher rate when insects are present in large numbers and turf is being damaged.

(3) SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. Apply at rate of 2.5 lbs. per 1,000 Sq.Ft.

(4) MOLE CRICKETS

Use 4 lbs. per 1,000 sq ft. Treat suspected problem areas when insects become a problem or as recommended by your local Ag. Extension Service Specialist.

The Best Time to Control Adult Mole Crickets

Jan. Feb. Mar. Apr. May Jun. July Aug. Sep. Oct. Nov. Dec.

1.

2.

1. Indicates Egg Laying time. 2. Best time to Control Crickets

(5) WHITE GRUBS; CRANE FLY LARVAE

To control White Grubs (Black Turfgrass Ateenius, European Chafer, Southern Masked Chafer, Japanese Beetle and June Beetle Larvae), use 4 lbs to treat 1,000 sq ft. To improve efficacy apply enough water to thoroughly wash the insecticide into the turf. For best results remove thatch first and treat when grubs are young and actively feeding near the soil surface, usually during late July & August. Consult your local Agricultural Extension Service Specialist for best time to treat. European Crane Fly larvae will be controlled using 3.5 lbs per 1,000 Sq Ft. or will be controlled simultaneously with the same application being applied for control of "White Grubs" etc. (4 lbs/1,000 Sq Ft.) Consult your Agricultural Agent for proper time to apply. For best results, first remove thatch. Water the granules into the soil.

(6) BLUEGRASS BILLBUGS

Use 4 lbs./1,000 sq ft. for Billbug grubs. To improve control apply enough water to wash granules into soil where insects are feeding.

(7) IMPORTED FIRE ANT CONTROL

MOUND TREATMENT: For treatment of Fire Ant mounds occurring in lawns, L-1 and L-2 areas, treat individual ant mounds measuring from 2 to 12 inches in diameter by applying 8 tablespoons (1 1/2 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment then by applying 2 tablespoons (0.4 oz.) sprinkled evenly over the center of the mound. After application of the granules, gently sprinkle the treated area with a minimum of 1 quart of water per 6 inches of mound diameter. Apply in cool weather (65 to 80 degrees F) or in early morning or late evening. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide. Repeat treatments as needed for additional control but not more than once every 7 days.

GENERAL DIRECTIONS FOR LAWNS, AND ANTS

Apply 1% DURSBAN granules, sod for non-crop land as type spreader or first appear at the insects. Increase as listed under turf with water. Repeat treatment graph for, No. (2)

For control of Yellow-striped, Cutworms, Crickets, Greenbugs, Leafhoppers, Sowbugs, Springtails

Refer to SPECIFIC DIRECTIONS panel for rates to surface. Observe

USE FOR CONTROL OF HOMES, WAREHOUSES, FOOD

To protect against be placed around panels, Ticks, Roaches (Asian) Use 1 pound to the structure. As they cause problems and then as needed. Keep children and the area is

lbs./Product	1,000 Sq. Ft.
1.0	
2.5	
3.5	
4.0	

NOTE: LESCO chemical description under its intended purpose and plant and soil factors affecting its efficacy. LESCO is a registered trademark of DowElanco. For directions or use, see the label. For more information, contact your local Agricultural Agent or LESCO.