

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Frank Vetter  
LESCO, Inc.  
15885 Sprague Rd.  
Strongsville, OH 44136-1772

NOV 30 2004

Dear Mr. Vetter:

Subject: Label Revision Amendment  
LESCO Pre-M 1.31% Plus Fertilizer  
EPA Reg. No. 10404-82  
Submission dated July 8, 2004

The labeling, referred to above, submitted in connection with the application under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the following changes:

1. "Under Precautionary Statements";  
"Under Caution";  
Add "Avoid breathing dust."
2. Under Suggested Spreader Settings";  
Replace "aaa," "bbb," etc., with the actual names of the spreaders.
3. Remove the PPE sentence from the Precautionary Statements. Add a section on "Personal Protective Equipment" with the following requirements:  
Applicators and other handlers must wear:
  - ◆ Long-sleeved shirt and long pants.
  - ◆ Chemical resistant gloves made of any waterproof materials, such as butyl rubber >14 mils, nitrile rubber >14 mils, or neoprene rubber >14 mils.
  - ◆ Shoes plus socks.
  - ◆ Follow manufacturers' instructions for changing/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.
4. Under Environmental Hazards  
Add "or rinse" after "equipment washwaters"
5. It is recommended that you add a section on "User Safety Recommendations" with the following entries:

- ◆ Users should wash hands before eating, drinking, chewing guns, using tobacco, or using the toilet.
- ◆ Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- ◆ Remove contaminated clothing and wash separately before reuse.

6. Under "Directions for Use";

- ◆ Remove "or sprays have dried" from the last bullet point.
- ◆ Add "Do not apply this product in a way that will contact any person or pets, either directly or by drift.
- ◆ Add the following spray drift management statement:  
"Do not allow dust to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals."

7. "Under Application Rates":


Specify a maximum rate of application (times or lbs./unit area) per year.

8. "Under Storage and Disposal";

Change "Storage" to "Pesticide Storage," "Product Disposal" to "Pesticide Disposal," and "Empty Container Disposal" to "Container Disposal."

The amended label supersedes all previously accepted labels. A stamped copy of the revised label is enclosed for your records. Please submit one copy of your final printed label before you release the product for shipment.

Sincerely yours,



Jim Tompkins  
Product manager (25)  
Herbicide Branch  
Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	Bien							
DATE ▶	11-16-04							




## PRE-M® 1.31% Plus Fertilizer

- Not for use on turf being grown for sale or for other commercial use as sod, or for commercial seed production, or for research purposes.
- For control of specified weeds in or on residential lawns, corporate lawns, parks, recreational areas, golf courses, cemeteries, and athletic fields.
- Preemergent control of crabgrass, goosegrass, foxtail, chickweed, ~~local spurge~~, knotweed, and Poa annua.
- Provides quick green-up from soluble nitrogen.
- Prevents annual grassy and broadleaf weeds.

ACCEPTED  
with COMMENTS

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.  
10404-82

NOV 30 2004



(OPTIONAL LANGUAGE: MINI, ELITE, NOVEX, PROFESSIONAL TURF FERTILIZER)  
GUARANTEED ANALYSIS

<b>TOTAL NITROGEN (N)</b>		XX.XX%
XX.XX%	Ammoniacal Nitrogen	
XX.XX%	Urea Nitrogen*	
XX.XX%	Water Insoluble Nitrogen	
XX.XX%	Water Soluble Controlled Release Nitrogen	
<b>AVAILABLE PHOSPHATE (P<sub>2</sub>O<sub>5</sub>)</b>		XX.XX%
<b>SOLUBLE POTASH (K<sub>2</sub>O)</b>		XX.XX%
<b>MAGNESIUM (Mg) total</b>		XX.XX%
XX.XX%	Water Soluble Magnesium	
<b>SULFUR (S) total</b>		XX.XX%
XX.XX%	Free Sulfur (S)	
XX.XX%	Combined Sulfur (S)	
<b>COPPER (Cu) total</b>		XX.XX%
XX.XX%	Soluble Copper (Cu)	
<b>IRON (Fe) total</b>		XX.XX%
XX.XX%	Soluble Iron (Fe)	
<b>MANGANESE (Mn) total</b>		XX.XX%
XX.XX%	Water Soluble Manganese	
<b>ZINC (Zn) total</b>		XX.XX%
XX.XX%	Soluble Zinc (Zn)	
DERIVED FROM:XXXX, XXXX, XXXX, ...		
<b>CHLORINE (Cl) Max.</b>		XX.XX%
*X.XX% Slowly available Nitrogen from LESCO® Poly Plus® Sulfur Coated Urea		

<b>Active Ingredient:</b>	
Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine*	1.31%
<b>Other Ingredients:</b>	
	98.69%
Total	100.00%

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441.

**SUGGESTED SPREADER SETTINGS**

Recommended Rates in lb per 1,000 sq ft

Spreader	1.8 lb	2.6 lb	3.5 lb
aaa	XX	XX	XX
bbb	XX	XX	XX
ccc	XX	XX	XX
ddd	XX	XX	XX
fff	XX	XX	XX
ggg	XX	XX	XX
hhh	XX	XX	XX
iii	XX	XX	XX

**IMPORTANT:** Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Applicators must wear long-sleeved shirt, long pants, shoes and socks when applying this product.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters.

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at time of pesticide application.
- Observe all cautions and limitations in this leaflet and on the labels of products used in combination with this product. Do not use this product other than in accordance with the instructions set forth on this label. The use of this product not consistent with this label can result in injury to turf, animals, or persons. Keep container closed to avoid spills and contamination.
- DO NOT apply this product through any type of irrigation system.
- Do not enter or allow others to enter the treated area until dusts have settled or sprays have dried.

### GENERAL INFORMATION

This product is a combination pre-emergent herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in turfgrass. This product is labeled for use on the following turfgrass species: Kentucky Bluegrass, Perennial Ryegrass, Fine Fescue, Tall Fescue, Bentgrass, *Poa annua* (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, and Zoysiagrass.

This product controls weeds as they germinate but will not control established weeds.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, this product will not cause turf injury. Over-application can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken turfgrass and increase the possibility of turf damage from this product.

### APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a pre-emergent herbicide barrier in the soil. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

### APPLICATION RATES: RATE AND APPLICATION CHART

Turfgrass Species	Weeds	Rates this Product	Comments:
<b>NORTHERN GRASSES</b>			
Kentucky Bluegrass	crabgrass	2.6 to 3.5 lb/1,000 sq ft	Apply a repeat application of 1.8 to 2.6 lb/1,000 sq ft (78 – 115 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
Perennial Ryegrass	foxtail	(115 – 153 lb/acre) as	
Fine Fescue	<i>Poa annua</i>	the initial application prior	
Tall Fescue	barnyardgrass	to weed germination in	
	fall panicum	the spring.	
	oxalis		
	prostrate spurge		
	purslane		
	knotweed		
	evening primrose		
	hop clover		
	goosegrass	2.6 to 3.5 lb/1,000 sq ft	Apply a repeat application of 2.6 lb/1,000 sq ft (115 lb/acre)
		(115 – 153 lb/acre) in the	
		spring prior to weed	
		germination.	

	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre)	Apply in late summer or early fall prior to weed germination.
Bentgrass and established <i>Poa annua</i> * (1/2 inch height or taller)	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	2.6 lb/1,000 sq ft (115 lb/acre) in the spring prior to weed germination.	Apply a repeat application of 1.8 to 2.6 lb/1,000 sq ft (78 – 115 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	2.6 lb/1,000 sq ft (115 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 2.6 lb/1,000 sq ft (115 lb/acre) after a minimum of 4 weeks.
	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.6 lb/1,000 sq ft (115- lb/acre)	Apply in late summer or early fall prior to weed germination.
<b>SOUTHERN GRASSES</b> Bermudagrass** St. Augustinegrass Bahia grass Zoysiagrass Centipedegrass Tall Fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 1.8 to 2.6 lb/1,000 sq ft (78 – 115 lb/acre) after a minimum of 4 weeks if necessary.
	goosegrass	2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre) prior to weed germination in the spring. Apply a repeat application of 2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre) after a minimum of 4 weeks.	An additional application of 2.6 lb/1,000 sq ft (115 lb/acre) may be applied for extended goosegrass control a minimum of 4 weeks after the second application.
	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre)	Apply in late summer or early fall prior to weed germination.

\*Not for use on greens or tees.

\*\*Rates do not apply to Bermudagrass greens and tees.

The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 14 days, erratic weed control may result.

### STORAGE AND DISPOSAL

**STORAGE:** Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PRODUCT DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call or [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

**EMPTY CONTAINER DISPOSAL:** Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.

**SPILL:** In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

### PRECAUTIONS

Use only on well-established turfgrass with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times.

On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application.

Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species.

Do not use on Bentgrass or *Poa annua* (greens and tees) or injury may occur.

Delay reseeding or winter overseeding treated turfgrass for at least 2 months following application. Do not sprig turfgrass for 3 months after application.

### WARRANTY

LESCO, Inc. warrants that this herbicide conforms to the chemical description on this label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 15885 Sprague Rd, Strongsville, OH 44136-1772, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-a1.com>.

LESCO, Pre-M and Poly Plus are registered trademarks of LESCO Technologies, LLC.

G:\REGUL\Product Labels\Maters\EPA Documents\PreM 1.31.doc

Created: xx/xx/xx (XXXXXX)

Rev. xx/xx/xx xxxx

**Net Weight: XX lb (XX kg)**

EPA REG. NO. 10404-82

EPA EST. NO. 10404-FL-1(S), 10404-OH-4(M)

First letter of lot code indicates manufacturing site.

**#XXXXXXXX**

FXXXX

Manufactured by: LESCO, Inc.

15885 Sprague Road  
Strongsville, Ohio 44136-1772

758073xxxxx

EPA Reg. No.: 10404-82  
 Product Name: LESCO Pre-M 1.31% Plus Fertilizer  
 Date: 7/8/2004

**Alternative Fertilizer Percentages**

The following information is to be made a part of the file for this product since it involves the alternate Fertilizer Fillers and Conditioners which will be utilized for this product as based upon the need for certain amounts of soil nutrients as indicated by a "Soil Test" supplied by the Customer or Market.

Product	CAS Number	Percentage	Source
Methylene urea	5/6/9011	0 - 98	KPAC Holdings (Disputanta, VA), Nu-Gro (Courtright, Ontario)
Sulfur Coated Urea	7704-34-9	0 - 98	LESCO, Inc. (Strongsville, OH); Nu-Gro (Courtright, Ontario)
Urea	57-13-6	0 - 98	CF Industries (Long Grove, IL); PCS Sales (Northbrook, IL)
Ammonium Sulfate	57-13-6	0 - 95	DSM Chemical (Augusta, GA); APF (Galina Park, TX)
Diammonium Phosphate	7783-20-2	0 - 95	CF Industries (Long Grove, IL); PCS Sales (Northbrook, IL)
Limestone	7783-28-0	0 - 95	National Lime & Stone (Findlay, OH); Franklin Minerals (Nashville, TN); Cargill Fertilizer (Riverview, FL)
Limestone	1317-65-3	0 - 95	National Lime & Stone (Findlay, OH); Franklin Minerals (Nashville, TN); Cargill Fertilizer (Riverview, FL)
Monoammonium Phosphate	14808-60-1	0 - 95	IMC Global (Lake Forest, IL)
Clay	7722-76-1	0 - 90	Englehard Minerals (Iselin, NJ)
Corn Cob Granules	NA	0 - 90	Timberlake Sales (Mt. Pulaski, IL); IBCP (Independence, IA)
Sulfur	NA	0 - 30	Sulfur Coated Urea (LESCO, Inc, Strongsville, OH); Ammonium Sulfate (DSM Chemical, (Augusta, GA); APF (Galina Park, TX)
Potassium Nitrate	7704-34-9	0 - 20	Hydro Agri (Tampa, FL)
Sodium Bentonite	7757-79-1	0 - 20	Bennett Minerals (Walheston, VA)
Sulfate of Potash	1302-78-9	0 - 20	PICC (New York City, NY); Great Salt Lake Minerals (Overland Park, KS)
Biosolids	7778-80-5	0 - 10	New England Fertilizer
Iron (Ferric) Oxide	NA	0 - 10	Micronutrient Mixture (Frit Ind., Ozark, AL)
Iron (Ferrous) Oxide	1309-37-1	0 - 10	Granusol 50% Iron (American Minerals, King of Prussia, PA)
Iron (Ferrous) Sulfate	8047-67-4	0 - 10	Micronutrient Mixture (Frit Ind., Ozark, AL)
Magnesium Carbonate	7720-78-7	0 - 10	Sulfate of Potash-Magnesium (IMC Global, Northbrook, IL)
Magnesium Oxide	39409-82-0	0 - 10	Granusol 45% Magnesium (American Minerals, King of Prussia, PA)
Magnesium Sulfate	1309-48-4	0 - 10	Sulfate of Potash-Magnesium (IMC Global, Northbrook, IL)
Manganese Compounds	7478-88-9	0 - 10	Granusol 45% Magnesium (American Minerals, King of Prussia, PA)
Manganese Oxide	7439-96-5	0 - 10	Sulfate of Potash-Magnesium (IMC Global, Northbrook, IL)
Muriate of Potash	1317-35-7	0 - 10	Granusol 45% Magnesium (American Minerals, King of Prussia, PA)
Nitroform UF	7447-40-7	0 - 10	Granusol 35% Manganese (American Minerals, King of Prussia, PA)
Potassium Magnesium Sulfate	5/6/9011	0 - 10	Granusol 35% Manganese (American Minerals, King of Prussia, PA)
Urea Formaldehyde Concentrate	14977-37-8	0 - 10	PCS Sales (Northbrook, IL); IMC Global (Lake Forest, IL)
Calcium Silicate	50-00-0	0 - 5	Nu-Gro (Courtright, Ontario);
Manganous Oxide	1344-95-2	0 - 3	IMC Global (Lake Forest, IL)
Phosphoric Acid	1344-43-0	0 - 3	Borden Chemical (Springfield, OR)
M-Pyrol	7664-38-2	0 - 1	J. M. Huber (Havre de Grace, MD); Celite Corp. (Lompoc, CA)
DCA 11 Paraffinic Oil	872-50-4	0 - 1	Chemetals (Baltimore, MD)
Copper	64741-89-5	0 - 1	PCS Sales (Northbrook, IL)
Zinc	7440-50-8	0 - 1	ISP Technologies (Wayne, NJ)
CHLORINE	1314-13-2		ISP Technologies (Wayne, IN)

01/8

9/10

Optional Marketing Claims

**PRE-EMERGENT WEED CONTROL  
For Use on Turfgrasses**

- Preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed, and Poa annua
- Provides quick green-up from soluble nitrogen.
- Prevents annual grassy and broadleaf weeds.

**• PROVEN TECHNOLOGY**

- (insert number)% Poly Plus® for controlled spring growth and extended color
- Contains pendimethalin for preemergent weed control

**• GREATER PERFORMANCE AND SAVINGS**

- Free-flowing product makes (insert fertilizer grade) easy to apply
- A cost effective solution providing wide spectrum weed control - crabgrass, foxtail, goosegrass
- Controls crabgrass and other annual weeds
- Contains potassium for plant hardiness and drought tolerance
- Removing particles will help prevent staining

**(Brand) ANNUAL TURF FERTILIZER PROGRAM**

**Step 1 Apply Late Winter-Early Spring**

**(Brand) Crabgrass Preventer + Lawn Fertilizer**

Prevents crabgrass, foxtail, poa annua, spurge and other annual grasses while feeding your lawn with long-lasting, deep greening controlled-release nitrogen,

**Step 2 Apply Mid-Late Spring**

**(Brand) Weed & Feed\***

Kills over 100 species of weeds, roots and all! Feeds with controlled-release nitrogen for longer lasting greening

**Step 3 Apply Late Spring-Summer**

**(Brand) Lawn Insect Control with Fertilizer**

Controls insects while greening your lawn with controlled-release nitrogen for dependable, predictable results!

**Step 4 Apply Late Summer-Early Fall**

**(Brand)-Lawn Fertilizer**

Provides extended greening throughout the summer months with controlled-released nitrogen for dependable, predictable results!

\* (Brand) St. Augustine Weed & Feed can be used as step 2 in your Annual Program ONLY on St. Augustine, and Centipede Lawns

**SATISFACTION GUARANTEED PANEL**

**(GUARANTEE SEAL)**

**SATISFACTION GUARANTEED OR YOUR MONEY BACK.** (Brand) is committed to providing the highest quality products. If you are not totally satisfied with our products, we will replace your product or refund your money. Just call us at 1-800-XXX-XXXX and we will be happy to help you.

**Start Using the (Brand) Annual Turf Fertilizer Program**

(Chart showing the time of year to use what products by agronomic zone)

Northern Zone

- Step 1 Late March - Early May
- Step 2 April 1 - Early June
- Step 3 Mid May-End of August
- Step 4 August 1- End of September

Transitional Zone

- Step 1 Mid March- Mid April
- Step 2 Late March - End of May
- Step 3 Mid May-Early September
- Step 4 First of September - Late October

Southern Zone

- Step 1 Mid January- Mid April
- Step 2 Mid March-Mid May
- Step 3 Late April-End of September
- Step 4 Mid August - Mid November

(Graphic of the map showing where the various zones are located)

***This product can cause staining. To prevent staining and/or unwanted tracking of yellow particles do not apply this product to moist turf. After application, brush or sweep accidentally applied materials off all clothing, shoes, cement, stone, metallic surfaces (automobiles or garden furniture). If needed, rinse off with water.***

***This product can cause staining. Do not track product onto linoleum flooring or carpeting. To help prevent movement of this product off-site and/or staining, do not apply this product to moist turfgrass. After application but before initial irrigation, brush or sweep particles off all surfaces including cement, stone, metal, pools, decks, etc. After initial watering, keep off treated areas until dry.- Wash clothing used for applying this product separately. Air dry outdoors.***