

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460 EPA Reg. Number:

Date of Issuance:

10404-82

07/03/07

NOTICE OF PESTICIDE:

__ Registration
__X Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

LESCO Pre-M 1.31% Plus Fertilizer

Name and Address of Registrant (include ZIP Code):

Mr. John Olizinger Lesco, Inc. 1301 East 9th St., Suite 1300 Cleveland, OH 44114-1849

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The label described above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This product is re-registered in accordance with FIFRA sec. 4(g)(2)(C). Reregistration under this section of the Act does not eliminate the need for continual assessment of the registration of this product.

This labeling supersedes all previous accepted labeling for this product.

A copy of the label stamped accepted is enclosed for your records. Please submit one copy of the final printed labeling when product is released for shipment.

Signature of Approving Official:
James A. Tompkins
Product Manager 25
Herbicide Branch

Registration Division (7505R)

Date

7/3/07

EPA Form 8570-6

•PRE-M[®] 1.31% Plus Fertilizer

Not for use on turf being grown for sale or for other commercial as sod, or for commercial seed production, or for research /0ses

For control of specified weeds in or on residential lawns, corporate lawns, parks, recreational areas, golf courses, cemeteries, and

- Preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalls. spurge, knotweed, and Poa annua.
- · Provides quick green-up from soluble nitrogen.
- · Prevents annual grassy and broadleaf weeds.
- · Contains LESCO Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding.
 Contains LESCO® Poly Plus® Sulfur Coated Urea to provide a uniform growth

with extended nitrogen feeding.

Active Ingredient

, thurst surgicular	
Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-	
2.6-dinitrobenzenamine*	1.31%
Other ingredients:	98.69%
Total	100.00%
1 icensed under one or more of the following U.S. Patenta: 3 920 742: 4 066 44	1.



		 	200100
HAR	ΔNI	 ANA	YSIS

GUARANTEED ANALYSIS	
TOTAL NITROGEN (N)	XX.XX%
XX.XX% Ammoniacal Nitrogen	
XX.XX% Urea Nitrogen*	
XXXX% Water Insoluble Nitrogen	
X.XX% Water Soluble Controlled Release Nitrogen	
F. JLABLE PHOSPHATE (P₂O₅)	XX.XX%
SOLUBLE POTASH (K20)	XX.XX%
MAGNESIUM (Mg) Total	XX.XX%
XX.XX% Water Soluble Magnesium	
SULFUR (S) Total	XX.XX%
XX.XX% Free Sulfur (\$)	
XX.XX% Combined Suffur (S)	
COPPER (Cu) Total	XX.XX%
XX,XX% Water Soluble Copper (Cu)	
IRON (Fe) Total	XX.XX%
XX.XX% Water Soluble Iron (Fe)	
MANGANESE (Mn) Total	XX.XX%
XX.XX% Water Soluble Manganese	
ZINC (Zn) Total	XX.XX%
XX.XX% Water Soluble Zinc (Zn)	
DERIVED FROM:	
CHLORINE (CI) Max.	
*X.XX% Slowly available Urea Nitrogen from Polymer Coated Sulfur C	oated Urea.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, ekin, or clothing. Harmful if absorbed through skin. Wash theroughly with seap and water after handling. Avoid breathing dust

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID			
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then		
	continue rinsing. Call a polson control center or doctor for treatment advice.		
, illowed	 Call a poison control center or doctor immediately for treatment advice. 		
-	 Have person sip a glass of water if able to swallow. 		
	 Do not induće vomiting unless told to by the poison control center or doctor. 		
	 Do not give anything by mouth to an unconscious person. 		
If on skin or	Take off contaminated clothing.		

clothing	Rinse skin immediately with pienty of water for 15-20 minutes. Call a poison control center or doctor for treatment edvice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a polson control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof materials, such as butyl rubber > 14 mils, nitrile rubber >14 mils, or neoprene rubber > 14 mils, shoes plus socks. Follow manufacturers' instructions for changing cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE Immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove contaminated clothing and wash separately before reuse.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at time of pesticide
- Observe all cautions and limitations in this leaflet and on the labels of products used in combination with this product. Do not use this product other than in accordance with the instructions set forth on this label. The use of this product not consistent with this label can result in injury to turf, animals, or persons. Keep container closed to avoid spills and contamination.
- DO NOT apply this product through any type of irrigation system.
- Do not enter or allow others to enter the treated area until dusts have settled.
- Do not apply this product in a way that will contact workers, other any persons, or pets, either directly or by through drift. Only protected handlers may be in the area during application.
- Do not allow dust to drift from the application site and contact people, structures people occupy at the time and associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rengelands, or animais.

DO NOT apply more than 3.5 lb of this product per 1,000 sq ft per application, and no more than 10.5 lb of this product per 1,000 sq ft per year.

GENERAL INFORMATION

This product is a combination pre-emergent herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in turfgrass. This product is labeled for use on the following turfgrass species: Kentucky Bluegrass, Perennial Ryegrass, Fine Fescue, Tall Fescue, Bentgrass, Poa annua (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, and Zoyslagrass.

This product controls weeds as they germinate but will not control established weeds. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control

Applied according to label directions and under normal growing conditions, this product will not cause turf injury. Over-application can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil saft concentration, or drought can weaken turfgrass and increase the possibility of turf damage from this product.

APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated appropries the spirit of the s prevent staining.

> Under the Federal Innecticide, Fungicide, and Rodenticide Act as amended, for the posticide registered under EPA Reg. No.



10404-82

PRE-M® 1.31% Plus Fertilizer

·			
CONTRACTOR DATES.	DATE AND	LAPPI ICATION	CHART

		E AND APPLICATION Rates of 1.31%	Comments:
Turfgrass Species	Weeds	PRE-M Plus Fertilizer	Comments:
NORTHERN GRASSES Kentucky Bluegrass Perennial Ryagrass Fine Fescue Tall Fescue	crebgrase foxiall Poe annue barnyardgrass fall panicum oxalle prosurate spurge purslane	2.8 to 3.5 lb/1,000 eq ft (115 – 153 lb/acre) as the initial application prior to weed germination in the spring.	Apply a repeat application of 1.8 to 2.6 to/1,000 sq ft (78 – 115 to/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	knotweed evening primrose hop clover goosegrass	2,6 to 3.5 lb/1,000 eq ft (115 – 153 lb/ecre)	Apply a repeat
	cudweed	in the spring prior to weed germination. 2,6 to 3.5 lb/1,000 eq	ib/1,000 sq ft (115 lb/acre) Apply in late summer
	Pos annua chickweed lawn burweed henbit corn speedwell	ft (1.15 – 153 lb/acre)	or sarly fall prior to weed germination.
Bentgrass and established <i>Pas ennus</i> * (1/2 Inch height or taller)	crabgrase foxual Pos annua barnyardgrass fall panicum oxalls prostrate spurge purslane knotweed evening primrose hop clover	2.8 ib/1,000 eq it (115 lb/acre) in the epring prior to weed germination.	Apply a repeat application of 1.8 to 2.8 ib/1,000 sq ft (78 – 115 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	2.6 lb/1,000 sq ft (115 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 2.8 lb/1,000 sq ft (115 lb/acre) after a minimum of 4 weeks.
	cudweed Poa annus chickweed lawn burweed henbit com speedwell	2.6 lb/1,000 sq ft (115 lb/scre)	Apply in late summer or early fall prior to weed germination.
SOUTHERN GRASSES Bermudagras SI. Augustinegrass Bahiagrass Zoyalagrass Centipedagrass Tall Feacue	crabgrass foxtail Poe annua barnyardgrass fall panicum oxalis prostrate apurge pursiane knotweed evening primrose hop clover	2.8 to 3.5 ib/1,000 sq ft (115 – 153 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 1.8 to 2.6 lb/1,000 eq ft (78 – 115 lb/acre) after a minimum of 4 weeks if necessary.
	goosegrass	2.6 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre) prior to weed gemination in the spring. Apply a repeat application of 2.5 to 3.5 lb/1,000 sq ft (115 – 153 lb/acre) after a minimum of 4 weeks.	An additional application of 2.6 ib/1,000 sq ft (115 ib/acre) may be applied for extended goossprass control a minimum of 4 weeks after the second application.
	cudweed Pos snnus chickweed tawn burweed henbit com speedwell	2.6 to 3.5 lb/1,000 sq ft (115 = 153 lb/acre)	Apply in late summer or early fall prior to weed germination.

*Not for use on greens or tees.

**Rates do not apply to Bermudagrass greens and tees.

The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 14 days, erratic weed control may result.

SUGGESTED SPREADER SETTINGS

Recommended Rates in lb per 1,000 sq ft

Spreader	1.8 16	2.6 lb	3.5 lb
LESCO Rotary Calibration Gauge	#XX	#XX	#X
Cyclone® or Spyker®	Х	X	X
SCOTTS® R8-A	X	X	X
Lely®	Χii	ХII	Χii

IMPORTANT: Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured atorage area. Do not contaminate water, foodstuffs, feed or seed by storage or disnocal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call [1-800-CLEANUP] for disposal instructions, Never place unused product down any Indoor or outdoor drain.

outdoor drain.

CONTAINER DISPOSAL: Do not reuse bag. Dispose of emptied bag(s) in a senitary landfill approved for pasticide disposal, or by incineration.

SPILL: In case of splil, awasp up material and dispose of material according to "peaticide disposal" directions listed above.

PRECAUTIONS

Use only on well-established turfgrass with a dense and uniform atand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times.

On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application.

Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species.

Do not use on Bentgrass or Pos annua (greens and tees) or injury may occur.

Delay reseeding or winter overseeding treated turigrass for at least 2 months following application. Do not sprig turigrass for 3 months after application.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY
NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to aliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors which are beyond the control of LESCO, inc., All such risks shall be assumed by Buyer and Buyer agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer assumes the risk of any such use. LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall LESCO, Inc. be liable for any Incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ONBREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC., REPLACEMENT OF THE PRODUCT.

LESCO, Inc. offers this product, and Buyer accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-a1,com.

LESCO, Pre-M and Poly Plus are registered trademarks and the aweeping design is a trademark of LESCO Technologies, LLC, SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC, Lely is a registered trademark of C. Van Der Lely N.V.

G:\REGUL\WP\Regul Private\MstrLb/_ERIEVIEW ADDRE\$\$\Naw EPA Approved\Pre-M
1.31% formatted.doo (113004) Rev. 5/24/07

Net Weight: XX lb (XX kg)

EPA REG. NO. 10404-82
EPA EST, NO.82757-FL-001 (S), 82757-QH-001 (M)
First letter of lat code indicates manufacturing site.

F1560 Distributed by: LESCO, Inc. 1301 East 9th Street Cleveland, OH 44114-1849