10404-8	4	/	-11							1/2
	Uni Environmental	ited States	Agency	Form A		Registr Amend Other	ation	OPF		x <u>pires 2-28-9</u> 5 ar Number
	A	Application f	for Pesti	cide - Se	ction	I				
1. Company/Product Numb				A Product Ma	inager		3.	3. Proposed Classification		
10404-84			Joanne Miller					None	•	Restricted
4. Company/Product (Nam LESCO Dimension 0.			23				-		••••••	I
5. Name and Address of A LESCO, Inc. 1301 East 9th Stree Cleveland, OH 4411	t	9)	(b)(i), to:	my product	t is sim	ilar or iden	itical in	compos	ition and	
Check if th	is is a new address		Proc	duct Name						
			Section -	- 11			····			
Amendment - Expla	in below. sponse to Agency letter d		[	Final print Agancy la "Ma Too"	tter date			LIFICA	TION	
		ateu	[	<u>.</u>				DTIFICATION, UL 1 1 2005		
Notification - Explain	n below.		<b>/</b>	Other - Ex	plain be	low.	JU	[]]	2000	
1. Material This Product Wi	ill Re Packaged (n:	S	Section -	111						·
Child-Resistant Packaging	Unit Packaging	w	ater Soluble	Packaging		2. Type of	Contain			
Yos	Yes		Yes				Metal			
No	No		No			_ Plasti Glass	Glass			
* Certification must be submitted	lf "Yes" Unit Packaging wgt.		"Yes" eckage wgt	No. per containe			_] Paper _] Other	(Specify	1	
3. Location of Net Contents	s Information 4 Container	. Size(s) Retail Co	ontainer	l	5. Loc	ation of La	bel Direc	tions		· · · · · · · ·
6. Manner in Which Label is	s Affixed to Product	Lithograph Paper plued Stenciled	1	Othe	er				······	
		S	ection -	IV						
1. Contact Point (Complete	e items directly below for	identification of i	individual to	be contacted,	, if nece	ssery, to pi	rocess ti	nis epplic	etion.)	
Name Susan Spencer			Title Manager, Regulatory Affairs			Telephone No. (Include Area Code) 216-706-9250 ext 2343		trea Code)		
I certify that the state I acknowledge that a both under applicable	ements I have made on th ny knowlinglly false or mi blaw.	Certification his form and all at isleading statement	tachments ti nt may be pi	herato are tru unishable by f	ie, accu fine or ir	rete and com mprisonmen	inplace. It or	1	te Applic ceived \Stamp	
2. Signature		3. Tit	10							
Aucan Expension			Manager, Regulatory Affairs							
4. Typed Name		5. Dat	te		·		•			
Susan E. Spencer			4/,	13,06						

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

statements are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

## KEEP OUT OF REACH OF CHILDREN CAUTION

	FIRST AID
If swallowed	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> </ul>
Swallowcu	
	<ul> <li>Have person sip a glass of water if able to swallow.</li> </ul>
	<ul> <li>Do not induce vomiting unless told to by the poison control center or doctor.</li> </ul>
	<ul> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> </ul>
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
	Call a poison control center or doctor for treatment advice.
lf on skin	Take off contaminated clothing.
or clothing	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give
	artificial respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.
	HOT LINE NUMBER
Have the pro-	oduct container or label with you when calling a poison control center or

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTIONI CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

#### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses, and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

Common Name Bentgrass, creeping\* Bluegrass, Kentucky Fescue, fine\*\* Fescue, tall Ryegrass, perennial <u>Scientific Name</u> Agrostis palustris Poa pratensis Festuca rubra Festuca arundinacea Lolium perenne



Net Weight: XX lb (XX kg)

"Beauly', 'Biljart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivaio', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waklorf', and 'Waklina'. Not all varieties of Fine Fescue have been tested.

\*\*\*DO NOT use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

# DIMENSION\* TURFGRASS REGIONS



#### Reseeding, Overseeding, or Sprigging


Reseeding, overseeding or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practice such as soil cultivation, irrigation and fertilization should be followed.

However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application.

For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

#### USE PRECAUTIONS

- Not for use on Golf Course Putting Greens.
- The early post-emergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.
- Apply this product directly to established lawns or ornamental turfs only.
- · DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.



XX.XX% Water Soluble Magnesium	
SULFUR (S) total	XX,XX%
XX.XX% Free Sulfur (S)	
XX.XX% Combined Sulfur (S)	
COPPER (Cu) total	XX.XX%
XX.XX% Water Soluble Copper (Cu)	
IRON (Fe) total	XX.XX%
XX.XX% Water Soluble Iron (Fe)	
MANGANESE (Mn) total	XX.XX%
XX.XX% Water Soluble Manganese	
ZINC (Zn) total	XX.XX%
XX.XX% Water Soluble Zinc (Zn)	
DERIVED FROM: XXXX, XXXX, XXXX,	
CHLORINE (CI) Max	XX.XX%
*X.XX% Slowly available Urea Nitrogen from Polymer Coated Sulfur Coated Urea	

# **DIMENSION\* 0.07% Plus Fertilizer**

### **APPLICATION DIRECTIONS - Control of Crabgrass**

Pre-emergence and Early Post-emergence Control

This product provides "pre-emergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early post-emergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early post-emergence activity is that (compared to strictly pre-emergence crabgrass products), Dimension controls crabgrass prior to and up to 4 weeks after germination.

#### **Application Frequency and Timing**

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 16.39 lb of this product per 1,000 sq ft per application, and

no more than 49.17 lb of this product per 1,000 sq ft per year.

Spring Applications For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be pre-emergence or early post-emergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program #1 for pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabarass control

Use Program #2 for pre-emergence control at sites where (a) turfgrass is cut low (e.g. golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program #3 for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or post-emergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

#### Table 1: Recommended Single Application Use Rates\*

PROGRAM	#1	#2	#3
USE	Pre-emergence	Pre-emergence	Pre-emergence
(turfgrass cut)	(high-cut turf)	(low-cut turf)	(low-cut turf)
		Post-emergence	Post-emergence
REGION		(high-cut turf)	(low-cut turf)
North	4.09 lb/1,000 sq ft <sup>‡</sup>	5.09 lb/1,000 sq ft <sup>‡</sup>	8.19 lb/1,000 sq ft
Transition	5.09 lb/1,000 sq ft <sup>‡</sup>	8.19 lb/1,000 sq ft	12.13 lb/1,000 sq ft
South	8.19 lb/1,000 sq ft	12.13 lb/1,000 sq ft	16.39 lb/1,000 sq ft
Coastal South	12.13 lb/1.000 sq ft	16.39 lb/1.000 sq ft	16.39 lb/1 000 sq ft**

Note: 8.19 lb per 1,000 sq ft is equal to 0.25 lb active ingredient per acre.

\* Particle distribution at this rate may not provide adequate control or suppression. \*DO NOT apply more than 16.39 lb of this product per 1,000 sq ft per application, and no more than 49.17 lb of this product per 1,000 sq ft per year. Post-emergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. See map of the United States. North: all areas not designated below.

	all alcas hot designated below.
Transition:	DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of
	IL, IN, OH, & coastal areas of CT, NY, & RI.
South:	AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX. ' May require split or sequential applications for full control.

### Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

#### Tips for Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

EPA REG. NO. 10404-84

EPA EST. NO. 82757-OH-001 (M), 82757-FL-001 (S)



**PPLICATION DIRECTIONS - Control of** Other Grasses and Broadleaf Weeds

#### Spring Application

Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

barnyardgrass	ryegrass (annual & perennial)
bluegrass (annual)	smutgrass
crowfootgrass	lespedeza (common)
foxtail (vellow & green)*	oxalis (buttercup, creeping & yellow woodsorrel)
goosegrass	purslane (common)
kikuyugrass	speedwell (corn)
	source (prostrate & spotted)

\*Also controlled at the "Program 2" rates.

#### **Fall Applications**

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence: bluegrass (annual) geranium (Carolina) blue na)

Didegrass (annual)	geranium (Carolina
bittercress	parsley-piert
chickweed	pineappleweed
henbit	shepherd's purse

#### SUGGESTED SPREADER SETTINGS\*

	(Recommended Rates in lb per 1,000 sq ft)					
Spreader	4.09 lb	5.09 lb	8.19 lb	12.13 lb	16.39 lb	
LESCO Calibration Gauge	, XX	XX	XX	XX	XX	
SCOTTS <sup>®</sup> R8A	XX	XX	XX	XX	XX	
Cyclone <sup>®</sup> or Spyker <sup>®</sup>	XX	XX	XX	XX	XX	
LESCO Pendulum	XX	XX	XX	XX	XX	

\*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

#### **APPLICATION EQUIPMENT AND INSTRUCTIONS**

Apply this product with drop or rotary spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

#### STORAGE AND DISPOSAL

STORAGE: Store this product only in its original container in a dry, cool, secured storage a. Do not contaminate water, foodstuffs, feed or seed by storage or disposal. PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be

used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call or [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.

SPILL: In case of spill, sweep up material and dispose of material according to \*Product Disposal" directions listed above.

#### WARRANTY

LESCO. Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. LESCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of LESCO and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL LESCO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE, OR USE OF THIS PRODUCT.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9<sup>th</sup> Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-a1.com.

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Poly Plus is comprised of Polymer Coated Suffur Coated Ures. DIMENSION is a registered trademark of Dow AgroSciences. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Streagers, LLC.

G \REGUL\WP\Regul Private\WstrLbf\New EPA ApprovedDimension .07%.doc Created: 5/27/03 (020503) 84. 07 05 BS

Net Weight: XX lb (XX kg) ..... F1560

Distributed by: LESCO, Inc. 1301 East 9th Street

Cleveland, OH 44114-1849





Phone: 216.706.9250 800.321.5325 www.lesco.com Nasdaq: LESCO

June 15, 2006

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Attn: Ms. Joanne Miller, PM 23

SUBJECT:Notification -- Optional Language and Fertilizer StatementsEPA #10404-840.07% Dimension Plus FertilizerEPA #10404-85Dimension 0.10% Plus FertilizerEPA #10404-86Dimension 0.15% Plus Fertilizer (Commercial & Non-Commercial)EPA #10404-86LESCO Dimension Crabgrass Pre-Emergent Plus Fertilizer (ABN)EPA #10404-87LESCO Scenic Green Crabgrass Pre-Emergent + Lawn Fertilizer (ABN)EPA #10404-87Dimension 0.21% Plus Fertilizer (Commercial & Non-Commercial)

#### Dear Ms. Miller:

Notifications are being submitted for your review and acceptance to address the following:

- Addition to Optional/Marketing statements:
  - Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding."
  - Clarification of 'Derived From' statement in the Guaranteed (Fertilizer) Analysis box:
  - o "Slowly Available Urea Nitrogen from Polymer Coated sulfur Coated Urea"
  - Clarification of 'Poly Plus' composition in the Trademarks section:
    - o "Poly Plus is comprised of Polymer Coated sulfur Coated Urea"

The following documentation is provided in support of the above:

- 8570-1: Notification request form for each EPA registration number
- One (1) highlighted label of each EPA registration noting the above items
- Five (5) copies of the clean label for each EPA registration

If you have any questions, or further clarification is required, please feel free to contact me at <u>sspencer@lesco.com</u> or 1-216-706-9250 ext 2343.

Thank you for your assistance on this matter.

Sincerely,

Susan E. Spencer

Susan E. Spencer Manager Regulatory Affairs

Enclosures

- 10404-84 LESCO Dimension 0.07% Plus Fertilizer
- 10404-85 LESCO Dimension 0.10% Plus Fertilizer
- 10404-86 LESCO Dimension 0.15% Plus Fertilizer (commercial & non-commercial)
- 10404-86 LESCO Dimension Crabgrass Pre-Emergent Plus Fertilizer (ABN)
- 10404-86 LESCO Scenic Green Crabgrass Pre-Emergent + Lawn Fertilizer (ABN)
- 10404-87 LESCO Dimension 0.21% Plus Fertilizer (commercial & non-commercial)

Notification to address the following:

- Addition to Optional/Marketing statements:
  - "Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding:"
- Clarification of 'Derived From' statement in the Guaranteed (Fertilizer) Analysis box:
  - "Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea"
- Clarification of 'Poly Plus' composition in the Trademark section:
  - o "Poly Plus is comprised of Polymer Coated Sulfur Coated Urea"