

10404-85

2/5/2003

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

FEB - 5 2003

Gloria M. Sieloff, Ph.D.  
Director, Regulatory Affairs and Corporate Quality  
LESCO, Inc.  
15885 Sprague Rd.  
Strongsville, Ohio 44138

Dear Dr. Sieloff:

Subject: 0.10% Dimension® Plus Fertilizer  
EPA Registration No. 10404-85  
Application Dated July 25, 2002, and Letters Dated  
October 8, and 25, 2002, Request for an Alternate  
Confidential Statement of Formula (CSF) and To  
Revise Labeling as Amendments to Registration

The propose alternate (A001), Confidential Statement of Formula (CSF), submitted on EPA Form 8570-4 and dated October 25, 2002, has been reviewed and found as an acceptable amendment to the registration of "0.10% Dimension® Plus Fertilizer" under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). The alternate CSF has been placed in this Agency's file for the subject pesticide product registration.

The proposed labeling amendments reflected on the labeling dated January 17, 2003 has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

- o Submit one (1) copy of the final printed label prior to shipment of this pesticide product under the enclosed revised label.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA,

-2-

section 6(e). Your release for shipment under this labeling constitutes acceptance of this condition. A stamped copy of the label is enclosed for your records.

Sincerely Yours,

*Eugene M. Wilson*  
for Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Enclosure

## DIMENSION® 0.10% Plus Fertilizer

**NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

In New York State this product may only be used by commercial applicators and at no more than 500 lb (0.5 lb of active ingredient) per acre per year (or 11.5 lb product per 1000 ft sq per year) and is prohibited from use in Nassau and Suffolk Counties.

### ACTIVE INGREDIENT:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester ..... 0.10%

**INERT INGREDIENTS:** ..... 99.90%

Total ..... 100.00%

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

Read the entire label before using this product. Use only according to label instructions.

**NOTICE:** Before using this product, read the Use Precautions, Warranty Statements, Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### FIRST AID

If swallowed	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION! CAUSES EYE IRRITATION. HARMFUL IF INHALED.** Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**#XXXXXX** (Front)      **Net Weight: XX lb (XX kg)**

### GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses, and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

##### Common Name

Bentgrass, creeping\*  
Bluegrass, Kentucky  
Fescue, fine\*\*  
Fescue, tall  
Ryegrass, perennial

##### Scientific Name

*Agrostis palustris*  
*Poa pratensis*  
*Festuca rubra*  
*Festuca arundinacea*  
*Lolium perenne*

#### Warm-Season Grasses

##### Common Name

Bahiagrass  
Bermudagrass  
Buffalograss\*\*\*  
Carpetgrass  
Centipedegrass  
Kikuyugrass  
St. Augustinegrass  
Zoysiagrass

##### Scientific Name

*Paspalum notatum*  
*Cynodon dactylon*  
*Buchloe dactyloides*  
*Axonopus affinis*  
*Eremochloa ophiuroides*  
*Pennisetum clandestinum*  
*Stenotaphrum secundatum*  
*Zoysia japonica*


DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.

\*Use of this product on certain varieties of Creeping Bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of Creeping Bentgrass have been tested.

\*\*Use of this product on certain varieties of Fine Fescue may result in undesirable turf injury. The following Fine Fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bijart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Kokel', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tajana', 'Waldorf', and 'Waldina'. Not all varieties of Fine Fescue have been tested.

\*\*\*DO NOT use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

### DIMENSION® TURFGRASS REGIONS



### Reseeding, Overseeding, or Sprigging


Reseeding, overseeding or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practice such as soil cultivation, irrigation and fertilization should be followed.

However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application.

For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

**USE PRECAUTIONS**

- Not for use on Golf Course Putting Greens.
- The early post-emergence crabgrass control described below is limited. Post-emergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.
- Apply this product directly to established lawns or ornamental turfs only.
- DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.



**(OPTIONAL LANGUAGE: MINI, ELITE, NOVEX, PROFESSIONAL TURF FERTILIZER)  
GUARANTEED ANALYSIS**

TOTAL NITROGEN (N) .....	XX.XX%
XX.XX% Ammoniacal Nitrogen	
XX.XX% Urea Nitrogen*	
XX.XX% Water Insoluble Nitrogen	
XX.XX% yyyyyy	
AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> ) .....	XX.XX%
SOLUBLE POTASH (K <sub>2</sub> O) .....	XX.XX%
MAGNESIUM (Mg) total .....	XX.XX%
XX.XX% Water Soluble Magnesium	
SULFUR (S) total .....	XX.XX%
XX.XX% Free Sulfur (S)	
XX.XX% Combined Sulfur (S)	
COPPER (Cu) total .....	XX.XX%
XX.XX% Soluble Copper (Cu)	
IRON (Fe) total .....	XX.XX%
XX.XX% Soluble Iron (Fe)	
MANGANESE (Mn) total .....	XX.XX%
XX.XX% Water Soluble Manganese	
ZINC (Zn) total .....	XX.XX%
XX.XX% Soluble Zinc (Zn)	
DERIVED FROM: XXXX, XXXX, XXXX, ...	
CHLORINE (Cl) MAX. ....	XX.XX%
*XX.XX% Slowly available Nitrogen from LESCO® Poly Plus® Sulfur Coated Urea.	

**APPLICATION DIRECTIONS - Control of Crabgrass  
Pre-emergence and Early Post-emergence Control**

This product provides "pre-emergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early post-emergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early post-emergence activity is that (compared to strictly pre-emergence crabgrass products), Dimension controls crabgrass prior to and up to 4 weeks after germination.

**Application Frequency and Timing**

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 11.47 lb of this product per 1,000 sq ft per application and no more than 34.41 lb of this product per 1,000 sq ft per year.

**Spring Applications**

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be pre-emergence or early post-emergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

**Use Program #1** for pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabgrass control.

**Use Program #2** for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

**Use Program #3** for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or post-emergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

**Table 1: Recommended Single Application Use Rates\***

PROGRAM	#1	#2	#3
USE (turfgrass cut)	Pre-emergence (high-cut turf)	Pre-emergence (low-cut turf) Post-emergence (high-cut turf)	Pre-emergence (low-cut turf) Post-emergence (low-cut turf)
REGION			
North	2.86 lb/1,000 sq ft <sup>†</sup>	4.13 lb/1,000 sq ft <sup>†</sup>	5.73 lb/1,000 sq ft
Transition	4.13 lb/1,000 sq ft <sup>†</sup>	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft
South	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft
Coastal South	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft	11.47 lb/1,000 sq ft**

Note: 5.73 lb per 1,000 sq ft is equal to 0.25 lb active ingredient per acre.  
<sup>†</sup> Particle distribution at this rate may not provide adequate control or suppression.  
 \*DO NOT apply more than 11.47 lb of this product per 1,000 sq ft per application, and no more than 34.41 lb of this product per 1,000 sq ft per year. Post-emergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. See map of the United States.  
 North: all areas not designated below.  
 Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.  
 South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.  
 Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.  
 \*\* May require split or sequential applications for full control.

**Fall Applications**

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

**Tips for Improved Control**

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.  
 Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

4/6

**APPLICATION DIRECTIONS - Control of  
Other Grasses and Broadleaf Weeds**

**Spring Application**

Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

- |                           |  |
|---------------------------|--|
| barnyardgrass             | ryegrass (annual & perennial)                    |
| bluegrass (annual)        | smutgrass  |
| crowfootgrass             | lespedeza (common)                               |
| foxtail (yellow & green)* | oxalis (buttercup, creeping & yellow woodsorrel) |
| goosegrass                | purslane (common)                                |
| kikuyugrass               | speedwell (corn)                                 |
|                           | spurge (prostrate & spotted)                     |

\*Also controlled at the "Program 2" rates.

**Fall Applications**

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

- |                    |                     |
|--------------------|---------------------|
| bluegrass (annual) | geranium (Carolina) |
| bittercress        | parsley-piert       |
| chickweed          | pineappleweed       |
| nenbit             | shepherd's purse    |

**SUGGESTED SPREADER SETTINGS\***

Spreader	(Recommended Rates in lb per 1,000 sq ft)				
	2.86 lb	4.13 lb	5.73 lb	8.49 lb	11.47 lb
aaa	XX	XX	XX	XX	XX
bbb	XX	XX	XX	XX	XX
ccc	XX	XX	XX	XX	XX
ddd	XX	XX	XX	XX	XX
fff	XX	XX	XX	XX	XX
ggg	XX	XX	XX	XX	XX
hhh	XX	XX	XX	XX	XX
iii	XX	XX	XX	XX	XX

\*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

**APPLICATION EQUIPMENT AND INSTRUCTIONS**

Apply this product with drop or rotary spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or clumps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

EPA REG. NO. 10404-85  
EPA EST. NO. 10404-OH-4 (M),  
10404-FL-1 (S), 59197-CA-1 (C)

First letter of lot code indicates manufacturing site.

FXXXX

Manufactured by: **LESCO, Inc.**  
15885 Sprague Road  
Strongsville, OH 44136-1799

758073xxxxx

**#XXXXXXXX** (Back)

5/6

**STORAGE AND DISPOSAL**

**STORAGE:** Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PRODUCT DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call or [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

**EMPTY CONTAINER DISPOSAL:** Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.

**SPILL:** In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

**WARRANTY**

LESCO, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. LESCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of LESCO and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL LESCO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE, OR USE OF THIS PRODUCT.**

Information concerning the raw materials composing this product can be obtained by writing to LESCO, Inc., Attn. RA Dept., 15885 Sprague Rd Strongsville, OH 44136-1799, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-a1.com>.

LESCO and Poly Plus are registered trademarks of LESCO Technologies, L.L.C.

DIMENSION is a registered trademark of DowAgrosciences

G:\REGUL\Product Labels\Masters\EPA Documents\Dimension010.clean.DOC Created xx/xx/xx (XXXXXX)

Rev xx/xx/xx xxxx

**Net Weight: XX lb (XX kg)**

Optional Language:

6/6

# Weed Control with Premium Lawn Food

DIMENSION 0.10% Plus Fertilizer is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

Dimension controls crabgrass prior to or after germination.

Provides up to 4 months crabgrass control.

Provides both pre- and post-emergence crabgrass control.

**ACCEPTED  
with COMMENTS  
in EPA Letter Dated**

**FEB - 5 2003**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

10404-85