

10404-88

7-11-2006

1/5

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

 Environmental Protection Agency United States Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 10404-88	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) LESCO Stonewall 0.20% Plus Fertilizer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) LESCO, Inc. 1301 East 9th Street Cleveland, OH 44114-1849 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input checked="" type="checkbox"/> Other - Explain below.
---	--

NOTIFICATION
JUL 11 2006


Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Optional Marketing and Fertilizer Statements
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Susan Spencer	Title Manager, Regulatory Affairs	Telephone No. (Include Area Code) 216-706-9250 ext 2243
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Manager, Regulatory Affairs	
4. Typed Name Susan E. Spencer	5. Date 6/13/06	

	<ul style="list-style-type: none"> • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.

SPILL: In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

DIRECTIONS FOR USE

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns; landscape ornamentals; and established perennials & wildflower plantings. This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application.
- **Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system.
- Do not apply aerially.
- Do not apply to golf course putting greens.

USE PRECAUTIONS

The following use precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over-seeded grasses.
- Do not apply to overseeded turf for 60 days after seeding or until after the second mowing, which ever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil.
- Do not cut treated sod before 120 days after application. Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.


Available Nitrogen (N) Total	XX.XX%
Soluble Potash (K ₂ O)	XX.XX%
Magnesium (Mg) Total	XX.XX%
XX.XX% Water Soluble Magnesium (Mg)	
Sulfur (S) Total	XX.XX%
XX.XX% Free Sulfur (S)	
XX.XX% Combined Sulfur (S)	
Copper (Cu) Total	XX.XX%
XX.XX% Water Soluble Copper (Cu)	
Iron (Fe) Total	XX.XX%
XX.XX% Water Soluble Iron (Fe)	
Manganese (Mn) Total	XX.XX%
XX.XX% Water Soluble Manganese (Mn)	
Zinc (Zn) Total	XX.XX%
XX.XX% Water Soluble Zinc (Zn)	
DERIVED FROM:	
CHLORINE (Cl) MAX.	XX.XX%
*X.XX% Slowly available Urea Nitrogen from Polymer Coated Sulfur Coated Urea.	

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated commercial application equipment.

ESTABLISHED TURF

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the information below for approximate crabgrass seed germination dates.



Region	Date
Upper North	After May 10
North	After April 20
Transition	After March 20
South	Jan 1 - March 20

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed below. Do not exceed the maximum yearly rate:

Turf Species	Maximum Application Rate per Calendar Year*	
	lb LESCO Stonewall 0.20% Plus Fertilizer / acre	lb AI per Acre
Bermudagrass ¹ Bahia grass Centipedegrass Seashore Paspalum St. Augustinegrass ² Tall Fescue (including turf-type) Zoysia	750 lb/acre (17.22 lb/1,000 sq ft)	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	500 lb/acre (11.48 lb/1,000 sq ft)	1.0
Creeping Red Fescue	375 lb/acre (8.60 lb/1,000 sq ft)	0.75
Creeping Bentgrass	330 lb/acre (7.57 lb/1,000 sq ft)	0.65

1. These are the maximum rates per calendar year by species limitations.
2. May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 280 lb/Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.
3. Use at an initial rate of 190 to 375 lb/acre per application followed by sequential applications at doses that would not exceed the maximum annual application rate of 750 lb/acre/year.

Do not apply more than 750 lb of this product per acre per calendar year. Use the higher rates of this product to achieve higher levels of fertility and weed control for each turf type.

SUGGESTED SPREADER SETTINGS*

Spreader	Recommended Rates in lb per 1,000 sq ft			
	4.40 lb	5.70 lb	7.57 lb	8.60 lb
LESCO Calibration Gauge	#XX	#XX	#XX	#XX
SCOTT'S® R8A	XX	XX	XX	XX
Cyclone® or Spyker®	XX	XX	XX	XX
LESCO Pendulum	XX	XX	XX	XX

*IMPORTANT: These settings are only approximate. Age, condition of the spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

NOTIFICATION

JUL 11 2006

#XXXXXX

Stonewall™ 0.20% Plus Fertilizer

3/5

APPLICATION TIMING AND INFORMATION

This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil. This product may be applied at any time to established plants. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand-weeding, cultivation, or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most established ornamental trees, shrubs, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species without injury.

Scientific Name	Common Name	Scientific Name	Common Name
<i>Abies</i> spp.	Fir Species, Balsam, etc.**	<i>Malosiphora luteola</i>	Ice Plant
<i>Acer palmatum</i>	Japanese Maple	<i>Malus</i> sp.	Crabapple*
<i>A. platanoides</i>	Norway Maple	<i>Nandina domestica</i>	Heavenly Bamboo
<i>Actinidia chinensis</i>	Kiwi* (Ornamental Only)	<i>Olea europaea</i>	Olive* (Ornamental Only)
<i>Agapanthus africanus</i>	Lily-of-the-Nile	<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita	<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Arctotheca calendula</i>	Cape Weed	<i>Oxydendrum arboreum</i>	Sourwood
<i>Aucuba japonica</i>	Japanese Aucuba	<i>Persea americana</i>	Avocado*
<i>Barberis gladwynensis</i>	Barberry	<i>Photinia fraseri</i>	Frasier's Photinia (Redtip)
<i>B. julianae</i>	Wintergreen Barberry	<i>Picea</i> spp.	Colorado Blue, Norway, etc.
<i>B. mentorensis</i>	Mentor Barberry	<i>Peris japonica</i>	Lily-of-the-Valley Shrub
<i>B. thunbergii</i>	Japanese Barberry	<i>Pinus brutia</i>	Calabrian Pine
<i>B. verticillata</i>	Warty Barberry	<i>P. canariensis</i>	Canary Island Pine
<i>Buxus microphylla</i>	Japanese Boxwood	<i>P. elliptica</i>	Slash Pine
<i>Callistemon viminalis</i>	Weeping Bottlebrush	<i>P. halapensis</i>	Aleppo Pine
<i>Calluna vulgaris</i>	Scotch Heather	<i>P. nigra</i>	Austrian Black Pine
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)	<i>P. radiata</i>	Monterey Pine
<i>Cassia artemisoides</i>	Feathery Cassia	<i>P. palustris</i>	Longleaf Pine
<i>Ceanothus rigidus</i>	Wild Lilac	<i>P. strobus</i>	Eastern White Pine
<i>Chamaecyparis pisifera</i>	False Cypress	<i>P. sylvestris</i>	Scotch Pine
<i>Cleyera japonica</i>	Cleyera	<i>P. taeda</i>	Loblolly Pine
<i>Citrus</i> spp.*	Citrus Species*	<i>P. thunbergiana</i>	Japanese Black Pine
<i>Cornus florida</i>	Flowering Dogwood	<i>Pistachio</i> sp.	Pistachio*
<i>C. stolonifera</i>	American Dogwood	<i>Pittosporum thombifolium</i>	Queenland Pittosporum
<i>Cortaderia selloana</i>	Pampas Grass	<i>P. tobira</i>	Japanese Pittosporum
<i>Cotoneaster spiculatus</i>	Cranberry Cotoneaster	<i>Podocarpus macrophylla</i>	Podocarpus sp.
<i>C. buxifolius</i>	Cotoneaster	<i>Prunus laurocerasus</i>	English Laurel
<i>C. dammeri</i>	Bearberry Cotoneaster	<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>C. microphyllus</i>	Rockspray Cotoneaster	<i>Pseudotsuga menziesii</i>	Douglas Fir (Orna. Sites)
<i>Cretaegus</i> spp.	Hawthorne*	<i>Pyraecantha coccinea</i>	Firethorn (Scarlet)
<i>Cupressus sempervirens</i>	Italian Cypress	<i>P. fortuneana; koidzumii</i>	Firethorn spp.
<i>Delosperma alba</i>	White Trailing Ice Plant	<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Dodonea viscosa</i>	Hop Bush	<i>Quercus</i> spp.	Oak spp.
<i>Eleaagnus pungens</i>	Silverberry	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Euonymus fortunei</i>	Wintercreeper	<i>Rhododendron (including azalea)</i>	'Coral Bells', 'Formosa', 'Hino-crimson', 'IPJM', 'Roseum Elegans'
<i>E. japonica</i>	Evergreen Euonymus	<i>Rosa banksiae</i>	Lady Bank's Rosa
<i>E. kiautschovica</i>	Spreading Euonymus	<i>Rosmarinus officinalis</i>	Rosemary
<i>Fatsia japonica</i>	Japanese Aralia	<i>Santolina virens</i>	Santolina
<i>Forsythia intermedia</i>	Border Forsythia	<i>Sedum album</i>	Stonecrop
<i>F. viridissima</i>	Greenstem Forsythia	<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Gardenia jasminoides</i>	Gardenia, Cape Jasmine	<i>Taxus cuspidata</i>	Japanese Yew
<i>Hedera helix</i>	English Ivy	<i>T. media</i>	Yew, Var.
<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus	<i>Thuja occidentalis</i>	American Arborvitae
<i>Ilex cornuta</i>	Chinese Holly	<i>Trechelosperrum asiaticum</i>	Star Jasmine
<i>I. crenata</i>	Japanese Holly	<i>Tsuga canadensis</i>	Common Hemlock
<i>I. opaca</i>	American Holly	<i>Viburnum japonicum</i>	Japanese Viburnum
<i>I. pernyi</i>	Holly	<i>V. odoratissimum</i>	Sweet Viburnum
<i>Jasminum nudiflorum</i>	Winter Jasmine	<i>V. plicatum</i>	Japanese Snowball
<i>Juglans</i> sp.	Walnut*	<i>V. rigidum</i>	Canary Island Viburnum
<i>Juniperus chinensis</i>	Chinese Juniper	<i>V. tinus</i>	Laurustinus
<i>J. conferta</i>	Shore Juniper	<i>V. trilobium</i>	Cranberry Bush
<i>J. horizontalis</i>	Creeping Juniper	<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Justica brandegeana</i>	Shrimp Plant	<i>Vinca minor</i>	Dwarf Periwinkle
<i>Lagerstromia indica</i>	Crape Myrtle	<i>Vitis</i> sp.	Grape*
<i>Ligustrum amurense</i>	Amur Privet	<i>Weigela florida</i>	Old Fashioned Weigela
<i>L. japonicum</i>	Japanese Privet	<i>Yucca aloifolia</i>	Spanish Bayonet
<i>L. lucidum</i>	Glossy Privet	<i>Yucca filamentosa</i>	Yucca, Adam's Needle
<i>Lunope muscari</i>	Big Blue Lily Turf		

*Ornamental species only. Do not use on food producing trees or vines.
**For use on landscape ornamental sites

WEEDS CONTROLLED

When used in accordance with this label, this product will provide control of the following weeds:

Barnyardgrass	Goosegrass ¹	Lovegrass	Speedwell
Bluegrass, Annual (<i>Poa annua</i>)	Henbit	Panicum (Texas, Fall, Brown Top)	Sprangletop
Carpetweed	Itchgrass	Pigweed	Spurge, Prostrate
Chickweed, Common, Mouseear	Johnsongrass (from seed)	Purslane, Common	Witchgrass
Crabgrass (large, smooth)	Junglerice	Pusley, Florida	Woodsorrel, Yellow (from seed)
Crowfoot grass	Knotweed	Rescuegrass ²	
Cupgrass, Woolly	Kochia	Shepherd's Purse ³	
Foxtails, Annual	Lambsquarter, Common	Signalgrass, Broadleaf	

- In many areas a single application of 330 to 750 lb/acre of this product will control the above listed weeds and goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control can be obtained by making an initial application of 330 to 500 lb/acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.
- Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of weed seeds.
- Applications for this weed should be made in late summer, fall or winter prior to germination.

WHEN TO APPLY AFTER OVERSEEDING

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Product	LESCO Stonewall 0.20% Plus Fertilizer Rate (lb/Acre)	Interval Before Overseeding		
		North	Transition	South
250 lb/acre	0.50 lb/acre	4 months	4 months	4 months
330 lb/acre	0.66 lb/acre	5 months	4 months	4 months
370 lb/acre	0.74 lb/acre	6 months	5 months	5 months
400 lb/acre	0.80 lb/acre	-	6 months	6 months
500 lb/acre	1.00 lb/acre	-	7 months	7 months
570 lb/acre	1.14 lb/acre	-	-	9 months
650 lb/acre	1.30 lb/acre	-	-	10 months

LANDSCAPE ORNAMENTALS

(Including Established Perennial and Wildflower Plantings)

This product may be applied for residual pre-emergence weed control in ornamentals.

Use Rates - Apply this product at 330 to 750 lb/acre in fall and/or spring. Use the higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 750 pounds, per acre, per year.

WARRANTY

LESCO, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. LESCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of LESCO and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL LESCO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE, OR USE OF THIS PRODUCT.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-01.com>

LESCO and Poly Plus are registered trademarks and Stonewall and the sweeping design are trademarks of LESCO Technologies, LLC. Poly Plus is comprised of Polymer Coated Sulfur Coated Urea. SCOTTS is a registered trademark of The SCOTTS Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.

G:\REGUL\WP\Regul Private\MstrLb\New EPA Approved\Stonewall 0.20% doc
Created: 6/12/03 (061103) Rev.: 11/11/05

Net Weight: XX lb (XX kg)

EPA REG. NO. 10404-88

EPA EST. NO. 82757-FL-001(S), 82757-OH-001 (M)

First letter of lot code indicates manufacturing site

#XXXXXX

F1560

Distributed by: LESCO, Inc.

1301 East 9th Street

Cleveland, OH 44114-1849

4/5



Phone: 216.706.9250
800.321.5325
www.lesco.com
Nasdaq: LESCO

June 15, 2006

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attn: Ms. Joanne Miller, PM 23

SUBJECT: Notification – Optional Language and Fertilizer Statements
EPA #10404-88 LESCO Stonewall 0.20% Plus Fertilizer
EPA #10404-89 LESCO Stonewall 0.43% Plus Fertilizer
EPA #10404-90 LESCO Stonewall 0.68% Plus Fertilizer
EPA #10404-91 LESCO Stonewall 2.0% Plus Fertilizer

Dear Ms. Miller:

Notifications are being submitted for your review and acceptance to address the following:

- ◆ Addition to Optional/Marketing statements:
 - Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding.”
- ◆ Clarification of ‘Derived From’ statement in the Guaranteed (Fertilizer) Analysis box:
 - “Slowly Available Urea Nitrogen from Polymer Coated sulfur Coated Urea”
- ◆ Clarification of ‘Poly Plus’ composition in the Trademarks section:
 - “Poly Plus is comprised of Polymer Coated sulfur Coated Urea”

The following documentation is provided in support of the above:

- ◆ 8570-1: Notification request form for each EPA registration number
- ◆ One (1) highlighted label of each EPA registration noting the above items
- ◆ Five (5) copies of the clean label for each EPA registration

If you have any questions, or further clarification is required, please feel free to contact me at sspencer@lesco.com or 1-216-706-9250 ext 2343.

Thank you for your assistance on this matter.

Sincerely,

Susan E. Spencer
Manager
Regulatory Affairs

Enclosures

- 10404-88 LESCO Stonewall 0.20% Plus Fertilizer
- 10404-89 LESCO Stonewall 0.43% Plus Fertilizer
- 10404-90 LESCO Stonewall 0.68% Plus Fertilizer
- 10404-91 LESCO Stonewall 2.00% Plus Fertilizer

Notification to address the following:

- ◆ Addition to Optional/Marketing statements:
 - “Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding:”
- ◆ Clarification of ‘Derived From’ statement in the Guaranteed (Fertilizer) Analysis box:
 - “Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea”
- ◆ Clarification of ‘Poly Plus’ composition in the Trademark section:
 - “Poly Plus is comprised of Polymer Coated Sulfur Coated Urea”