	<u>n reverse before comple</u>	eting form.		Eorm Appro	ved. OMB No). 2070-0 0	060, Appro	yaiexcoires 2-28
€PA	Environmenta	United States			Regist	ration		intifier Number
		Applicatio	on for Pesticid	e - Sectio	nl		······································	
1. Company/Product Number 10404-93		•	2. EPA Product Menager Joanne Miller			3. Proposed Classification		
Company/Product (Nam LESCO Ronstar 0.63			РМ# 23			L	None	Restricted
5. Name and Address of A LESCO, Inc. 1301 East 9th Stree Cleveland, OH 441	t 4-1849	nde)	(b)(i), my to: EPA Re	product is s	imilar or ide	ntical in o	omposition	ection 3(c)(3) n and labeling
Check if th	is is a new address		the second se	t Name				
Amendment - Expla	in below.	, <u> </u>		inal printed lat		se to	NOTIF	ICATION
Resubmission in response to Agency letter dated				Agency letter dated "Me Too" Application. JUL 1 1 2005				
Notification - Explai	n below.		✓	Other - Explain below.				
Material This Product Wi	I Be Packaged In:		Section - III					
hild-Resistant Peckaging	Unit Packaging		Water Soluble Paci	caging	1	Containa		
Yes	Yes		Yes			Metal Plastic		
No	No		No		_ ↓ ├──	Glass		
Certification must submitted	if "Yes" Unit Packaging wgt.	No. per conteiner	lf "Yes" Package wgt	No. per container		Paper Other (Specify)	· · · · · · · · · · · · · · · · · · ·
	Information	4. Size(s) Rota	il Container	5. L	ocation of La	bel Directio	្រាទ	
1 (1)	Container							
Label		Lithogra Paper g Stencie	aph [lued d	Other				
Label () Manner in Which Label is	Affixed to Product	Paper g	Section - IV					
Label () C Manner in Which Label is	Affixed to Product	Paper g	Section - IV		cessery, to pr	ocess this	appl; sation,	;
Location of Net Contents Label [] (Manner in Which Label is Contact Point <i>(Complete</i> Une Usan Spencer	Affixed to Product	Papei g Stencia r identification	Section - IV	ontected, if nea	cessery, to pr	Telephon		le Area Code)
Lebel (Manner in Which Lebel is Contact Point <i> Complete</i> me usan Spencer I certify that the state	Affixed to Product items directly below for ments I have made on the provingily false or m	r identification T Certificati his form and a	Section - IV of individual to be co itle Manager, Regulatory on Il attachments therete	y Affairs	urate and cor	Telephone 216-706-	в No. ((лэ'ц) 9250 ехt 234 6. Date Ap Ruceive	le Area Code) C plication
Label (Manner in Which Label is Contact Point <i> Complete</i> me usan Spencer ! certify that the state ! acknowledge that an both under applicable Signature	Affixed to Product items directly below for ments I have made on ti y knowlinglly false or m aw.	Paper g Stencils r identification T N Certificati his form and al isleading state	Section - IV of individual to be co itle Manager, Regulatory on Il attachments therete	y Affairs	urate and cor	Telephone 216-706-	в No. ((лэ'ц) 9250 ехt 234 6. Date Ap Ruceive	le Area Code) C plication d
Label (Manner in Which Label is Contact Point <i> Complete</i> me Usan Spencer I certify that the state I acknowledge that an both under applicable Signature	Affixed to Product items directly below for ments I have made on ti y knowlinglly false or m aw.	r identification T Certificati his form and al isleading state 3.	itie Manager, Regulator of attachments therete ment may be punishe	y Affairs o are true, acc able by fine or	urate and cor	Telephone 216-706-	в No. ((лэ'ц) 9250 ехt 234 6. Date Ap Ruceive	le Area Code) C plication d
Label (Manner in Which Label is Contact Point <i> Complete</i> me Usan Spencer ! certify that the state ! acknowledge that an both under applicable Signature	Affixed to Product items directly below for ments I have made on ti y knowlinglly false or m aw.	Paper g Stencie r identification T Certificati his form and al isleading state 3.	Section - IV of individual to be co itle Manager, Regulatory on Il attachments thereto ment may be punishe Title	y Affairs o are true, acc able by fine or	urate and cor	Telephone 216-706-	в No. ((лэ'ц) 9250 ехt 234 6. Date Ap Ruceive	le Area Code) C plication d

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

10404-93	LESCO Ronstar 0.63% Plus Fertilizer
10404-63	LESCO Ronstar 0.95% Plus Fertilizer

Notification to address the following:

- Addition to Optional/Marketing statements:
 - "Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding:"
- Clarification of 'Derived From' statement in the Guaranteed (Fertilizer) Analysis box:
 - "Slowly Available Urea Nitrogen from Polymer Coated Sulfur Coated Urea"
- Clarification of 'Poly Plus' composition in the Trademark section:
 - o "Poly Plus is comprised of Polymer Coated Sulfur Coated Urea"

RONSTAR® 0.63% Plus Fertilizer JUL 1 1 2006

For Sale to, Use and Storage by Commercial Nursery, Turf and Landscape Personnel Only.

Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENT: Oxadiazon; 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)	
Δ1,3,4-oxadiazolin-5-one	0.63%
INERT INGREDIENTS:	
Total	

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

	FIRST AID	
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. 	
lf on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 	
lf in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
If swallowed	 Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
	HOT LINE NUMBER	
control cente	oduct container or label with you when calling a poison or doctor, or going to treatment. You may also contact 9300 for emergency treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield, or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing dust, contact with skin, eyes, or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear, and chemical resistant apron when cleaning equipment, mixing or loading. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

NON-AGRICULTURAL USES REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pasticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forests, nurseries, sod farms or greenhouses. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, locked place out of the reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows:

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA regional

office for guidance. <u>CONTAINER DISPOSAL</u>: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	xx.xx%
xx.xx% Ammoniacal Nitrogen	
xx.xx% Urea Nitrogen*	
AVAILABLE PHOSPHATE (P2O5)	xx.xx%
SOLUBLE POTASH (K2O)	xx.xx%
SULFUR (S) Total	xx.xx%
xx.xx% Free Sulfur (S)	
xx.xx% Combined Sulfur (S)	
IRON (Fe) Total	xx.xx%
xx.xx% Water Soluble Iron	
MANGANESE (Mn) Total xx.xx% Water Soluble Manganese	
xx.xx% Water Soluble Manganese	•
DERIVED FROM:	•
CHLORINE (CI) MAX.	
CHLORINE (CI) MAX. *XX.XX% Slowly available Urea Nitrogen from Polymer Coated Suk	ur Čoaled Urea

FOR LAWNS AND ORNAMENTAL TURF (NOT FOR USE ON HOME LAWNS).

(NOT FOR USE ON HOME LAW/NS) • • When applied according to directions, this product carbo psed safety on the following established turgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, TALL FESCUE, & ZOYSIAGRASS jawns and of mamental furf. It may be used for Apartment, Condominium and Corporate complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turigrass areas. May also be used on Sod Farms

CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS Annual Bluegrass (*Poa ennua*), Barnyardgrass, Crabgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed.

#XXXXXX (Front)

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader applications on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After application, if rain is not expected within 24 hours, sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control.

RATES FOR WEED CONTROL IN TURF GRASSES

FOR CRABGRASS: Apply 50 lb per 6,865 sq ft when turf has been in a yearly maintenance program (equal to 2 lb A.I. herbicide per acre). Apply 50 lb to treat 4,575 sq ft of lawn if it has been heavily infested with crabgrass (equal to 3 lb A.I. herbicide per acre). Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When to Apply" chart. <u>GOOSEGRASS_AND_PROSTRATE_SPURGE</u>: These weeds do not usually

germinate until late spring or early summer. Apply 50 lb to treat 3,430 sq ft (4 lb A.I./acre) and apply 1 to 2 weeks prior to seed germination. See chart for "When to Apply". Do not exceed maximum rate for St. Augustinegrass which is 50 lb per 4,575 sq ft.

FOR ANNUAL BLUEGRASS (Poa annua): Apply 50 lb per 3,430 to 4,575 sq ft of lawn (equal to 3 to 4 lb active herbicide per acre). Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control Annual Bluegrass present at time of application; therefore, only germinating seeds. Refer to chart for proper timing in your locale. Annual Barnyardgrass and Prostrate Spurge may also be controlled by this same rate. Do not exceed maximum rate for St. Augustinegrass, which is 3 lb active ingredient / acre.

Poa annua "ANNUAL BLUEGRASS" CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY: Apply 50 lb to treat 6,865 sq ft (2 lb A.I./acre). Apply in late summer to early fall prior to Poa annua spp. annua seed germination. Established plants will not be controlled. This product should be applied no closer than 60 days prior to overseeding. NOTE: Do not use where Poa trivialis is present in overseed mix. IMPORTANT: In many areas Annual Bluegrass exists as at least two biotypes or subspecies of Poa annua. The true annual biotype, "Poa annua spp. annua" which germinates from seed every year is controlled, while the biotype, "Poa annua spp. raptans" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning you locale.

FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE, AND CARPETGRASS: Where these weeds are a problem, apply 50 lb to treat 3,430 to 3,920 sq ft of lawn (equal to 3.5 to 4 lb active herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control emerged and growing weeds; only effective against germinating weed seeds. Observe maximum rate for St. Augustinegrass (50 lb / 4 575 sq ft).

SUGGESTED SPREADER SETTINGS Recommended Rates in lb per 1,000 sq ft

Spreader	Ground Speed	7.3 lb	10.9 lb	12.75 lb	14.60 lb
LESCO	3 mph	XX	XX*	XX*	XX*
Cyclone® or Spyker®	3 mph	XX	XX*	XX*	XX*
Lely®	41⁄2 mph	XX	XX*	XX*	XX*
Vicon®	4½ mph	XX	XX*	XX*	XX*
	•		* 2	naeeae	

Settings are approximate. The age and condition of spreader and operator's walking speed can cause variations. Be sure spreader is properly calibrated. Check spreader over 500 sq ft using 1/4 the "sq ft rate" then adjust spreader to apply exact amount.

WHEN TO APPLY

Appropriate Times to Apply This Pre-emergent Herbicide:

Oct. Nov. Dec.

За.

For CRABGRASS and other Grasses and Broadleaf Weeds germinating in the Spring in 2. For GOOSEGRASS and other Late Germinating Annual Grasses and Broadleaf Weeds such as

PROSTRATE SPURGE.

2a. GOOSEGRASS in Southern Turf. See PRECAUTIONS IN USING. 3. For Poa annua in Northern Turf. 3a. For Poa annua in Southern Turf.

4. For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipedegrass since damage will occur. Do not use on putting greens or "tees" due to varying cultural practices used. Use only the rate of 50 lb to treat 4,575 sq ft on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended in the Ornamental Plants section for this label. For newly seeded lawns, do not apply this product until after the

(S)

First letter of lot code indicates manufacturing site

second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

RATES FOR WEED CONTROL IN ORNAMENTAL PLANTS

If treatment is desired in non-turf areas, where weeds are presently growing, first remove existing weeds. Rake and smooth the soil by "breaking-up" clumps. Apply at rate as directed for specific weeds (50 lb to treat either 4,575 or 6,865 sq ft). This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

WEED CONTROL IN CONTAINER GROWN PLANTS

This product is recommended on a wide variety of Container and Field Grown Woody Ornamental Shrubs, Vines and Trees. It may be used on the following list of Container Grown Plants. Use at the rate of 50 lb per 3,430 sq ft of pot soil surface area. Plants in 14-Inch containers set out in 12-foot wide rows x 100 feet long will require 1/5 lb of this product per row of 10 pots (equivalent to approx, 4 lb A.I. per acre). Apply with a "shoulder strap held" grass-seeder or granule applicator that will direct the granules evenly to each pot soil surface for Agapanthes, Andromeda, Arborvita, Archtostaphylos, Azalea, Barberry, Boxwood, Common-Japanese, Calistemon, Camellia, Ceanothus, Cedar, Cotoneaster, Cypress, Dogwood, Douglas Fir, Eucalyptus, Euonymus-evergreen, Forsythia, Gardenia, Ginko, Heath, Heather, Hedera, Holly, Honeysuckle, Jasmine, Juniper-varieties, Justica, Leucothoe, Magnolia, Mahonia, Myrtle, Osmanthus, Periwinkle, Photina, Pine-varieties, Pittosporum, Podocarpus, Privet, Pyracantha, Quercus, Raphiolepus, Rhododendron, Rose, Star Jasmine, Syzygium, Viburnum, Xvlosma

FOR FIELD GROWN ORNAMENTALS *Apply at the rate of 50 ib per 4,575 or 6,865 sq ft of area.

This product may be used around the following Established Plants.

Arborvita, Ash, Azalea, Barberry-Japanese, Birch, Camellia, Cotoneaster, Crabapple, Dogwood, Euonymus, Forsythia, Holly, Honeysuckle, Juniper, Lilac, Maple, Oak, Periwinkle, Pine, Pittosporum, Privet, Rhododendron, Rose, Russian Olive, Spruce, Taxus, Viburnum, Xylosma. Ground Covers: Carpobrotus, Osteospermum, Sedum, Algerian Ivy, Vinca Major, English Ivy, Pachysandra, Ajuga, Euonymus.

*Use higher rate (50 lb /4.575 sq ft) for hard to kill weeds.

May be used around the following <u>Newly Transplanted Ornamentals</u>. Camellia, Dogwood, Forsythia, Hedera, Holly, Juniper, Oak, Periwinkle, Pine, Privet, Rhododendron, Russian Olive, Spruce, Taxus, Tsuga.

To obtain continued season-long weed control on the following list of plants, additional applications may be made 60 days after the previous application in the following situations.

APPLY 2 REPEAT TREATMENTS 50 Ib/ 4,575 SQ FT ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Common Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron. <u>Newly Transplanted - Container Grown Plants</u>: Common Boxwood, English Ivy, Juniper, Pine, Rhododendron.

APPLY 3 REPEAT TREATMENTS 50 lb / 4,575 SQ FT ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Holly, Rhododendron. Forest Nurseries - For Pines - Lobiolly, Slash, Eastern White.

WARRANTY

NOTICE: LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions of Use, subject to the inherent risks referred to above. LESCO, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall LESCO, Inc. be liable for consequential, special, or indirect damages resulting from the use of handling of this product.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the item number found on this bag

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-a1.com

Contains RONSTAR Herbicide, a registered trademark of Bayer. LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Poly Plus is comprised of Polymer Coated Sulfur Coated Urea. Spyker and Cyclone are registered trademarks of Spyker Spreaders LLC. Lely is a registered trademark of C Van Der Lay N.V. Vicon is a registered trademark of C Van Der Lay N.V. trademark of Ransomes Cushman-Ryan Company.

G:\REGUL\WP\STRONGSV\Labels in Progress\069173 - temp.doc Rev.11/11/05 85 Created: 10/4/02 (8/16/02)

Net Weight: 50 lb (22.7 kg)





Phone: 216.706.9250 800.321.5325 www.lesco.com Nasdag: LESCO

June 15, 2006

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Attn: Ms. Joanne Miller, PM 23

SUBJECT: Notification – Optional Language and Fertilizer Statements EPA #10404-93 LESCO Ronstar 0.63% Plus Fertilizer EPA #10404-63 LESCO Ronstar 0.95% Plus Fertilizer

Dear Ms. Miller:

Notifications are being submitted for your review and acceptance to address the following:

- Addition to Optional/Marketing statements:
 - Contains LESCO® Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding."
- Clarification of 'Derived From' statement in the Guaranteed (Fertilizer) Analysis box:
 - "Slowly Available Urea Nitrogen from Polymer Coated sulfur Coated Urea"
 - Clarification of 'Poly Plus' composition in the Trademarks section:
 - o "Poly Plus is comprised of Polymer Coated sulfur Coated Urea"

The following documentation is provided in support of the above:

- ♦ 8570-1: Notification request form for each EPA registration number
- One (1) highlighted label of each EPA registration noting the above items
- Five (5) copies of the clean label for each EPA registration

If you have any questions, or further clarification is required, please feel free to contact me at <u>sspencer@lesco.com</u> or 1-216-706-9250 ext 2343.

Thank you for your assistance on this matter.

Sincerely,

luse E Spencer

Susan E. Spencer Manager Regulatory Affairs

Enclosures