

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

SEP 14 1998

H.R. McLANE, INC.  
Agent  
LESCO, INC.  
7510 Red Road, Suite 206  
Miami, FL 33143-5321

Dear Mr. McLane:

SUBJECT: Submission of Label Amendment (Commercial/Non-commercial Label)  
Atrazine Weed & Feed 105  
EPA Reg. 10404-94  
Your Submission Dated April 29, 1998

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit a copy of the toxicity profile for the new formulation as requested by the Technical Review Branch Similarity Clinic.
2. Make the following changes listed below for the Non-Commercial Label before you release the product for shipment bearing the amended labeling:
  - a. Under the "Precautionary Statements" remove "Wash thoroughly with soap and water after handling or using".
  - b. Retain the "Personal Protective Equipment (PPE)" statement and the "User Safety Recommendations" box on the label.
  - c. Under the "Environmental Hazards" statement retain "to areas where surface water is present or to intertidal areas below the mean high water mark".

CONCURRENCES

|           |              |  |  |  |  |  |  |  |
|-----------|--------------|--|--|--|--|--|--|--|
| SYMBOL ▶  | 750EC        |  |  |  |  |  |  |  |
| SURNAME ▶ | HINDR, E.    |  |  |  |  |  |  |  |
| DATE ▶    | Sep 14, 1998 |  |  |  |  |  |  |  |

d. Retain the "Non-Agricultural Requirements" box.


e. Under the "Storage and Disposal" box retain "Do not contaminate water, food or feed by storage or disposal".

3. Submit one (1) copy of your Non-Commercial and one (1) copy of your Commercial final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



Jim Tompkins  
Product Manager (25)  
Herbicide Branch  
Registration Division (7505C)

Enclosure

Revisions required in EPA Letter dated 5/25/98 and commercial/non-commercial label split with optional marketing statements

(Commercial Label)

# Atrazine Weed & Feed 105

## For St. Augustine and Centipede Turfgrasses

A LESCO TURF FOOD WITH ATRAZINE FOR CONTROL OF BROADLEAF WEEDS AND WEEDY GRASSES

16 1/2 Pounds covers up to 5,000 Sq.Ft. of Turfgrass  
Prevents Crabgrass, Paspalums, and Spotted Spurge.  
Controls Florida Pusley, Lippia, Pigweed.  
Over Forty Five More Weeds as Listed.

A WEED & FEED FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS LAWNS

### ACTIVE INGREDIENTS:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine..... 1.01%

Related Compound..... 0.04%

INERT INGREDIENTS:..... 98.95%

Total ..... 100.00%

Contains 21.0 lbs of Atrazine per ton.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

SEP 14 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

10464-94

KEEP OUT OF REACH OF CHILDREN

CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid inhalation or contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling or using. Remove clothing and wash clothing before reuse. Keep away from feed or food products.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. IF ON SKIN: Wash with plenty of soap and water. Get medical attention. IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks and a Dust/Mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. If product is spilled; while wearing a dust mask, sweep up product and use as this label instructs (as a Weed & Feed) or dispose of in accordance with the STORAGE AND DISPOSAL section on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes and socks.

**WEEDS CONTROLLED OR SUPPRESSED**

- |  |   |  |
|--|---|--|
| Annual Bluegrass<br>( <i>Poa Annua</i> )   | Cutleaf Evening Primrose                      | Oxalis ( <i>Common and Yellow Woodsorrel</i> ) |
| Annual Sowthistle                          | Dichondra                                     | Pigweed  |
| Clover                                     | Dollar Weed                                   | Purslane                                       |
| Black Medic                                | Florida Beggarweed                            | Ragweed  |
| Blue Toadflax                              | Florida Betony<br>( <i>Rattlesnake Weed</i> ) | Red Sorrel                                     |
| Bull Paspalum                              | Florida Pusley                                | Rescuegrass                                    |
| Buckhorn Plantain                          | Fragrant Cudweed<br>( <i>Rabbit Tobacco</i> ) | Sandspur                                       |
| Bur Clover                                 | Henbit  | Sedge*   |
| Carolina Geranium<br>( <i>Cranesbill</i> ) | Knotweed                                      | Shepards Purse                                 |
| Chickweed ( <i>Common and Mouse Ear</i> )  | Lambsquarter                                  | Sicklepod                                      |
| Cocklebur                                  | Lespedeza                                     | Spotted Spurge                                 |
| Common Dandelion                           | Lippia ( <i>Matchweed</i> )                   | Spurweed                                       |
| Common Purslane                            | Moneywort                                     | Stinging Nettle                                |
| Crabgrass                                  | Morning Glory                                 | Swinecrest                                     |
| Crowfoot Grass                             | Narrowleaf Vetch                              | Venus Looking Glass                            |
| Cudweed                                    | Parsley-Piert                                 | Virginia Creeper                               |
|  |   | Wild Mustard                                   |
|  |   | Wood Sorrel                                    |

**NON-AGRICULTURAL REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

**IMPORTANT:** In addition to providing turfgrass with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from seeds".

**FOR WEEDS THAT ARE PRESENT:** Includes broadleaf weeds from the list such as; Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.

**FOR WEEDS GERMINATING FROM SEEDS:** Applied before weed seeds germinate; includes grassy weeds from the list such as; Crabgrass, Paspalum, \*Sedge, (an additional application may be required to control sedge) and Spotted Spurge. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

**WHERE TO USE**

This material can be used on St. Augustinegrass and Centipedegrass grown in home and corporate lawns, parks and recreational areas, golf courses, cemeteries, athletic fields, and sod farms. Do not plant any crop (flowers or vegetables) to treated areas for 18 months since plant injury may result. **DO NOT USE ON ANY VARIETIES OF BLUEGRASS, FESCUE, BENTGRASS, BAHIAGRASS, RYEGRASS, BERMUDAGRASS, OR DICHONDRA LAWNS.**

**WHEN TO APPLY**

Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good weed control.

For pre-emergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

**HOW MUCH TO APPLY**

This product may be applied with a broadcast rotary type spreader. Apply 3.27 lbs to 4.37 lbs uniformly over 1,000 sq.ft. of lawn area. (equivalent to the required 1.5 lb. to 2.0 lbs Atrazine active ingredient per acre). This is also equal to 16 1/2 pounds of this product to treat 3,775 sq.ft. up to 5,000 sq.ft. Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommended rates. Apply to dry grass. For best results, rainfall or watering is necessary within 7 to 10 days after treatment to activate the herbicide and fertilizer by washing the product off the grass blades down to the soil surface.

**APPROXIMATE ROTARY SPREADER SETTINGS**

| Pounds of Product/1,000 sq.ft. | LESCO | Cyclone | Ortho | True Temp. 5000 |
|--------------------------------|-------|---------|-------|-----------------|
| To apply 3.27#/ 1,000 sq.ft.   | #8    | 2 3/4   | 4 1/2 | 2 3/4           |
| To apply 4.37#/ 1,000 sq.ft.   | #9    | 3       | 5     | 3               |

Settings are approximate. The age and condition of spreader, and operator's walking speed can cause variations. Be sure spreader is properly calibrated. Check spreader over 500 sq.ft. using 1/2 the "1,000 sq.ft. rate"; adjust spreader to apply exact amount.

**USE PRECAUTIONS**

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

**GUARANTEED FERTILIZER ANALYSIS**

- Total Nitrogen (N) ..... XX %
- Ammoniacal Nitrogen -XX %
- Water Soluble Nitrogen -XX %
- Coated Slow Release Nitrogen -XX %
- Water Insoluble Nitrogen -XX %
- Available Phosphate (P<sub>2</sub>O<sub>5</sub>)XX %
- Soluble Potash (K<sub>2</sub>O)... XX %
- Chlorine not more than XX %

Derived from Urea, Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to \_\_\_ lbs of Calcium Carbonate per ton.

**Statement of Secondary Plant Foods**

Total Magnesium as Mg - \_\_\_%; Iron as Fe - \_\_\_%; Water Soluble Magnesium as Mg - \_\_\_%;  
 Zinc as Zn - \_\_\_% Manganese as Mn - \_\_\_%; Boron as B - \_\_\_% Copper as Cu - \_\_\_%;  
 Molybdenum as Mo - \_\_\_% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide,  
 Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

**NOTICE:** LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. LESCO, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall LESCO, Inc. be liable for consequential, special, or indirect damages resulting from the use or handling of this product. (ATRA-105.p65\02-05-98)

EPA REG. NO. 10404-94

EPA EST. NO. 10404-OH-1, 10404-FL-1, 35512-FL-1, 49991-TX-1

NET WEIGHT: 16 1/2 ; 25; 33; 50 POUNDS (7.48; 11.34; 14.97; 22.68 kg's)

LESCO, Inc.

20005 Lake Road

Rocky River, OH 44116-1545 U.S.A.

(Non-Commercial Label)

# Atrazine Weed & Feed 105

## For St. Augustine and Centipede Turfgrasses

A LESCO TURF FOOD WITH ATRAZINE FOR CONTROL OF BROADLEAF WEEDS AND WEEDY GRASSES

16 1/2 Pounds covers up to 5,000 Sq.Ft. of Turfgrass  
Prevents Crabgrass, Paspalums, and Spotted Spurge.  
Controls Florida Pusley, Lippia, Pigweed.  
Over Forty Five More Weeds as Listed.

A WEED & FEED FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS LAWNS

### ACTIVE INGREDIENTS:

|  |         |
|--|---------|
| Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine..... | 1.01%   |
| Related Compound.....  | 0.04%   |
| INERT INGREDIENTS:.....  | 98.95%  |
| Total .....  | 100.00% |

Contains 21.0 lbs of Atrazine per ton.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

SEP 14 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended for the pesticide  
registered under EPA Reg. No.

10404-94

KEEP OUT OF REACH OF CHILDREN

### CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid inhalation or contact with eyes, skin or clothing. Wear long-sleeved shirt and long pants, socks, and shoes and waterproof gloves. Wash thoroughly with soap and water after handling or using. Remove clothing and wash clothing before reuse. Keep away from feed or food products.

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

~~Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks and a Dust/Mist filtering respirator (MSHA/NIOSH) approval number prefix TC 21C. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.~~

#### User Safety Recommendations

~~Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.~~

**ENVIRONMENTAL HAZARDS**

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, ~~to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. If product is spilled; while wearing a dust mask, sweep up product and use as this label instructs (as a Weed & Feed) or dispose of in accordance with the STORAGE AND DISPOSAL section on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. ~~Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.~~

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**AGRICULTURAL USE REQUIREMENTS**

~~Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.~~

~~Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes and socks.~~

**WEEDS CONTROLLED OR SUPPRESSED**

- |  |   |  |
|--|---|--|
| Annual Bluegrass<br>( <i>Poa Annua</i> )   | Cutleaf Evening Primrose                      | Oxalis ( <i>Common and Yellow Woodsorrel</i> ) |
| Annual Sowthistle                          | Dichondra                                     | Pigweed  |
| Clover                                     | Dollar Weed                                   | Purslane                                       |
| Black Medic                                | Florida Beggarweed                            | Ragweed  |
| Blue Toadflax                              | Florida Betony<br>( <i>Rattlesnake Weed</i> ) | Red Sorrel                                     |
| Bull Paspalum                              | Florida Pusley                                | Rescuegrass                                    |
| Buckhorn Plantain                          | Fragrant Cudweed                              | Sandspur                                       |
| Bur Clover                                 | ( <i>Rabbit Tobacco</i> )                     | Sedge*   |
| Carolina Geranium<br>( <i>Cranesbill</i> ) | Henbit  | Shepards Purse                                 |
| Chickweed ( <i>Common and Mouse Ear</i> )  | Knotweed                                      | Sicklepod                                      |
| Cocklebur                                  | Lambsquarter                                  | Spotted Spurge                                 |
| Common Dandelion                           | Lespedeza                                     | Spurweed                                       |
| Common Purslane                            | Lippia ( <i>Matchweed</i> )                   | Stinging Nettle                                |
| Crabgrass                                  | Moneywort                                     | Swinecrest                                     |
|  | Morning Glory                                 | Venus Looking Glass                            |
|  | Narrowleaf Vetch                              | Virginia Creeper                               |



Crowfoot Grass  
Cudweed

Parsley-Piert

Wild Mustard  
Wood Sorrel

**NON-AGRICULTURAL REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

**IMPORTANT:** In addition to providing turfgrass with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from seeds".

**FOR WEEDS THAT ARE PRESENT:** Includes broadleaf weeds from the list such as; Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.

**FOR WEEDS GERMINATING FROM SEEDS:** Applied before weed seeds germinate; includes grassy weeds from the list such as; Crabgrass, Paspalum, \*Sedge, (an additional application may be required to control sedge) and Spotted Spurge. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

**WHERE TO USE**

This material can be used on St. Augustinegrass and Centipedegrass grown in home and corporate lawns, parks and recreational areas, ~~golf courses, cemeteries, athletic fields, and sod farms.~~ Do not plant any crop (flowers or vegetables) to treated areas for 18 months since plant injury may result. DO NOT USE ON ANY VARIETIES OF BLUEGRASS, FESCUE, BENTGRASS, BAHIAGRASS, RYEGRASS, BERMUDAGRASS, OR DICHONDRA LAWNS.

**WHEN TO APPLY**

Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good weed control.

For pre-emergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

**HOW MUCH TO APPLY**

This product may be applied with a broadcast rotary type spreader. Apply 3.27 lbs to 4.37 lbs uniformly over 1,000 sq.ft. of lawn area. (equivalent to the required 1.5 lb. to 2.0 lbs Atrazine active ingredient per acre). This is also equal to 16 1/2 pounds of this product to treat 3,775 sq.ft. up to 5,000 sq.ft. Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommended rates. Apply to dry grass. For best results, rainfall or watering is necessary within 7 to 10 days after treatment to activate the herbicide and fertilizer by washing the product off the grass blades down to the soil surface.

**APPROXIMATE ROTARY SPREADER SETTINGS**

| Pounds of Product/1,000 sq.ft. | LESCO | Cyclone | Ortho | True Temp. 5000 |
|--------------------------------|-------|---------|-------|-----------------|
| To apply 3.27#/ 1,000 sq.ft.   | #8    | 2 3/4   | 4 1/2 | 2 3/4           |
| To apply 4.37#/ 1,000 sq.ft.   | #9    | 3       | 5     | 3               |

10/12

Atrazine Weed & Feed 105 (Non-Commercial Label)

Page 8

Settings are approximate. The age and condition of spreader, and operator's walking speed can cause variations. Be sure spreader is properly calibrated. Check spreader over 500 sq.ft. using 1/2 the "1,000 sq.ft. rate"; adjust spreader to apply exact amount.

USE PRECAUTIONS

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

- Total Nitrogen (N) ..... XX %
- Ammoniacal Nitrogen -XX %
- Water Soluble Nitrogen -XX %
- Coated Slow Release Nitrogen -XX %
- Water Insoluble Nitrogen -XX %
- Available Phosphate (P<sub>2</sub>O<sub>5</sub>)XX %
- Soluble Potash (K<sub>2</sub>O)... XX %
- Chlorine not more than XX %

Derived from Urea, Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to \_\_\_ lbs of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

- Total Magnesium as Mg - \_\_\_%; Iron as Fe - \_\_\_%; Water Soluble Magnesium Mg - \_\_\_%;
- Zinc as Zn - \_\_\_% Manganese as Mn - \_\_\_%; Boron as B - \_\_\_% Copper as Cu - \_\_\_%;
- Molybdenum as Mo - \_\_\_% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax. Sodium Molybdate.

STORAGE AND DISPOSAL

~~Do not contaminate water, food or feed by storage or disposal.~~

**STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

~~Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.~~

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**HOUSEHOLD:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

NOTICE: LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. LESCO, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall LESCO, Inc. be liable for consequential, special, or indirect damages resulting from the use or handling of this product. (ATRA-105-p65\02-05-98)

EPA REG. NO. 10404-94

EPA EST. NO. 10404-OH-1, 10404-FL-1, 35512-FL-1, 49991-TX-1

NET WEIGHT: 16 1/2 ; 25; 33; 50 POUNDS (7.48; 11.34; 14.97; 22.68 kg's)

LESCO, Inc.

20005 Lake Road

Rocky River, OH 44116-1545 U.S.A.

11/12

Optional Marketing Statements

- Kills dollarweed and other common lawn weeds
- Contains TruRelease Nitrogen for Extended Greening
- Use on St. Augustine and Centipede turf only.

Call 1-300-XXX-XXXX for more information.

**SATISFACTION GUARANTEED PANEL**

(GUARANTEE SEAL)

**SATISFACTION GUARANTEED OR YOUR MONEY BACK.** TruGreen is committed to providing the highest quality products. If you are not totally satisfied with our products, we will replace your product or refund your money. Just call us at 1-800-TRU-GREEN and we will be happy to help you.

**TRUGREEN ANNUAL TURF FERTILIZER PROGRAM**

**Step 1 Apply Late Winter-Early Spring**

TruGreen by ChemLawn Crabgrass Control plus Fertilizer Prevents crabgrass, foxtail, poa annua, spurge and other annual grasses while feeding your lawn with long-lasting, deep greening controlled-release nitrogen,

**Step 2 Apply Mid-Late Spring**

TruGreen by ChemLawn Weed & Feed\* Kills over 100 varieties of weeds, roots and all! Feeds with controlled-release nitrogen for longer lasting greening

**Step 3 Apply Late Spring-Summer**

TruGreen by ChemLawn Lawn Insect Control with Fertilizer Controls insects while greening your lawn with controlled-release nitrogen for dependable, predictable results

**Step 4 Apply Late Summer-Early Fall**

TruGreen by ChemLawn Fertilizer Provides extended greening throughout the summer months with controlled-released nitrogen for dependable, predictable results!

\* TruGreen by ChemLawn St. Augustine Weed & Feed can be used as step 2 in your Annual Program ONLY on St. Augustine, and Centipede Lawns

**Start Using the TruGreen by ChemLawn Annual Turf Fertilizer Program**

(Chart showing the time of year to use what products by agronomic zone)

**Northern Zone**

- Step 1 Late March - Early May
- Step 2 April 1 - Early June
- Step 3 Mid May-End of August
- Step 4 August 1- End of September

**Transitional Zone**

- Step 1 Mid March- Mid April
- Step 2 Late March - End of May
- Step 3 Mid May-Early September
- Step 4 First of September - Late October

**Southern Zone**

- Step 1 Mid January- Mid April
- Step 2 Mid March-Mid May
- Step 3 Late April-End of September
- Step 4 Mid August - Mid November

(Graphic of the map showing where the various zones are located)

---

**SPREADER CALIBRATION & PATTERNS**

Rotary spreaders, like the kind our technicians use, are the most preferred. These spreaders are easier to use and reduce the "streaking" problems associated with drop spreaders. Drop spreaders work well, but require more patience to avoid gaps and streaks. Streaks appear as irregular light green areas in the lawn a week or so after application. This comes from uneven fertilizer coverage. If using a drop spreader, be extra careful to avoid gaps in coverage.

Always apply with a properly calibrated rotary or drop spreader. Never apply by hand.  
(photo of technician and spreader)

| <u>Rotary Spreader</u>     | <u>Settings</u> | <u>Drop Spreader</u>   | <u>Settings</u> |
|----------------------------|-----------------|------------------------|-----------------|
| Scotts SpeedyGreen 1000    | 4.5             | Scotts AccuGreen 1000  | 7.75            |
| Scotts SpeedyGreen 2000    | 5.75            | Scotts AccuGreen 3000  | 6.5             |
| Scotts SpeedyGreen 3000    | 5.25            | Craftsman/Republic 530 | 7.75            |
| Scotts EasyGreen           | 27.75           |                        |                 |
| Craftsman/Republic 540     | 2.5             |                        |                 |
| Vigoro SB4000V             | 11              |                        |                 |
| Precision/ Red Devil SB550 | 6.75            |                        |                 |

To avoid gaps or streaks, first spread boarder of lawn. Fill in center with uniform, parallel passes.

For best results, apply with a fertilizer spreader at the settings listed . [delivers 4.2#/1000 sqft]

---

Available Exclusively Through  
Tru-Green-ChemLawn  
Memphis, TN 38120-9792

---

5/25/98 EPA Stamped Label

8/24/98 Revisions required in EPA Letter dated 5/25/98 and commercial/non-commercial label split