# 10404-94 Atrazine 1.05% Plus rertilizer

Weed Control with Premium Lawn Food

50 Pounds covers up to 15,240 Sq Ft of Turfgrass Prevents Crabgrass, Paspalums, and Spotted Spurge Controls Dollarweed, Florida Pusley, Lippia, Pigweed Over Forty-Fve More Weeds as Listed

& FEED FOR ST. AUGUSTINEGRASS AND **CENTIPEDEGRASS LAWNS** 

Contains LESCO Poly Plus® Sulfur Coated Urea to provide uniform growth with extended nitrogen feeding

### **ACTIVE INGREDIENTS:**

Atrazine; 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	1.059
Related active triazine compounds	0.049
OTHER INGREDIENTS:	98.919
Total	000 000

# KEEP OUT OF REACH OF CHILDREN CALITION

FIRST AID			
Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by the poison control cente or doctor On ort give anything by mouth to an unconscious person.			
Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.     Call a poison control center or doctor for treatment advice.			
Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.			
Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.     Call a poison control center or doctor for further treatment advice.			
ict container or label with you when calling a poison control center or			
for treatment. You may also contact Chem-Trec at 1-800-424-9300			
medical treatment information. PRECAUTIONARY STATEMENTS			

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Loaders, applicators and other handlers must wear long-sleeved shirt, long pants, shoes and socks. Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handheld rotary applicator. Such equipment may only be used for

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators and other handlers must wear long-sieeved shirt and long pants, shoes and socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

# **User Safety Recommendations**

User Sarety Recommendations
Users should wash hands before eating, dinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

# ENVIRONMENTAL HAZAROS

Atrazine can travel (seep or leach) through soil and can enter ground wa Arrazine can travel (seep or teach) through soil and can enter ground water which may be used as drinking water. Attrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Product must not be mixed or loaded within 50 feet of intermittent streams and the recognition of the applied.

rivers, natural or impounded lakes and reservoirs. Product must not be applied within 68 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 68 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed or loaded or used within 50 feet of all wells, including Additional State imposed requirements regarding well-head setbacks and

operational area containment must be observed

One of the following restrictions must be used in applying atrazine to tile-outletted fields containing standpipes.

- Do not apply within 86 feet of standpipes in tile-outletted fields. Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- product to the entire tile-outletted field under a no-till practice only where a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash

# **#XXXXXX**

DERIVED FROM: CHLORINE (CI) Max..



GUARANTEED ANALYSIS	ı
TOTAL NITROGEN (N)	XX.XX%
XX.XX% Ammoniacal Nitrogen	
XX.XX% Urea Nitrogen*	
XX.XX% Water Insoluble Nitrogen	
AVAILABLE PHOSPHATE (P2O6)	
SOLUBLE POTASH (K <sub>1</sub> O)	
MAGNESIUM (Mg) Total	XX XX%
XX.XX% Water Soluble Magnesium (Mg)	
SULFUR (S) Total	XX.XX%
XX.XX% Free Sulfur (S)	
XX.XX% Combined Sulfur (S)	
COPPER (Cu) Total	XX.XX%
XX.XX% Water Solubie Copper (Cu)	
	XX.XX%
XX.XX% Water Soluble Iron (Fe)	
MANGANESE (Mn) Total	XX.XX%
XX.XX% Water Soluble Manganese (Mn)	
ZINC (Zn) Total	XX.XX%
XX.XX% Water Soluble Zinc (Zn)	

\*X.XX% Slowly available Nitrogen from LESCO Poly Plus® Sulfur Coated This product has different N-P-K fertilizer ratios to meet the needs for different geographies, varieties, and temperatures.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

When applying multiple products containing atrazine, the total pounds of atrazine

applied in pounds of atrazine per acre may not exceed 4 pounds of atrazine from

Turfgrass at Residential Sites (including homes, daycare facilities, schools, playgrounds, parks, recreational areas, and sports fields.)
This product must be watered in immediately after application. Watering-in must be

rmed by the commercial applicator or the commercial applicator must provide

med by the commercial applicator or the commercial applicator must provide slidwing instructions to the resident or owner in writing:

-This product must be watered in immediately.

- Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A ANY DSE OF THIS PRODUCT IN AN AREA WHERE USE IS PROFIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.strazine-watershed.info or 1-886-385-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact. LESCO for a refund

# WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poe annual) Annual Sowthistie Clover Black Medic Rive Toedfiles **Buckhorn Plantein** Bur Clover Carolina Geranium (Crenesbill) eed (Con and Mouse Ear) Cocklebur Common Dandelion Common Puntiene Crc wfoot G. ass

Cutteef Evening Primrose Dichondre Dollarweed Floride Begge Florida Batchy (Rattiosnake We Florida Pusiey Fragrant Cudweed (Rebbit Tobacco) Henioit

Lippin (Matchy Viorning Glory

Oxalis (Common and Yellow Woodsorrel) Pigweed Purslane Sandsour Shapharda Pursa Sickiapodi Spotted Source Stinging Nettic Venua Looking Glass Virginia Creep Wild Mustard

**ACCEPTED** Under the Federal Insecticide, Fungicide, and Rodonticide Act as amended, for the pesticide registered under EPA Reg. No. 0404-94

# Atrazine 1.05% Plus Fertilizer

# NON-AGRICULTURAL REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nursenes, or greenhouses <u>Do not enter</u> or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

IMPORTANT: In addition to providing turfgrass with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from

FOR WEEDS THAT ARE PRESENT: Includes broadleaf weeds from the list such as; Chickwed, Creeping Charle, Dandellon, Evening Primrose, Florida Betony, Florida Pustry, Lippia, Oxalls, Pigweed, Plantain, Rabbit Tobacco, and Toadflax. FQR\_WEEDS\_GEMINATING\_FROM\_SEEDS; Applied before weed seeds germinate, includes grassy weeds from the list such as, Crabgrass, Paspalum, and Spotted Sourge.

### WHERE TO USE

FOR CONTROL of weeds in St. Augustine and Centipede grass in or on residential lawns, corporate lawns, parks, recreational areas, golf course, cerneteries, and

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes

NOT FOR USE on Bluegrass, Fescue, Bentgrass, Bahiagrass, Ryegrass, Bermudagrass, or Dichondra lawns.

### WHEN TO APPLY

WHEN TO APPLY

WHEN TO APPLY

Best waed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post-emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadlead weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good week

control.

For pre-amergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Flonda Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

## **USE PRECAUTIONS**

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of omamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

DO NOT APPLY THIS PRODUCT IN A WAY THAT WILL CONTACT ANY PERSON OR PET, EITHER DIRECTLY OR THROUGH DRIFT. KEEP PEOPLE AND PETS OUT OF THE AREA DURING APPLICATION. THIS PRODUCT MUST BE WATERED IN IMMEDIATELY AFTER APPLICATION. DO NOT ENTER OR ALLOW OTHERS (INCLUDING CHILDREN OR PETS) TO ENTER THE TREATED AREAS (EXCEPT THOSE INVOLVED IN THE WATERING) UNTIL THE WATERING-IN IS COMPLETE AND THE SURFACE IS DRY

Apply 3.27 to to 4.37 to uniformly over 1.000 sq ft of lawn area, (equivalent to the required 1.5 – 2.0 to Atrazine per acre). Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommendate rates. Do not apply more than 4 to a 1 Atrazine (2 applications at 4.37 to product) per acre per year. Apply to dry grass. This product must be watered in immediately

### SUGGESTED SPREADER SETTINGS\*

	(Kecommecoed Kates	in no per 1,υυυ sq πj	
Spreader		3.27 lb	4.37 lb
LESCO	Calibration Gauge	#XX	#XX
SCOTTS* R	B-A	XX	XX
Cyclone® or a	Spyker <sup>®</sup>	XX	XX
Lelv	• •	XX II	XX II

\*Settings are approximate. Age, condition of spreader and operator speed can cause variations. Be sure to calibrate your epreader with each application.

# STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

<u>PESTICIDE STORAGE</u>: Store product in original labeled container in a cool, dry,

locked place out of reach of children.

locked place out of reach of children.

PESTICIPE\_DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Dispose of emptied bag(s) in a sanitary landfill approved for pesticide disposal, or by incineration.

SPILL. In case of spill, sweep up material and dispose of material according to product disposal directions listed above.

# WARRANTY

WARRANTY
LESCO, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use LESCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of LESCO and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or reliabling to flow label directions will be assumed by the Buyer or User IN NO CASE WILL LESCO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE, OR USE OF THIS PRODUCT.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9<sup>th</sup> Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the Item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.rsquistory-info-e1.com">http://www.rsquistory-info-e1.com</a>

LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. SCOTTS is a registered trademark of The SCOTT Company, Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. Lely is a registered trademark of C Van Der Leiy N.V.

G:\REGUL\WP\Regul Private\MatrLb\\\_ERIEV\\ ADDRESS\New EPA Approved \Atrazine 105FL formatted.doc (0705) Rev 3/26/07

Net Weight: XX lb (XX kg)

EPA REG. NO. 10404-94 EPA EST. NO. 82757-OH-001 (M), 82757-FL-001 (S)



F1560

Distributed by: LESCO, Inc. 1301 East 9th Street Cleveland, OH 44114-1849