

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

- Registration
- Amendment
- Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 10404-95	2. EPA Product Manager Jim Thompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Atrazine Weed & Feed 45	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) LESCO, Inc. 15885 Sprague Rd. Strongsville, Ohio 44136-1799 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

- Amendment - Explain below.
- Resubmission in response to Agency letter dated \_\_\_\_\_
- Notification - Explain below.
- Final printed labels in response to Agency letter dated \_\_\_\_\_
- "Me Too" Application.
- Other - Explain below.

NOTIFICATION

SEP 11 2001

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Combination of Redundant Labeling Statements per PR Notice 98-10.  
This notification is consistent with the provision of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted			If "Yes" Unit Packaging wgt.	No. per container	<input type="checkbox"/> Glass
			If "Yes" Package wgt	No. per container	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Bernadine Javorek	Title Manager, Regulatory Affairs	Telephone No. (Include Area Code) 440-783-9250
2. Signature <i>Bernadine Javorek</i>		6. Date Application Received (Stamp(s))
3. Title Manager, Regulatory Affairs		
4. Typed Name Bernadine Javorek	5. Date 8/29/01	



# Atrazine Weed & Feed 45

For St. Augustine and Centipede Turfgrasses

A LESCO TURF FOOD WITH ATRAZINE FOR CONTROL OF BROADLEAF WEEDS AND WEEDY GRASSES

19 1/2 Pounds covers up to 2,500 Sq.Ft. of Turfgrass  
Prevents Crabgrass, Paspalums, and Spotted Spurge.  
Controls Florida Pusley, Lippia, Pigweed.  
Over Forty Five More Weeds as Listed.

### A WEED & FEED FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS LAWNS

#### ACTIVE INGREDIENTS:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine..... 0.43%  
Related Compound..... 0.02%

INERT INGREDIENTS:..... 99.55%

Total ..... 100.00%

Contains 9 lbs. of Atrazine per ton.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if symptoms or irritation persist. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists. **IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMAN AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid inhalation or contact with eyes, skin or clothing. Wear long-sleeved shirt and long pants, socks and shoes and waterproof gloves. Wash thoroughly with soap and water after handling. Remove clothing and wash clothing before reuse.

##### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; waterproof gloves; and shoes plus socks and a Dust/Mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

##### User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. If product is spilled, while wearing a dust mask, sweep up product and use as this label instructs (as a Weed & Feed) or dispose of in accordance with the STORAGE AND DISPOSAL section on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes and socks.

**WEEDS CONTROLLED OR SUPPRESSED**

- |                                            |                                               |                                                |
|--------------------------------------------|-----------------------------------------------|------------------------------------------------|
| Annual Bluegrass<br>( <i>Poa Annua</i> )   | Cutleaf Evening Primrose                      | Oxalis ( <i>Common and Yellow Woodsorrel</i> ) |
| Annual Sowthistle                          | Dichondra                                     | Pigweed                                        |
| Clover                                     | Dollar Weed                                   | Purslane                                       |
| Black Medic                                | Florida Beggarweed                            | Ragweed                                        |
| Blue Toadflax                              | Florida Betony<br>( <i>Rattlesnake Weed</i> ) | Red Sorrel                                     |
| Bull Paspalum                              | Florida Pusley                                | Rescuegrass                                    |
| Buckhorn Plantain                          | Fragrant Cudweed                              | Sandspur                                       |
| Bur Clover                                 | ( <i>Rabbit Tobacco</i> )                     | Sedge*                                         |
| Carolina Geranium<br>( <i>Cranesbill</i> ) | Henbit                                        | Shepards Purse                                 |
| Chickweed ( <i>Common and Mouse Ear</i> )  | Knotweed                                      | Sicklepod                                      |
| Cocklebur                                  | Lambsquarter                                  | Spotted Spurge                                 |
| Common Dandelion                           | Lespedeza                                     | Spurweed                                       |
| Common Purslane                            | Lippia ( <i>Matchweed</i> )                   | Stinging Nettle                                |
| Crabgrass                                  | Moneywort                                     | Swinecrest                                     |
| Crowfoot Grass                             | Morning Glory                                 | Venus Looking Glass                            |
| Cudweed                                    | Narrowleaf Vetch                              | Virginia Creeper                               |
|                                            | Parsley-Piert                                 | Wild Mustard                                   |
|                                            |                                               | Wood Sorrel                                    |

**NON-AGRICULTURAL REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

**IMPORTANT:** In addition to providing turfgrass with required plant nutrients this product will control both "weeds that are present" and "weeds germinating from seeds".

**FOR WEEDS THAT ARE PRESENT:** Includes broadleaf weeds from the list such as; Chickweed, Creeping Charlie, Dandelion, Evening Primrose, Florida Betony, Florida Pusley, Lippia, Oxalis, Pigweed, Plantain, Rabbit Tobacco, and Toadflax.

**FOR WEEDS GERMINATING FROM SEEDS:** Applied before weed seeds germinate; includes grassy weeds from the list such as; Crabgrass, Paspalum, \*Sedge, (an additional application may be required to control sedge) and Spotted Spurge. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

**WHERE TO USE**

This material can be used on St. Augustinegrass and Centipedegrass grown in home and corporate lawns, parks and recreational areas, golf courses, cemeteries, athletic fields, and sod farms. Do not plant any crop (flowers or vegetables) to treated areas for 18 months since plant injury may result.

**DO NOT USE ON ANY VARIETIES OF BLUEGRASS, FESCUE, BENTGRASS, BAHIAGRASS, RYEGRASS, BERMUDAGRASS, OR DICHONDRA LAWNS.**

**WHEN TO APPLY**

Best weed control will result if applied before weeds sprout since the product acts as a "pre-emergent" and also as a "post emergent" herbicide. Established weeds are harder to kill and may require an additional application in the fall or spring. For newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. For broadleaf weeds apply anytime the weeds are present and actively growing. Early spring application is recommended for best control such as January 1st up to March 1st, otherwise it may be applied up to June 15th with good weed control.

For pre-emergence weed control (see above) apply before seeds germinate. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeat application may be necessary. Wait 2 months for a repeat application. Apply when air is calm to prevent particle wind drift. Apply this product only when grass blades are dry. To verify proper application timing consult your State Agricultural Extension Service, Turfgrass Specialist.

**HOW MUCH TO APPLY**

This product may be applied with a broadcast rotary type spreader. Apply 7.65 lbs. to 10 lbs. uniformly over 1,000 sq.ft. of lawn area. (equivalent to the required 1.5 lb. to 2.0 lbs. Atrazine active ingredient per acre). This is also equal to 19 1/2 pounds of this product to treat 1,950 sq.ft. up to 2,500 sq.ft. Use the lower rate for average weed problems; the higher rate for heavy weed population. Do not exceed recommended rates. Apply to dry grass. For best results, rainfall or watering is necessary within 7 to 10 days after treatment to activate the herbicide and fertilizer by washing the product off the grass blades down to the soil surface.

**APPROXIMATE ROTARY SPREADER SETTINGS**

Pounds of Product/1,000 sq.ft.	LESCO	Cyclone	Ortho	True Temp. 5000
To apply 7.65#/ 1,000 sq.ft.	#12	3 3/4	6	7
To apply 10.0#/ 1,000 sq.ft.	#13	4	7	8

Settings are approximate. The age and condition of spreader, and operator's walking speed can cause variations. Be sure spreader is properly calibrated. Check spreader over 500 sq.ft. using 1/2 the "1,000 sq.ft.rate"; adjust spreader to apply exact amount.

575

### USE PRECAUTIONS

Sweep product off walks and painted surfaces following application to avoid possible discolorations. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses, since injury to those other grasses may result.

### GUARANTEED FERTILIZER ANALYSIS

- Total Nitrogen (N) ..... XX %
- Ammoniacal Nitrogen -XX %
- Water Soluble Nitrogen -XX %
- Coated Slow Release Nitrogen -XX %
- Water Insoluble Nitrogen -XX %
- Available Phosphate (P<sub>2</sub>O<sub>5</sub>)XX %
- Soluble Potash (K<sub>2</sub>O)... XX %
- Chlorine not more than XX %

Derived from Urea, Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to \_\_\_lbs. of Calcium Carbonate per ton.

#### Statement of Secondary Plant Foods

Total Magnesium as Mg - \_\_\_\_%; Iron as Fe - \_\_\_\_%; Water Soluble Magnesium Mg - \_\_\_\_%;  
 Zinc as Zn - \_\_\_\_% Manganese as Mn - \_\_\_\_%; Boron as B - \_\_\_\_% Copper as Cu - \_\_\_\_%;  
 Molybdenum as Mo - \_\_\_\_% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide,  
 Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

NOTICE: LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. LESCO, Inc. makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall LESCO, Inc. be liable for consequential, special, or indirect damages resulting from the use or handling of this product. (ATRA-45.p65\02-03-98)

EPA REG. NO. 10404-95

EPA EST. NO. 10404-OH-1

EPA EST. NO. 10404-FL-1

EPA EST. NO. 35512-FL-1

EPA EST. NO. 49991-TX-1

NET WEIGHT: 19 1/2 ; 25; 39; 50 POUNDS (8.84; 11.34; 17.69; 22.68 kg's)



20005 Lake Road - Rocky River, OH 44116 U.S.A.

8/29/01 EPA Stamped Label