

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 10404-98	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) LESCO Pre-M 1.5% Fertilizer Broadleaf Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) LESCO, Inc. 15885 Sprague Road Strongsville, Ohio 44136-1799 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

JUN 24 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Name Per PR Notice(s) 95-2: LESCO Pre-M 1.5% Plus Fertilizer. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged in:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Gloria M. Sieloff	Title Director, Regulatory Affairs	Telephone No. (Include Area Code) 440-783-9250 ext. 2597
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		8. Date Application Received (Stamped)
2. Signature <i>Gloria Sieloff</i>	3. Title Director, Regulatory Affairs	
4. Typed Name Gloria M. Sieloff	5. Date 6/11/03	

Manganese (Mn) TotalXX.XX%
 XX.XX% Water Soluble Manganese (Mn)
 DERIVED FROM:
 Chlorine (Cl) Max.XX.XX%
 700.XX% Slowly Available Nitrogen from LESCO Poly Plus® Sulfur Coated Urea.
 Made in U.S.A. Produced under the following U.S. Patent 5,750,130 F1486

Active Ingredient:
 Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,5-dinitrobenzenamine* 1.50%
 Inert Ingredients: 98.50%
 Total 100.00%

*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,068,441.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand this label, find someone to explain it to you in detail.)
 See below for additional precautionary statements.

In case of emergency endangering life or property involving this product, call day or night, Chem-Trec 1-800-424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

Personal Protective Equipment (PPE):

- Applicators and other handlers must wear:
- Long-sleeved shirt and long pants
 - Chemical resistant gloves
 - Shoes plus socks
- Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at time of pesticide application. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter treated area until dusts have settled.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR RESIDUES.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
Storage: DO NOT store above 120°F. Store in original containers and keep closed; reseal opened bag by folding top down and securing. Store in a cool, dry place.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Observe all cautions and limitations on this label and on the labels of products used in combination with this herbicide. Do not use this herbicide other than in accordance with the instructions set forth on this label. The use of this herbicide not consistent with this label can result in injury to turf, animals, or persons. Keep Container closed to avoid spills and contamination.

GENERAL INFORMATION

This product is a combination pre-emergent herbicide and fertilizer formulated to provide pre-emergence control of most annual grasses and certain broadleaf weeds as they germinate in any turfgrass site (lawns, sod, turf areas). Examples of

turf injury. Seeding diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken turfgrass and increase the possibility of turf damage from this product.

APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). Check application routinely to determine proper calibration. Avoid overlaps that will increase rates above those recommended. Avoid application when winds may cause drift. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

SUGGESTED SPREADER SETTINGS

Spreader	Ground Speed	Spreading Width	Recommended Rates lb per 1000 sq ft (lb per acre)			
			1.5 (67)	2.3 (100)	5.4 (233)	
LESCO Rotary	3 mph	9 ft	Calibration Gauge	#XX	#XX	#XX
Cyclone® or Spyr®				XX	XX	XX

*Delivers 1.5 lb a/A and X.XX lb N/1,000 sq ft
 **Delivers 3.5 lb a/A and X.XX lb N/1,000 sq ft
 IMPORTANT: Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

TURFGRASS APPLICATION CHART

Turfgrass Species	Weeds	Rate of PRE-M 1.5% Plus Fertilizer	Comments:	
NORTHERN GRASSES Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail Poa annua barnyardgrass fall panicum oxalis prostrate spurge purslane knockweed evening primrose hop clover	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre) as the (initial application prior to weed germination in the spring.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67-100 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.	
	goosegrass	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre) in the spring prior to weed germination.	Apply a repeat application of 1.5 lb/1,000 sq ft (67 lb/acre)	
	cudweed Poa annua chickweed lawn burweed henbit corn speedwell	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre)	Apply in late summer or early fall prior to weed germination.	
	crabgrass foxtail Poa annua barnyardgrass fall panicum oxalis prostrate spurge purslane knockweed evening primrose hop clover	2.3 to 4.8 lb/1,000 sq ft (100-200 lb/acre) in the spring prior to weed germination.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67-100 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.	
	goosegrass	2.3 to 3.1 lb/1,000 sq ft (100-133 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 2.3 lb/1,000 sq ft (100 lb/acre) after a minimum of 4 weeks.	
	cudweed Poa annua chickweed lawn burweed henbit corn speedwell	2.3 to 3.1 lb/1,000 sq ft (100-133 lb/acre)	Apply in late summer or early fall prior to weed germination.	
	SOUTHERN GRASSES Bermudagrass St. Augustinegrass Bahia grass Buffalograss Zoysia grass Centipede grass Tall fescue	crabgrass foxtail Poa annua barnyardgrass fall panicum oxalis prostrate spurge purslane knockweed evening primrose hop clover	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67-100 lb/acre) after a minimum of 4 weeks if necessary.
		goosegrass	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre) prior to weed germination in the spring. Apply a repeat application 2.3 to 3.1 lb/1,000 sq ft (100-133 lb/acre) after minimum of 4 weeks.	An additional application of 1.5 lb/1,000 sq ft (67 lb/acre) may be applied for extended goosegrass control a minimum of 4 weeks after the second application.
		cudweed Poa annua chickweed lawn burweed henbit corn speedwell	2.3 to 5.4 lb/1,000 sq ft (100-233 lb/acre)	Apply in late summer or early fall prior to weed germination.

*Not for use on grasses or less
 The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 30 days, erratic weed control may result. This product can be applied before or after post-emergence herbicides used to control emerged weeds in turfgrasses. For annual grass control applications can be made with Atrazine Extra® or MSMA to control emerged weeds. Broadleaf weeds can be controlled using Trifluralin, Trifluralin, 2,4-D, and other similar products.

TURFGRASS PRECAUTIONS

Use only on well established turfgrass with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times. On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application. Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species. Do not use on bentgrass or Poa annua (greens and tees) or injury may occur. Delay reseeding or winter overseeding treated turfgrass for at least 2 months following application. Do not sprig turfgrass for 3 months after application.

INDUSTRIAL (unimproved) TURF

Industrial, or unimproved turf areas often have a different spectrum of weeds to be controlled than those found in fine turf as described elsewhere in this leaflet. This product will control annual grasses and broadleaf weeds mentioned in that section of this label as well as the following weeds that might germinate in established grasses in rights-of-way, roadsides, construction sites, parks, substations, lots, or similar areas:
 Crowfootgrass, fitchgrass, Johnsongrass (from seed), Junglerice, Lovegrass, Browntop Panicum, Texas Panicum, Field Sandbur, Signalgrass, Mexican Sprangletop, Red Sprangletop, Wishgrass, Woolly Cupgrass, Carpetweed, Fiddleneck, Flaree, Kochia, Lambquarters, Pigweed, Purslane, Florida Pusley, London Rocket, Shepherd's purse, Pennsylvania Smartweed, Annual Spurge, and Velvetleaf.

Apply before weeds germinate. A post-emergence herbicide such as 2,4-D, MSMA, or similar product is recommended to control established weeds.

#XXXXXX (front)

Net Weight: XX lb (XX kg)

ORNAMENTALS - RECOMMENDED SPECIES

PRE-M 1.5% Plus Fertilizer is safe to use around and over the top of the established plants listed below.

This product may be used on plant species not listed on this label. The suitability for such uses should be determined by treating a small number of such plants at the recommended rate. Treated plants should be evaluated 1-2 months following treatment for possible injury. THE USER ASSUMES RESPONSIBILITY FOR ANY CROP DAMAGE OR OTHER LIABILITY. Do not treat plants grown for food or feed. Do not use treated plants for food or feed.

Table with 4 columns: Common Name, Scientific Name, Common Name, Scientific Name. Lists various trees and shrubs such as Alder, European Black, Apple, Arbutus, American, etc.

Table with 4 columns: Common Name, Scientific Name, Common Name, Scientific Name. Lists various ground covers and bedding plants such as Acacia, Adonis, Agapanthus, etc.

ORNAMENTAL BULBS

This product may be applied for control of susceptible annual weeds in ornamental bulbs listed on the label (crocus, daffodil, narcissus), gladiolus, tulip, etc.). Apply this product prior to, during or after bulb emergence. If weeds have already germinated use a labeled post-emergence herbicide to control emerged weeds.

WILDFLOWERS

This product may be applied for control of susceptible annual weeds in plantings of wildflowers listed on the label (Black-eyed Susan, California Poppy, Coreopsis, Oxyeye Daisy, etc.). This product may be applied to established perennial wildflowers before emergence of weeds or wildflowers. For wildflowers being established from seed, apply this product after wildflowers have emerged but prior to weed germination. If weeds have already germinated, add a labeled post-emergence product to control emerged weeds.

ORNAMENTAL PRECAUTIONS

Apply this product to established plantings. DELAY applying this product to seedbeds, transplant beds, or liners until plants have become well rooted. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of this product with roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Plant only those desirable plant species listed on this label into soil treated the previous season with this product or injury may occur. For container grown ornamentals, delay first application of this product to bareroot liners 2-4 weeks.

LANDSCAPE AND GROUNDS MAINTENANCE

This product can be incorporated into landscape and grounds maintenance programs to provide extended pre-emergence control of most annual grasses and certain broadleaf weeds. Areas to be treated, such as much beds, parking areas and roadways, fencelines and borders, around statutory or monuments, and similar areas, should be free of emerged weeds before application. To remove emerged weeds, treat the area with a post-emergence product labeled for such use. Refer to APPLICATION RATE TABLE for rates. Avoid unintentional contact of granules with stone, wood, or other porous surfaces as staining may occur. Sweep immediately to avoid staining. If granules are crushed into surface, then rinse thoroughly.

NON-BEARING FRUIT AND NUT CROPS AND VINEYARDS

This product may be applied for pre-emergence control of most annual grasses and certain broadleaf weeds on the following non-bearing crops:

Table with 4 columns: Crop Name, Citrus, Olive, Pistachio. Lists crops like Almond, Apple, Fig, Peach, Plum, Apricot, Grape, Pear, Prune, Cherry, Naclarine, Pecan, Walnut, English.

NON-CROPLAND AREAS INCLUDING TREE PLANTATIONS

This product is recommended for grounds maintenance in non-cropland areas; pre-emergence control of the weed species listed in and around established tree plantations (including Christmas trees); in and around established ornamentals planted in non-cropland areas such as highway rights-of-way, and utility substations. This product may be used for hardwood and conifer regeneration on conservation reserve program land or similar areas. Refer to APPLICATION RATE TABLE for rates. This product may be applied at planting or to established trees. When making an application at planting, it is important that slit closure be achieved to prevent this product from directly contacting the tree roots or being washed into the root zone via the open slit or root stunting may occur.

TOTAL VEGETATIVE CONTROL

This product may be mixed with a granular formulation, or used in sequence with any formulation, of ARSENAL, Roundup, Karmex, Finale, Oust, diuron, or other products to provide bare ground, or total vegetation control. This product can be used to provide greater plant selectivity in areas where such action may be desired. Such sites might have roots of landscape vegetation, ornamentals, or desirable trees encroaching into the treated zone. Applications may be made to existing weeds controlled by the partner herbicide. Recommended rates should be determined from the product labels prior to use. Follow the most restrictive label instructions. For Kochia: Combinations of this product with ARSENAL or diuron are recommended if control has been a problem for other herbicides.

APPLICATION RATE TABLE

For pre-emergence control of the weed species listed using broadcast spreader equipment, apply PRE-M 1.5% Plus Fertilizer at the following rates:

Table with 3 columns: Length of Control, Pounds required to treat one acre, Pounds required to treat 1000 sq ft. Shows rates for Short Term (2-4 months) and Long Term (6-8 months).

Rate settings for a specific type of spreader device must be determined by applicator. To deliver the proper rate (3.1-6.1 lb/1000 sq ft) calibrate prior to application of product.

One bag (50 lb) will cover 1/3 acre (16,130 sq ft) at a rate of 3.1 lb per 1000 sq ft. The efficacy of this product will improve if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 30 days, erratic weed control may result. This product will not control established weeds.

If weeds should develop prior to activation of herbicide, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. This product will be used before or after applications of herbicides registered for post-emergence use (i.e. Roundup or Finale) for the control of established weeds. Do not apply sprays containing Roundup or Finale over the top of desirable plants. A PRE-M 1.5% Plus Fertilizer treatment may be followed by any registered herbicide to control weeds not listed on this label.

WEED SPECIES CONTROLLED

This product is recommended for pre-emergence control of the weed species listed. Applications can be made around and over the top of the ornamentals and to the sites listed on this label.

Table with 4 columns: Common Name, Scientific Name, Common Name, Scientific Name. Lists various grasses and broadleaf weeds controlled such as Barnyardgrass, Bluegrass, Annual, Crabgrass, etc.

WARRANTY

LESCO, Inc. warrants that this herbicide conforms to the chemical description on this label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, matters with other chemicals, and other factors affecting the use of this product are beyond the control of the manufacturer, the user assumes the responsibility for any crop damage or other liability resulting from the use of this product contrary to label directions or other conditions which are specified or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept, 15885 Sprague Rd., Strongsville, OH 44136-1772, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-a1.com>

Finale is a registered trademark of Aventis. Trimec is a registered trademark of PBI Gordon Corp. Roundup is a registered trademark of Monsanto Company. ARSENAL is a trademark of BASF. Karmex is a registered trademark of E. I. DuPont de Nemours and Company. Cyclone and Spiker are registered trademarks of Spiker Spreaders, LLC. LESCO and PRE-M are registered trademarks of LESCO Technologies, LLC.

G:REGULWPSTRONGSVIAprv1 Lbl - WorldControl Products/PRE-MXXXXXX.doc Rev: XXXXXX WBW (12/2002) E.P.A. Reg No. 10404-08 E.P.A. Est No. 10404-0H-04 (M, 10404-FL-01(S)) Net Weight: XX lb (XX kg) First label of lot code indicates manufacturing site

Refer to the APPLICATION RATE TABLE for recommended amount to use. *After planting bedding plants and before this product is applied, area to be treated should be watered sufficiently to pack soil around roots.



F1406 Made by LESCO, Inc. 15885 Sprague Road Strongsville, Ohio 44136-1772

758073XXXXX