



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 2, 2020

Viviana Torres
Assistant Regulatory Manager
LESCO, Inc
1385 East 36th Street
Cleveland, OH 44114-4114

Subject: Notification per PRN 98-10 – Addition of optional marketing claims
Product Name: LESCO PRE-M 1.5% PLUS FERTILIZER
EPA Registration Number: 10404-98
Application Date: May 20, 2019
Decision Number: 552554

Dear Ms. Torres:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

The primary brand name has been corrected to “**LESCO PRE-M 1.5% PLUS FERTILIZER**”.

If you have any questions, please contact Curtis Hildebrandt at 703-347-8198 or by email at hildebrandt.curtis@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Emily Schmid".

Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

10404-98

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

3/2/2020

PENDIMETHALIN

GROUP

3

HERBICIDE

LESCO® PRE-M® 1.5% Plus Fertilizer

PRE-EMERGENT WEED CONTROL

For Use on Turfgrasses, Ornamentals, Landscape or Grounds Maintenance, and Non-cropland Areas.

Contains LESCO® Poly Plus® Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.

Active Ingredient:

Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine* 1.50%

Other Ingredients: 98.50%

Total 100.00%

*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441.

X-X-X

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)XX.XX%

XX.XX% Ammoniacal Nitrogen

XX.XX% Urea Nitrogen*

XX.XX% Water Insoluble Nitrogen

XX.XX% Water Soluble Controlled Release Nitrogen (N)

AVAILABLE PHOSPHATE (P₂O₅)XX.XX%

SOLUBLE POTASH (K₂O)XX.XX%

MAGNESIUM (Mg) TotalXX.XX%

XX.XX% Water Soluble Magnesium (Mg)

SULFUR (S) TotalXX.XX%

XX.XX% Free Sulfur (S)

XX.XX% Combined Sulfur (S)

COPPER (Cu) TotalXX.XX%

XX.XX% Water Soluble Copper (Cu)

IRON (Fe) TotalXX.XX%

XX.XX% Water Soluble Iron (Fe)

MANGANESE (Mn) TotalXX.XX%

XX.XX% Water Soluble Manganese (Mn)

ZINC (Zn) TotalXX.XX%

XX.XX% Water Soluble Zinc (Zn)

DERIVED FROM:

CHLORINE (Cl) Max.XX.XX%

*X.XX% Slowly available Urea Nitrogen from Polymer Coated Sulfur Coated Urea.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See below for additional precautionary statements.

In case of emergency endangering life or property involving this product, call day or night, Chem-Trec 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:
Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters.

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions.

Resistance Management

For resistance management, this product contains Group 3 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural, biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of plants and weed biotypes.

Contact your local sales representative, or extension agent to find out if suspected resistant weeds to this product have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at time of pesticide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT apply more than 3.5 lb of this product per 1,000 sq ft per application, and no more than 9.5 lb of this product per 1,000 sq. ft. (6.00 lb. ai per acre) per year.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

• Coveralls • Chemical resistant gloves • Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter treated area until dusts have settled.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR RESIDUES.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures. Or call [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to "pesticide disposal" directions listed above.

Observe all cautions and limitations on this label and on the labels of products used in combination with this herbicide. Do not use this herbicide other than in accordance with the instructions set forth on this label. The use of this herbicide not consistent with this label can result in injury to turf, animals, or persons. Keep Container closed to avoid spills and contamination.

GENERAL INFORMATION

This product is a combination pre-emergent herbicide and fertilizer formulated to provide pre-emergence control of most annual grasses and certain broadleaf weeds as they germinate in any turfgrass site (lawns, sod, turf areas). Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multifamily dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries; golf courses; prairiegrass areas; and sod farms.

This product can be used in and around field, liner, and container grown ornamental nurseries; established landscape ornamentals and ornamental gardens; listed groundcovers and bulbs; nonbearing fruit and nut trees; conifer and hardwood seedling nurseries; and for plantation site preparation and maintenance.

In addition, this product can be applied for general grounds maintenance around areas such as parking lots, driveways and roadsides, alleyways, bike and jogging paths, vacant lots, buildings, stone gardens and gravel yards, markers and fence lines, mulch beds and other similar areas. It may be used under asphalt or concrete treatments as part of a site preparation program.

This product is recommended for pre-emergence control of most annual grasses and certain broadleaf weeds as they germinate in any non-cropland area such as railroad, utility, highway, and pipeline rights-of-way; highway guardrails, delineators, and sign posts; bridge abutments and approaches; utility substations; petroleum tank farms; pumping installations; storage areas; fence rows; windbreaks and shelterbelts; paved or gravel surfaces; established wildflower plantings, and other similar areas where weed control is desired.

This product controls weeds as they germinate but will not control established weeds. Therefore, areas to be treated should be free of established weeds. For the control of established weeds, this product may be used in conjunction with herbicides registered for post-emergence use. Consult the labels of those herbicides for use rates timings, and precautions or restrictions.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

The efficacy of this product will improve if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

Applied according to label directions and under normal growing conditions, this product will not cause turf injury. Over-application can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken turfgrass and increase the possibility of turf damage from this product.

APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). Check application routinely to determine proper calibration. Avoid overlaps that will increase rates above those recommended. Avoid application when winds may cause drift. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

SUGGESTED SPREADER SETTINGS

Recommended Rates in lb per 1,000 sq ft

Spreader	1.5 lb	2.3 lb*	3.1 lb	4.6 lb	5.4 lb**
LESCO Rotary Calibration Gauge	#XX	#XX	#XX	#XX	#XX
Cyclone® or Spyker®	XX	XX	XX	XX	XX

*Delivers 1.5 lb ai/A and X.XX lb N/1,000 sq ft

**Delivers 3.5 lb ai/A and X.XX lb N/1,000 sq ft

IMPORTANT: Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

TURFGRASS APPLICATION CHART

Turfgrass Species	Weeds	Rates of LESCO PRE-M 1.5% Plus Fertilizer	Comments:
NORTHERN GRASSES Kentucky Bluegrass Perennial Ryegrass Fine Fescue Tall Fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre) as the initial application prior to weed germination in the spring.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67 – 100 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre) in the spring prior to weed germination.	Apply a repeat application of 4.0 lb/1,000 sq ft (174 lb/acre)
	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre)	Apply in late summer or early fall prior to weed germination.
Bentgrass and established <i>Poa annua</i> * (1/2 inch height or taller)	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane	2.3 lb to 4.6/1,000 sq ft (100-200 lb/acre) in the spring prior to weed germination.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67 - 100 lb/acre) after a minimum of 4 weeks for extended control or where heavy weed infestations

	knotweed evening primrose hop clover		are expected.
	goosegrass	2.3 lb to 3.1/1,000 sq ft (100-133 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 2.3 lb/1,000 sq ft (100 lb/acre) after a minimum of 4 weeks.
	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.3 to 3.1 lb/1,000 sq ft (100-133 lb/acre)	Apply in late summer or early fall prior to weed germination.
SOUTHERN GRASSES Bermudagrass* St. Augustinegrass Bahia grass Zoysiagrass Centipedegrass Tall Fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre) prior to weed germination in the spring.	Apply a repeat application of 1.5 to 2.3 lb/1,000 sq ft (67 – 100 lb/acre) after a minimum of 4 weeks if necessary.
	goosegrass	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre) prior to weed germination in the spring. Apply a repeat application of 2.3 to 3.1 lb/1,000 sq ft (100 – 233 lb/acre) after a minimum of 4 weeks.	An additional application of 1.5 lb/1,000 sq ft (67 lb/acre) may be applied for extended goosegrass control a minimum of 4 weeks after the second application.
	cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell	2.3 to 5.4 lb/1,000 sq ft (100 – 233 lb/acre)	Apply in late summer or early fall prior to weed germination.

*Not for use on greens or tees.

The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If this product is not activated by rainfall or irrigation within 30 days, erratic weed control may result. This product can be applied before or after post-emergence herbicides used to control emerged weeds in turfgrasses. For annual grass control applications can be made with Acclaim Extra 1 or MSMA to control emerged weeds. Broadleaf weeds can be controlled using Trimec2, Three-Way3, 2,4-D, and other similar products.

TURFGRASS PRECAUTIONS

Use only on well established turfgrass with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times. On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application.

Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species.

Do not use on bentgrass or *Poa annua* (greens and tees) or injury may occur.

Delay reseeding or winter overseeding treated turfgrass for at least 2 months following application. Do not sprig turfgrass for 3 months after application.

INDUSTRIAL (unimproved) TURF

Industrial, or unimproved turf areas often have a different spectrum of weeds to be controlled than those found in fine turf as described elsewhere in this leaflet. This product will control annual grasses and broadleaf weeds mentioned in that section of this label as well as the following weeds that might germinate in established grasses in rights-of-way, roadsides, construction sites, parks, substations, lots, or similar areas:

Crowfootgrass, Itchgrass, Johnsongrass (from seed), Junglerice, Lovegrass, Browntop Panicum, Texas Panicum, Field Sandbur, Signalgrass, Mexican Sprangletop, Red Sprangletop, Witchgrass, Woolly Cupgrass, Carpetweed, Fiddleneck, Filaree, Kochia, Lambsquarters, Pigweed, Puncturevine, Florida Pusley, London Rocket, Shepherd's purse, Pennsylvania Smartweed, Annual Spurge, and Velvetleaf.

Apply before weeds germinate. A post-emergence herbicide such as 2,4-D, MSMA, or similar product is recommended to control established weeds.

ORNAMENTALS - RECOMMENDED SPECIES

PRE-M 1.5% Plus Fertilizer is safe to use around and over the top of the established plants listed below.

Common Name	Scientific Name	Common Name	Scientific Name
Trees			
Alder, European black	<i>Alnus glutinosa</i>	Oak, Chinquapin	<i>Quercus muehlenbergii</i>
Apple	<i>Malus</i> spp.	Oak, Live	<i>Quercus virginiana</i>
Arborvitae, American	<i>Thuja occidentalis</i>	Oak, Pin	<i>Quercus palustris</i>
Arbutus	<i>Arbutus</i> spp.	Oak, Red	<i>Quercus rubra</i>
Ash, Red	<i>Fraxinus pennsylvanica</i>	Oak, Swamp chestnut	<i>Quercus michauxii</i>
Ash, White	<i>Fraxinus americana</i>	Oak, Water	<i>Quercus nigra</i>
Aspen, Bigtooth	<i>Populus grandidentata</i>	Oak, White	<i>Quercus alba</i>
Aspen, Quaking	<i>Populus tremuloides</i>	Olive	<i>Olea europaea</i>
Basswood	<i>Tilia</i> spp.	Palm, Date	<i>Phoenix</i> spp.
Birch, European Weeping	<i>Betula pendula</i>	Palm, Fan	<i>Washingtonia</i> spp.
Birch, River	<i>Betula nigra</i>	Palm, Pindo	<i>Butia</i> spp.
Cedar, White	<i>Thuja occidentalis</i>	Palm, Washington	<i>Washingtonia</i> spp.
Chamaecyparis, Boulevarde	<i>Chamaecyparis pisifera</i>	Peach	<i>Prunus persica</i>
Cherry, Black	<i>Prunus serotina</i>	Pear, Bradford	<i>Pyrus calleryana</i> 'Bradford'
Cherry, Choke	<i>Prunus virginiana</i>	Pecan	<i>Carya illinoensis</i>
Cherry, Kwanzan	<i>Prunus serrulata</i>	Pine, Austrian	<i>Pinus nigra</i>
Cherry, Nanking	<i>Prunus tomentosa</i>	Pine, Italian Stone	<i>Pinus pinea</i>
Cottonwood	<i>Populus deltoides</i>	Pine, Loblolly	<i>Pinus taeda</i>
Crabapple	<i>Malus</i> spp.	Pine, Monterey	<i>Pinus radiata</i>
Crepe Myrtle	<i>Lagerstroemia indica</i>	Pine, Red	<i>Pinus resinosa</i>
Cryptomeria, Japanese cedar	<i>Cryptomeria japonica</i>	Pine, Scotch	<i>Pinus sylvestris</i>
Cypress, Bald	<i>Taxodium distichum</i>	Pine, Slash	<i>Pinus elliotii</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	Pine, Virginia	<i>Pinus virginiana</i>
Dogwood, Flowering	<i>Cornus florida</i>	Pine, White	<i>Pinus strobus</i>
Dogwood, Korean	<i>Cornus kousa</i>	Plum, Purple Leaf	<i>Prunus cerasifera</i>
Dogwood, Silky	<i>Cornus amomum</i>	Poplar, Black	<i>Populus nigra</i>
Dogwood, Shrub	<i>Cornus</i> spp.	Redcedar, Eastern	<i>Juniperus virginiana</i>
Elm	<i>Ulmus japonica</i>	Redcedar, Western	<i>Thuja plicata</i>
Fir, Balsam	<i>Abies balsamae</i>	Red Ironbark	<i>Eucalyptus sideroxylon</i> 'Rosea'
Fir, Douglas	<i>Pseudotsuga menziesii</i>	Redwood, Dawn	<i>Metasequoia glyptostroboides</i>
Fir, Fraser	<i>Abies fraseri</i>	Sequoia, Giant	<i>Sequoiadendron giganteum</i>
Fir, White	<i>Abies concolor</i>	Serviceberry	<i>Amelanchier laevis</i>
Franklinia	<i>Franklinia</i> spp.	Spruce, Norway	<i>Picea abies</i>
Ginkgo	<i>Ginkgo biloba</i>	Spruce, White	<i>Picea glauca</i>
Gum, Black	<i>Nyssa sylvatica</i>	Sweetgum	<i>Liquidambar styraciflua</i>
Gum, Sour	<i>Nyssa sylvatica</i>	Sycamore	<i>Platanus occidentalis</i>
Haw, Black	<i>Viburnum prunifolium</i>	Trachycarpus	<i>Trachycarpus</i> spp.
Hawthorn	<i>Crataegus</i> spp.	Tulip tree	<i>Liriodendron tulipifera</i>
Hemlock, Canada	<i>Tsuga canadensis</i>	Walnut, Black	<i>Juglans nigra</i>
Hemlock, Eastern	<i>Tsuga canadensis</i>	Willow, Weeping	<i>Salix babylonica</i>
Holly, American	<i>Ilex opaca</i>	Yellowwood	<i>Cladrastis lutea</i>
Honeylocust	<i>Gleditsia triacanthos</i>		
Lilac, Common	<i>Syringa vulgaris</i>		
Lilac, Japanese Tree	<i>Syringa reticulata</i>		
Linden	<i>Tilia</i> spp.		
Magnolia, Saucer	<i>Magnolia soulangiana</i>		
Magnolia, Southern	<i>Magnolia grandiflora</i>		
Magnolia, Star	<i>Magnolia stellata</i>		
Maidenhair Tree	<i>Ginkgo biloba</i>		
Maple, Norway	<i>Acer platanoides</i>		
Maple, Japanese	<i>Acer palmatum</i>		
Maple, Red	<i>Acer rubrum</i>		
Maple, Sugar	<i>Acer saccharum</i>		
Shrubs			
Abelia, Glossy	<i>Abelia grandiflora</i>	Leucothoe, Coast	<i>Leucothoe axillaris</i>
Aucuba, Gold	<i>Aucuba japonica</i>	Lilac, Cut-leaf	<i>Syringa laciniata</i>

Azalea	<i>Rhododendron</i> spp.	Lily-of-the-Nile	<i>Agapanthus africanus</i>
Bamboo, Heavenly	<i>Nandina domestica</i>	Mahonia	<i>Mahonia aquifolium</i>
Barberry	<i>Berberis gladywensis</i>	Mock Orange	<i>Pittosporum tobira</i>
Barberry, Japanese	<i>Berberis thunbergii</i>	Myrtle, Compact	<i>Myrtus communis</i>
Blue Indigo Bush	<i>Dalea gregii</i>	Myrtle, Wax	<i>Myrica cerifera</i>
Bottlebrush, Lemon	<i>Callistemon citrinus</i>	Nandina	<i>Nandina domestica</i>
Boxwood, Common	<i>Buxus sempervirens</i>	Oleander	<i>Nerium oleander</i>
Boxwood, Japanese	<i>Buxus microphylla</i>	Oregon Grape	<i>Mahonia aquifolium</i>
Brittlebush	<i>Encelia farinosa</i>	Osmanthus	<i>Osmanthus fragrans</i>
Buttonbush	<i>Cephalanthus occidentalis</i>	Palm, European Fan	<i>Chamaerops humilis</i>
Camellia	<i>Camellia japonica</i>	Palm, Mediterranean Fan	<i>Chamaerops</i> spp.
Cape jasmine	<i>Gardenia jasminoides</i>	Phlox, Prickly	<i>Leptodactylon californicum</i>
Cassia, Feathery	<i>Cassia artemisioides</i>	Photinia, Fraser	<i>Photinia x Fraseri</i>
Cordylone	<i>Cordylone</i> spp.	Pteris, Japanese	<i>Pteris japonica</i>
Correa	<i>Correa</i> spp.	Pine, Mugo	<i>Pinus mugo</i>
Cotoneaster	<i>Cotoneaster apiculatus</i>	Plum, Natal	<i>Carissa grandiflora</i>
Cotoneaster, Bearberry	<i>Cotoneaster dammeri</i>	Privet, California	<i>Ligustrum ovalifolium</i>
Cotoneaster, Rock	<i>Cotoneaster horizontalis</i>	Privet, Glossy	<i>Ligustrum lucidum</i>
Cypress, Italian	<i>Cupressus sempervirens</i>	Privet, Variegated	<i>Ligustrum sinensis</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	Privet, Waxleaf	<i>Ligustrum japonicum</i>
Deutzia, Slender	<i>Deutzia gracilis</i>	Pyracantha	<i>Pyracantha coccinea</i>
Dogwood, Red Twig	<i>Cornus sericea</i>	Quince, Flowering	<i>Chaenomeles japonica</i>
Elaeagnus	<i>Elaeagnus ebbingei</i>	Ranger, Texas	<i>Leucophyllum frutescens</i>
Escallonia	<i>Escallonia fradesii</i>	Redroot	<i>Ceanothus</i> spp.
Euonymus	<i>Euonymus fortunei</i>	Rhododendron	<i>Rhododendron</i> spp.
Euonymus, Golden	<i>Euonymus japonica</i>	Robira	<i>Pittosporum tobira</i>
Euonymus, Winged	<i>Euonymus alata</i>	Rose	<i>Rosa</i> spp.
Firethorn	<i>Pyracantha coccinea</i>	Spice Plant	<i>Illicium parviflorum</i>
Forsythia, Border	<i>Forsythia intermedia</i>	Spiraea	<i>Spiraea vanhouttei</i>
Fragrant Olive	<i>Osmanthus fragrans</i>	Spiraea, Anthony Waterer	<i>Spiraea x bumalda</i>
Fuschia, California	<i>Zauschneria californica</i>	Spiraea, Japanese	<i>Spiraea japonica</i>
Gardenia	<i>Gardenia jasminoides</i>	Sweet Bay	<i>Laurus nobilis</i>
Hawthorne, Indian	<i>Raphiolepis indica</i>	Trumpet Bush	<i>Tecoma stans</i>
Hibiscus	<i>Hibiscus syriacus</i>	Verbena, Lemon	<i>Aloysia triphylla</i>
Holly, Chinese	<i>Ilex cornuta</i>	Viburnum	<i>Viburnum suspensum</i>
Holly, Japanese	<i>Ilex crenata</i>	Vitex	<i>Vitex</i> spp.
Holly, Fosters	<i>Ilex attenuata</i> 'Fosteri'	Weigela	<i>Weigela florida</i>
Holly, Savannah	<i>Ilex attenuata</i>	Wild Lilac	<i>Ceanothus</i> spp.
Holly, Yaupon	<i>Ilex vomitoria</i>	Xylosma	<i>Xylosma congestum</i>
Holly, Yaupon	<i>Ilex vomitoria</i>	Yellowbells	<i>Tecoma stans</i>
Honeysuckle, Bush	<i>Diervilla lonicera</i>	Yew	<i>Taxus media</i>
Hopseed Bush	<i>Dodonaea viscosa</i>	Yew, Japanese	<i>Taxus cuspidata</i>
Hopbush	<i>Dodonaea viscosa</i>	Yew, Southern	<i>Podocarpus macrophyllus</i>
Hydrangea	<i>Hydrangea macrophylla</i>	Yucca, Adam's Needle	<i>Yucca filamentosa</i>
Juniper	<i>Juniperus</i> spp.	Yucca, Weeping	<i>Yucca pendula</i>
Juniper, Chinese	<i>Juniperus chinensis</i> v. <i>pfitzer</i>		
Juniper, Shore	<i>Juniperus conferta</i>		
Juniper, Trailing	<i>Juniperus horizontalis</i>		
Laurel, Cherry	<i>Prunus laurocerasus</i>		
Laurel, Mountain	<i>Kalmia latifolia</i>		
Laurel, Otto Luyken	<i>Prunus laurocerasus</i>		
Laurel, Schipka	<i>Prunus schipkanensis</i>		
Laurustinus	<i>Viburnum tinus</i>		
Lavender, English	<i>Lavandula angustifolia</i>		
Leucothoe	<i>Leucothoe fontanesiana</i>		

Perennials			
Acacia	<i>Acacia redolens</i>	Ground Covers	
Asparagus	<i>Asparagus</i> spp.	Ajuga	<i>Ajuga reptans</i>
Aster, New York	<i>Aster novi-belgii</i>	Baby Sun Rose	<i>Aptenia cordifolia</i>
Aster, Stokes	<i>Stokesia laevis</i>	Beach Strawberry	<i>Fragaria chiloensis</i>
Babys Breath	<i>Gypsophila elegans</i>	Capeweed	<i>Arctotheca calendula</i>
Beard-Tongue	<i>Penstemon</i> spp.	Cinquefoil, Spring	<i>Potentilla verna</i>
Bellflower	<i>Campanula</i> spp.	Coyotebrush, Dwarf	<i>Baccharis pitularis</i>
Bird of Paradise	<i>Caesalpinia pulcherrima</i>	Daisy, Trailing African	<i>Osteospermum fruticosum</i>
Black-eyed Susan	<i>Rudbeckia hirta</i>	Dymondia	<i>Dymondia margaretae</i>
Blanket Flower	<i>Gaillardia X grandiflora</i>	Gazania	<i>Gazania splendens</i>
Bleeding Heart	<i>Dicentra spectabilis</i>	Icelandic, Large Leaf	<i>Carpobrotus edulis</i>
California Poppy	<i>Eschscholzia californica</i>	Ivy, English	<i>Hedera helix</i>
Calla lily	<i>Zantedeschia aethiopica</i>	Ivy, Geranium	<i>Pelargonium peltatum</i>
Canna, common garden	<i>Canna generalis</i> 'Lucifer'	Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>
Carex	<i>Carex</i> spp.	Jasmine, Primrose	<i>Jasminum mesnyi</i>
Chincherinchee	<i>Ornithogalum thyrsoides</i>	Jessamine, Carolina	<i>Gelsemium sempervirens</i>
Clover, Crimson	<i>Trifolium incarnatum</i>	Manzanita, Bearberry	<i>Arctostaphylos uva-ursi</i>
Columbine	<i>Aquilegia 'McKana Giant'</i>	Miscanthus	<i>Miscanthus</i> spp.
Coreopsis	<i>Coreopsis lanceolata</i>	Mondograss	<i>Ophiopogon japonica</i>
Crinum Lily	<i>Crinum</i> spp.	Morningglory	<i>Convolvulus</i> spp.
Crocus	<i>Crocus</i> spp.	Myoporum	<i>Myoporum parvifolium</i>
Daffodil	<i>Narcissus</i> spp.	Pachysandra	<i>Pachysandra terminalis</i>
Daylily	<i>Hemerocallis</i> spp.	Potentilla	<i>Potentilla fruticosa</i>
Fairy Duster	<i>Calliandra eriophylla</i>	Red Apple	<i>Aptenia cordifolia</i>
Fern, Asparagus	<i>Asparagus officinalis</i>	Rosemary	<i>Rosemarinus officinalis</i>
Fern, Boston	<i>Nephrolepis exaltata</i>	Rose-Of-Sharon	<i>Hypericum calycinum</i>
Fern, Leatherleaf	<i>Rumohra adiantiformis</i>	Sand Strawberry	Sand Strawberry
Fortnight Lily	<i>Moraea</i> spp.	Sedum	<i>Sedum spurium</i>
Foxglove	<i>Digitalis purpurea</i>	St. Johnswort, creeping	<i>Hypericum calycinum</i>
Freesia	<i>Freesia X hybrida</i>	Stonecrop	<i>Sedum spurium</i>
Gaillardia	<i>Gaillardia pulchella</i>	Verbena, Peruvian	<i>Verbena peruviana</i>
Geum	<i>Geum</i> spp.	Vervain	<i>Verbena peruviana</i>
Gladiolus	<i>Gladiolus</i> spp.	Vetch, Crown	<i>Vicia sativa</i>
Heather, Dwarf	<i>Calluna vulgaris</i>	Vinca	<i>Vinca minor</i>
Hosta	<i>Hosta</i> spp.	Wintercreeper	<i>Euonymus fortunei</i>
Indian Blanket	<i>Gaillardia pulchella</i>	Ornamental Grasses	
Lantana, Weeping	<i>Lantana montevidensis</i>	Beach Grass	<i>Ammophila breviligulata</i>
Leopards Bane	<i>Doronicum cordatum</i>	Fescue, Blue	<i>Festuca ovina</i>
Lily	<i>Lilium</i> spp.	Fescue, Sheep	<i>Festuca ovina</i>
Liriope, Big Blue	<i>Liriope muscari</i>	Fountain Grass	<i>Pennisetum setaceum</i>
Liriope, Creeping	<i>Liriope spicata</i>	Pampas Grass	<i>Cortaderia selloana</i>
Liriope, Variegated	<i>Liriope muscari</i>	Reed Canary Grass	<i>Phalaris arundinacea</i>
Moonbeam	<i>Coreopsis verticillata</i>	Reed, Giant	<i>Arundo</i> spp.
Montbretia	<i>Crocsmia crocosmiiflora</i>	Ribbon Grass	<i>Phalaris arundinacea</i>
Nightshade	<i>Solanum</i> spp.	Tufted Hair Grass	<i>Deschampsia caespitosa</i>
Orchid, Peacock	<i>Acidanthera bicolor</i>	Bedding Plants *	
Oxeye Daisy	<i>Chrysanthemum leucanthemum</i>	Ageratum	<i>Ageratum houstonianum</i>
Peony, Chinese	<i>Paeonia lactiflora</i>	Alyssum	<i>Alyssum saxatile</i>
Purple Coneflower	<i>Echinacea purpurea</i>	Anemone, poppy-flowered	<i>Anemone coronaria</i>
Purple Loosestrife	<i>Lythrum virgatum</i>	Artemesia	<i>Artemesia</i> spp.
Rosemary	<i>Rosmarinus officinalis</i>	Balloonflower	<i>Platycodon grandiflorum</i>
Sedge	<i>Carex</i> spp.	Begonia	<i>Begonia</i> spp.
Shata Daisy	<i>Chrysanthemum X superbum</i>	Cabbage, Ornamental	<i>Brassica oleracea</i>
Sweet Flag	<i>Acorus calamus</i>	Caladium	<i>Caladium</i> spp.
Tickseed	<i>Coreopsis lanceolata</i>	Cast-Iron Plant	<i>Aspidistra elatior</i>
Tulip	<i>Tulipa</i> spp.	China Aster	<i>Callistephus chinensis</i>

Wisteria	<i>Wisteria</i> spp.	Crocsmia, Montebretia	<i>Crocsmia X crocosmiiflora</i>
Wonder Flower	<i>Ornithogalum thyrsoides</i>	Dahlia	<i>Dahlia</i> spp.
Yarrow	<i>Achillea millefolium</i>	Dusty Miller	<i>Senecio cineraria</i>
		Gayfeather	<i>Liatris</i> spp.
		Gazania Treasure Flower	<i>Gazania rigens</i>
		Gazania Trailing	<i>Gazania rigens leucolaena</i>
		Geranium	<i>Geranium</i> spp.
		Gloxinia	<i>Gloxinia simningia</i>
		Impatiens	<i>Impatiens</i> spp.
		Kale, Ornamental	<i>Brassica napus</i>
		Marigold, African	<i>Tagetes erecta</i>
		Moss Rose	<i>Portulaca grandiflora</i>
		Mum, Garden	<i>Chrysanthemum</i> spp.
		Pansy	<i>Viola tricolor</i>
		Periwinkle	<i>Vinca major</i>
		Periwinkle, Rose	<i>Catharanthus roseus</i>
		Petunia	<i>Petunia</i> spp.
		Plumosa Cockscomb	<i>Celosia cristata</i>
		Portulaca	<i>Portulaca grandiflora</i>
		Salvia	<i>Salvia splendens</i>
		Snapdragon	<i>Antirrhinum majus</i>
		Statice	<i>Limonium</i> spp.
		Sweet William	<i>Dianthus barbatus</i>
		Vinca	<i>Vinca major</i>
		Zinnia	<i>Zinnia</i> spp.

Refer to the **APPLICATION RATE TABLE** for recommended amount to use.

*After planting bedding plants and before this product is applied, area to be treated should be watered sufficiently to pack soil around roots.

This product may be used on plant species not listed on this label. The suitability for such uses should be determined by treating a small number of such plants at the recommended rate. Treated plants should be evaluated 1-2 months following treatment for possible injury. THE USER ASSUMES RESPONSIBILITY FOR ANY CROP DAMAGE OR OTHER LIABILITY. Do not treat plants grown for food or feed. Do not use treated plants for food or feed.

ORNAMENTAL BULBS

This product may be applied for control of susceptible annual weeds in ornamental bulbs listed on the label [crocus, daffodil (narcissus), gladiolus, tulip, etc.]. Apply this product prior to, during or after bulb emergence. If weeds have already germinated use a labeled post-emergence herbicide to control emerged weeds.

WILDFLOWERS

This product may be applied for control of susceptible annual weeds in plantings of wildflowers listed on the label (Black-eyed Susan, California Poppy, Coreopsis, Oxeye Daisy, etc.). This product may be applied to established perennial wildflowers before emergence of weeds or wildflowers. For wildflowers being established from seed, apply this product after wildflowers have emerged but prior to weed germination. If weeds have already germinated, add a labeled post-emergence product to control emerged weeds.

Due to the diversity of species and varieties which exist in areas where wildflowers are grown, the response to this product may vary greatly. Careful testing on desirable species is recommended to determine if area-wide applications can be made.

ORNAMENTAL PRECAUTIONS

Apply this product to established plantings. DELAY applying this product to seedbeds, transplant beds, or liners until plants have become well rooted. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of this product with roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. Plant only those desirable plant species listed on this label into soil treated the previous season with this product or injury may occur. For container grown ornamentals, delay first application of this product to bareroot liners 2-4 weeks.

LANDSCAPE AND GROUNDS MAINTENANCE

This product can be incorporated into landscape and grounds maintenance programs to provide extended pre-emergence control of most annual grasses and certain broadleaf weeds. Areas to be treated, such as mulch beds, parking areas and roadsides, fencelines and borders, around statuary or monuments, and similar areas, should be free of emerged weeds before application. To remove emerged weeds, treat the area with a post-emergence product labeled for such use.

Refer to APPLICATION RATE TABLE for rates. Avoid unintentional contact of granules with stone, wood, or other porous surfaces as staining may occur. Sweep immediately to avoid staining. If granules are crushed into surface, then rinse thoroughly.

NON-BEARING FRUIT AND NUT CROPS AND VINEYARDS

This product may be applied for pre-emergence control of most annual grasses and certain broadleaf weeds on the following non-bearing crops:

Almond	Citrus	Olive	Pistachio
Apple	Fig	Peach	Plum
Apricot	Grape	Pear	Prune
Cherry	Nectarine	Pecan	Walnut, English

Refer to APPLICATION RATE TABLE for rates.

Apply this product directly to the ground below the trees or vines. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of this product and roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and plant tolerance. For newly transplanted and one year old grapevines, apply only when they are dormant. Do not apply if buds have started to swell. This product may be used where the roots of a fruit, vine, nut, or ornamental plant encroach into a treatable area.

TOTAL VEGETATIVE CONTROL

This product may be mixed with a granular formulation, or used in sequence with any formulation, of ARSENAL, Roundup, Karmex, Finale, Oust, diuron, or other products to provide bare ground, or total vegetation control. This product can be used to provide greater plant selectivity in areas where such action may be desired. Such sites might have roots of landscape vegetation, ornamentals, or desirable trees encroaching into the treated zone.

Applications may be made to existing weeds controlled by the partner herbicide. Recommended rates should be determined from the product labels prior to use. Follow the most restrictive label instructions.

For Kochia: Combinations of this product with ARSENAL or diuron are recommended if control has been a problem for other herbicides.

APPLICATION RATE TABLE

For pre-emergence control of the weed species listed using broadcast spreader equipment, apply **LESCO PRE-M 1.5% Plus Fertilizer** at the following rates:

Length of Control	Pounds required to treat one acre	Pounds required to treat 1,000 sq. ft.
Short Term (2-4 months)	133lb	3.1lb
Long term (6-8 months)	267lb	6.1lb

Rate settings for a specific type of spreader device must be determined by applicator. To deliver the proper rate (3.1 -6.1 lb/1000 sq ft) calibrate prior to application of product.

One bag (50 lb) will cover 1/3 acre (16,130 sq ft) at a rate of 3.1 lb per 1000 sq ft.

The efficacy of this product will improve if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If This product is not activated by rainfall or irrigation within 30 days, erratic weed control may result. This product will not control established weeds.

If weeds should develop prior to activation of herbicide, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. This product may be used before or after applications of herbicides registered for post-emergence use (i.e. Roundup or Finale) for the control of established weeds. Do not apply sprays containing Roundup or Finale over the top of desirable plants. A **LESCO PRE-M 1.5% Plus Fertilizer** treatment may be followed by any registered herbicide to control weeds not listed on this label.

WEED SPECIES CONTROLLED

This product is recommended for pre-emergence control of the weed species listed. Applications can be made around and over the top of the ornamentals and to the sites listed on this label.

Grasses Controlled		Broadleaf Weeds Controlled	
Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>	Burweed, Lawn	<i>Soliva pterosperma</i>
Bluegrass, Annual	<i>Poa annua</i>	Carpetweed	<i>Mollugo verticillata</i>
Crabgrass	<i>Digitaria</i> spp.	Chickweed, Common	<i>Stellaria media</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	Chickweed, Mouseear	<i>Cerastium vulgatum</i>
Foxtail, Giant	<i>Setaria faberi</i>	Clover, Hop	<i>Trifolium procumbens</i>
Foxtail, Green	<i>Setaria viridis</i>	Cudweed	<i>Gnaphalium</i> spp.
Foxtail, Yellow	<i>Setaria glauca</i>	Eveningprimrose	<i>Oenothera biennis</i>
Goosegrass	<i>Eleusine indica</i>	Fiddleneck	<i>Amsinckia intermedia</i>
Itchgrass	<i>Rottboellia exaltata</i>	Filaree	<i>Erodium</i> spp.
Johnsongrass (from seed)	<i>Sorghum halepense</i>	Henbit	<i>Lamium amplexicaule</i>
Junglerice	<i>Echinochloa colona</i>	Knotweed, prostrate	<i>Polygonum aviculare</i>
Lovegrass (from seed)	<i>Eragrostis</i> spp.	Kochia	<i>Kochia scoparia</i>
Panicum, Browntop	<i>Panicum fasciculatum</i>	Lambsquarters	<i>Chenopodium album</i>
Panicum, Fall	<i>Panicum dichotomiflorum</i>	Pigweed	<i>Amaranthus</i> spp.
Panicum, Texas	<i>Panicum texanum</i>	Puncturevine	<i>Tribulus terrestris</i>
Sandbur, Field	<i>Cenchrus incertus</i>	Purslane	<i>Portulaca oleracea</i>
Signalgrass	<i>Brachiaria platyphylla</i>	Pusley, Florida	<i>Richardia scabra</i>
Sprangletop, Mexican	<i>Leptochloa uninervia</i>	Rocket, London	<i>Sisymbrium irio</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>	Shepherd's purse	<i>Capsella bursa-pastoris</i>
Witchgrass	<i>Panicum capillare</i>	Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Woolly Cupgrass	<i>Eriochloa villosa</i>	Speedwell, Corn	<i>Veronica arvensis</i>
		Spurge, Annual	<i>Euphorbia</i> spp.
		Spurge, Prostrate	<i>Euphorbia humistrata</i>
		Woodsorrel, Yellow	<i>Oxalis stricta</i>
		Velvetleaf (Buttonweed)	<i>Abutilon theophrasti</i>

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>. Poly Plus is comprised of Polymer Coated Sulfur Coated Urea.

LESCO, Pre-M and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Inc. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. Lely is a registered trademark of C. Van Der Lely N.V. (100209) Rev. 2/26/20 VT

Net Weight: XX lb (XX kg)

E.P.A. Reg No. 10404-98

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H)

F1560

Manufactured for: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

OPTIONAL MARKETING CLAIMS

Contains LESCO® Poly Plus® OPTI™ Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

Contains LESCO® NOS™ (Nitrogen Optimizing System) for increased nitrogen use efficiency and extended greening.

Contains LESCO® NOS™ Plus (Nitrogen Optimizing System) for increased nitrogen use efficiency and extended greening.

Contains LESCO® SulfRX™ Polymer Coated Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.