



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

10404-113

Date of Issuance:

03/04/2015

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

LESCO® Stonewall® 0.29% Plus  
 Fertilizer

Name and Address of Registrant (include ZIP Code):

George Meindl  
 LESCO Inc.  
 1385 East 36<sup>th</sup> Street  
 Cleveland, OH 44114-4114

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kathryn V. Montague, Product Manager 23  
 Herbicide Branch, Registration Division (7505P)

Date:

03/04/2015

2. You are required to comply with the data requirements described in the DCI Order identified as: Prodiamine GDCI-110201-1153.

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: [http://www.epa.gov/oppsrrd1/contacts\\_prd.htm](http://www.epa.gov/oppsrrd1/contacts_prd.htm).


3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 10404-113."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following Confidential Statements of Formula (CSFs): Basic, Alternate #1, Alternate #2, Alternate #3 and Alternate #4 CSFs dated 11/04/2014.

If you have any questions, please contact Terri Stowe at (703) 305-6117 or [stowe.terri@epa.gov](mailto:stowe.terri@epa.gov).

Sincerely,



Kathryn V. Montague, Product Manager 23  
Herbicides Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

# LESCO® Stonewall® 0.29% Plus Fertilizer

For selective preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns and landscape ornamentals (including established perennials and wildflower plantings).

**ACTIVE INGREDIENT:**

Prodiamine: (N3, N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine ..... 0.29%

**INERT INGREDIENTS:** ..... 99.71%  
Total ..... 100.00%

Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See the following sections: Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID	
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advise.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or when going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment advice.	

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

**STORAGE AND DISPOSAL**

**STORAGE:** Store this product in its original container in a dry, cool, secured area.

**Do not contaminate water, foodstuffs, feed or seed by storage or disposal.**

**PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

**CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns;
- Landscape ornamentals;
- Established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall, irrigation, or shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

**Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses, or lawns and grounds.**

- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system.
- Do not apply aerially.
- Do not apply to golf course putting greens.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**USE PRECAUTIONS**

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly over-seeded grasses. (2) Do not apply to new or overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. (4) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (5) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.


**APPLICATION DIRECTIONS**

Apply uniformly with suitable, calibrated application equipment.

**ESTABLISHED TURF**


This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

**CRABGRASS SEED GERMINATION DATES**



**RATES OF APPLICATION**

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.



**MAXIMUM ANNUAL RATES**

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate per Calendar Year of Product by Turf Grass Species <sup>1</sup>			
Turf Species	Lbs LESCO Stonewall 0.29% Plus Fertilizer / Acre	Lbs. product/1000 sq. ft.	lbs AI/Acre
Bermudagrass <sup>2</sup> Bahigrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	517	11.9	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	345	7.9	1.0
Creeping Red Fescue	259	5.9	0.75
Creeping Bentgrass	224	5.1	0.65

- These are the maximum rates per calendar year by species limitations.
- May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb a.i./acre (equal to 172 lbs./A of this product) [or] equal to 4.0 lbs./1000 sq.ft. of this product). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.
  - Do not apply more than 1.5 lbs a.i. per calendar year per acre (equal to 517 lbs./A of this product) [or] equal to 11.9 lbs./1000-sq.ft. of this product).
  - Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates** table.



**GUARANTEED ANALYSIS**

Total Nitrogen (N) XX.XX%  
 XX.XX% Ammoniacal Nitrogen  
 XX.XX% Urea Nitrogen  
 XX.XX% Water Insoluble Nitrogen  
 Available Phosphate (P<sub>2</sub>O<sub>5</sub>) XX.XX%  
 Soluble Potash (K<sub>2</sub>O) XX.XX%  
 Magnesium (Mg) Total XX.XX%  
 XX.XX% Water Soluble Magnesium (Mg)  
 Sulfur (S) Total XX.XX%  
 XX.XX% Free Sulfur (S)  
 XX.XX% Combined Sulfur (S)  
 Copper (Cu) Total XX.XX%  
 XX.XX% Water Soluble Copper (Cu)  
 Iron (Fe) Total XX.XX%  
 XX.XX% Water Soluble Iron (Fe)  
 Manganese (Mn) Total XX.XX%  
 XX.XX% Water Soluble Manganese (Mn)  
 Zinc (Zn) Total XX.XX%  
 XX.XX% Water Soluble Zinc (Zn)

**DERIVED FROM:**

\*X.XX% Slowly available Urea Nitrogen from Polymer Coated Urea.

**SPREADER SETTINGS**

Spreader settings vary by make and model of spreader. It is recommended that individual spreaders are calibrated for the specific product that is to be applied. A walking speed of 3 miles per hour is recommended.

Spreader	Application Rates <sup>1</sup> a.i. lbs./A (Product lbs./1000 sq ft)			
	0.5 (4.0)	0.75 (5.9)	1.0 (7.9)	1.5 (11.9)
LESCO – All Rotary Models	#XX	#XX	#XX	#XX
Permagreen	XX	XX	XX	XX
Cyclone <sup>®</sup> Spyker <sup>®</sup> Z-Spray	XX	XX	XX	XX
Vicon (LESCO Pendulum)	XX	XX	XX	XX

<sup>1</sup> Application Rates for example only - do not exceed the maximum annual application rate of 1.5 lb a.i./acre. (11.9 lb product/1,000 sq. ft.)

**LANDSCAPE ORNAMENTALS**

**Including Established Perennials and Wildflower Plantings**

This product may be applied for residual preemergence weed control in ornamentals.

**Use Rates and When to Apply**

Apply in fall and/or spring, 0.65 to 1.5 lbs active ingredient per acre, which is equal to 5.1 to 11.9 lbs. of this product per 1,000 sq. ft. of ornamental plantings; therefore 20 lbs of this product will treat 1,681 to 3,922 sq.ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs a.i. per acre.

**ESTABLISHED LANDSCAPE ORNAMENTALS**

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

**Newly Transplanted Landscape Ornamentals**

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

**Tolerant Ornamental Species**

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name	Scientific Name	Common Name
Abies spp.	Fir	Osleospermum fruticosum	Trailing African Daisy
Abelia grandifolia	Abelia; Sherwood	Oxydendron luteum	Sourwood
Acer palmatum	Japanese Maple	Persea Americana*	Avocado*
A. platanoides	Norway Maple	Photinia fraseri	Frasier's Photinia; Redtip
Achillea spp.	Yarrow (King Edward)	Physotegia Virginiana	False dragonhead (Vivid)
Actinidia chinensis*	Kiwi*	Picea spp.**	Spruce species** (Colorado Blue, Norway, etc.)
Agapanthus africanus	Lily-of-the-Nile; African Lily	Pieris japonica	Lily-of-the-Valley Shrub
Akebia quintata	Five-leaf or Chocolate vine	Pinus brutia	Calabrian Pine
Allium cernuum	Lady's leek or Nodding onion	Pinus canariensis	Canary Island Pine
Anemone hybrida	Japanese anemone	Pinus elliotii	Slash Pine
Aquilegia spp.	Columbine (Red and gold)	Pinus halepensis	Aleppo Pine
Arctostaphylos densiflora	Vine Hill manzanita	Pinus nigra	Austrian Pine
Arctotheca calendula	Cape Weed	Pinus radiata	Monterey Pine
Artemisia spp.	Wormwood (Silver mound; Castle)	Pinus palustris	Longleaf Pine
Aster sp. or Symphitrichium sp.	Aster (Bonny blue, Purple dome)	Pinus strobus	Eastern White Pine
Aster X frikartii		Pinus sylvestris	Scotch Pine
<i>Athyrium felix-femina</i>	Lady fern	Pinus taeda	Loblolly Pine
Aucuba japonica	Japanese Aucuba	Pinus thunbergiana	Japanese Black Pine
Begonia spp.	Fibrous bigonia (Hardy Grandis)	Pinus virginiana	Virginia pine
Berberis gladywensis	Barberry	Pistachio sp.*	Pistachio*
Berberis julianae	Wintergreen Barberry	Pittosporum rhombifolium	Queensland Pittosporum
Berberis mentorensis	Mentor barberry	Pittosporum tobira	Japanese Pittosporum
Berberis thunbergii	Japanese Barberry	Podocarpus macrophyllus	Japanese Yew
Berberis verruculosa	Warty Barberry	Prunus laurocerasus	English Laurel
Bergenia cordifolia		Prunus spp.*	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
Boltonia asteroides	Snowbank	Pseudotsuga menziesii**	Douglas Fir**
Bougainvillea spp.	Bougainvillea	Pyracantha coccinea	Firethorn, Scarlet
Buddie ia davidii	Butterfly-bush (Dwarf Blue, Royal Red)	Pyracantha fortuneana	Firethorn

Buxus microphylla	Japanese Boxwood	Pyracantha koidzumii	Firethorn
Callistemon viminalis	Weeping Bottlebrush	Pyrus spp.	Bradford Pear
Calluna vulgaris	Scotch Heather	Quercus rubra	Red Oak
Campanula carpatica	Tussock bellflower (White Clips)	Quercus shumardii	Shumard oak
Campsis X tagliabuana	Trumpet creeper; Trumpet flower (Madame Gaten)	Raphiolepis indica	Indian Hawthorne
Carpobrotus edulis	Hottentot Fig; Ice Plant	Rhododendron (including Azalea)	Rhododendron & Azalea (Coral Bells, Delaware valley white, Flame creeper, George L. Tabor, PJM, Roseum Elegans, Formosa, Hino-crimson, Wakaebisu, Gumpo)
Cassia artemisoides	Feathery Cassia	Rosa banksiae	Lady Bank's Rose
Ceanothus rigidus	Wild Lilac	Rosmarinus officinalis*	Rosemary*
Ceratostigma plumbaginoides		Rudbeckia, spp.	Black-eyed Susan
Chamaecyparis pisifera	False Cypress	Rumohra adiantiformis	Leatherleaf Fern
Chrysanthemum nipponicum		Santolina wrens	Goldstrum Santolina
Cleyera japonica	Cleyera	Saxifraga spp.	Saxifrage, Purple dome
Citrus spp. *	Citrus species*	Sedum album	Stonecrop
Coreopsis spp.	Coreopsis; calliopsis (Early Sunrise, Moonbeam)	Sedum caulicola	Stonecrop, likadense
Cornus florida	Flowering Dogwood	Sedum dasyphyllum	Stonecrop
Cornus stolonifera	American Dogwood	Syzygium paniculatum	Japanese Boxcherry
Cortaderia selloana	Pampas Grass	Taxus cuspidata	Japanese Yew
Cotoneaster apiculatus	Cranberry Cotoneaster	Taxus media	Yew
Cotoneaster buxifolius	Cotoneaster	Teucrium spp.	Germander
Cotoneaster dammeri	Bearberry Cotoneaster	Thalictrum dipterocarpum	Meadow rue
Cotoneaster microphyllus	Rockspray Cotoneaster	Thuja occidentalis	American Arborvitae
Cretaeagus spp.	Hawthorne	Trachelospermum asiaticum	Star Jasmine
Crocsmia spp.	Lucifer	Tsuga canadensis	Canada Hemlock
Cupressus sempervirens	Italian Cypress	Tulipa spp.	Tulip
Delasperma alba	White Trailing Ice Plant	Viburnum japonicum	Japanese Viburnum
Delphinium spp.	Larkspur (Blue elf)	Viburnum odoratissimum	Sweet Viburnum
Dianthus deltooides	Maiden pinks (Zing)	Viburnum plicatum	Japanese Snowball
Dianthus gratianopolitanus	Cheddar pinks	Viburnum rigidum	Canary Island Viburnum
Dodonea viscosa	Hop Bush	Viburnum tinus	Laurustinus
Echinacea purpurea	Coneflower, purple (Magnus)	Viburnum trilobium	Cranberry Bush
Elaeagnus pungens	Silverberry	Viburnum wrightii	Leatherleaf Viburnum
Euonymus fortunei	Wintercreeper	Vinca major	Greater periwinkle, Vinca
Euonymus japonica	Japanese spindle tree; Evergreen Euonymus	Vinca minor	Common Periwinkle
Euonymus kiautschovica	Spreading Euonymus	Vitis spp.*	Grape*
Fatsia japonica	Japanese Aralia	Weigela florida	Old Fashioned Weigela
Forsythia intermedia	Border Forsythia	Yucca aloifolia	Spanish Bayonet
Forsythia suspense	Weeping forsythia	Yucca filamentosa	Yucca, Adam's Needle
Forsythia viridissima	Greenstem Forsythia		
Gail lard ia spp.	Blanket flower (Goblin)		
Gardenia jasminoides	Gardenia, Cape-Jasmine		
Gentiana dahurica	Gentian		
Geranium cinereum	Cranesbill		
Gladiolus spp.**	Gladiolus		
Gypsophila repens	Baby's breath		
Hedera helix	English Ivy		
Helianthemum spp.	Sunrose		
Hemerocallis spp.	Daylily (Aztec gold, Stella d'Oro, Tender love)		
Hibiscus spp. **	Rose of Sharon**		
Hibiscus spp.	Mallow (Disco belle white)		
Hibiscus Rosa-sinensis**	Chinese hibiscus**		
Hosta plantaginea	Hosta, Plantain lily		
H. sieboldiana	Hosta (Searsucker)		
Houttuynia cordata var. variegata			
Hydrangea macrophylla	Bigleaf hydrangea		

Ilex cornuta**	Chinese holly**		
Ilex crenata	Japanese Holly		
Ilex opaca	American Holly		
Ilex pernyi	Holly		
Ilex vomitoria	Yaupon holly		
Inula ensifolia			
Iris ensata	Sword-leaved iris (Jodlesong)		
Iris siberica	Siberian iris (Cabernet)		
Iris spp.	Iris **		
Jasminum nudiflorum	Winter Jasmine		
Juniperus chinensis	Chinese Juniper (Nick's Compact)		
Juniperus conferta	Shore Juniper (Blue Pacific)		
Juniperus davurica	Juniper		
Juniperus horizontalis	Creeping Juniper		
Juglans sp.*	Walnut*		
Justica brandegeana	Shrimp Plant		
Lagerstromia indica	Crape Myrtle		
Lantana montevidensis	Weeping lantana		
Lavender spp.	Lavender (Munstead)		
Leontopodium alpinum	Edelweiss		
Ligustrum amurense	Amur Privet		
Ligustrum japonicum	Japanese Privet		
Ligustrum lucidum	Glossy Privet (Wax-Leaf)		
Ligustrum sinense	Chinese privet (Variegata)		
Lilium spp.	Lily (Jazz)		
Liriope muscari	Liriope (Big Blue, Lily Turf, Silver Mound, Evergreen Giant)		
Liriope muscari var. variegata	Liriope, variegated		
Liriope spicata	Liriope, creeping		
Lobelia cardinalis	Cardinal flower		
Lonicera japonica	Japanese Honeysuckle		
Lonicera tatarica	Tatarian Honeysuckle		
Loropetalum chinense	Burgundy		
Ly thrum spp.	Loosestrife (Modern Pink)		
Magnolia spp.**	Magnolia**		
Maleophora luteola	Ice Plant		
Malus spp.*	Crabapple*		
Miscanthus sinensis	Yaku jima, Silberfeder		
Nandina domestica	Heavenly Bamboo		
Narcissus spp.**	Narcissus**		
Nerium Oleander*	Oleander		
Olea europaea*	Olive*		
Ophiopogon japonicus**	Mondo Grass		
Osmanthus heterophyllus	Osmanthus; False holly (Gulf tide)		

\* Do not use on food producing trees, vines or plants.

\*\*Not for use on container grown plants.

### WEEDS CONTROLLED

When used in accordance with this label, this product will provide control of the following weeds:

Barnyardgrass	Goosegrass <sup>1</sup>	Lovegrass	Speedwell, Persian
Bluegrass, Annual (Poa annua)	Henbit	Panicum (Texas, Fall, Brown Top)	Sprangletop
Carpetweed	Itchgrass	Pigweed	Spurge, Prostrate
Chickweed, Common	Johnsongrass (from seed)	Purslane, Common	Witchgrass
Chickweed, Mouse ear (from seed)			
Crabgrass (large, smooth)	Junglerice	Pusley, Florida	Woodsorrel, Yellow (from seed)
Crowfoot grass	Knotweed	Rescuegrass <sup>3</sup>	
Cupgrass, Woolly	Kochia	Shepherd's Purse <sup>2</sup>	
Foxtails, Annual	Lambsquarter, Common	Signalgrass, Broadleaf	

1. In many areas a single application of 0.65 lb. to 1.5 lb. active ingredient per acre (equal to 224 lbs. to 517 lbs./A of this product) [or] equal to 5.1 to 11.9 lbs./1000 sq. ft. of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to 224 lbs. to 345 lbs./A of this product) [or] equal to 5.1 to 7.9 lbs./1000 sq. ft. of this product) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.

2. Applications for this weed should be made in late summer, fall or winter prior to germination.

3. Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

**WHEN TO APPLY AFTER OVERSEEDING TURF**

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

**WHEN TO OVERSEED AFTER APPLICATION**

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

LESCO Stonewall 0.29% Plus Fertilizer Rate (lb/A)	Lbs product/1000 sq. ft.	Lbs a.i./A	Months Before Overseeding		
			North	Transition	South
172	4.0	.50	4	4	4
224	5.1	.65	5	4	4
259	5.9	.75	6	5	5
276	6.3	.80	-	6	6
345	7.9	1.00	-	7	7
393	9.0	1.14	-	-	9
448	10.3	1.30	-	-	10
517	11.9	1.50	-	-	12

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>.

LESCO and Stonewall and the sweeping design are trademarks of LESCO Technologies, LLC. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.

G:\REGUL\WP\Regul Private\MstrLbl\ERIEVIEW ADDRESS\New EPA Approved\Stonewall 0.29%.doc (050914) Rev.: 10/16/14 VT

**Net Weight: XX lb (XX kg)**

**EPA REG. NO. 10404-RRG**

EPA EST.NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E)

First letter of lot code indicates manufacturing site.

**#XXXXXX**

**F1560**

Distributed by: LESCO, Inc.  
1385 East 36<sup>th</sup> Street  
Cleveland, OH 44114-4114

### **OPTIONAL MARKETING CLAIMS**

- o Feeds [the lawn] up to (2, 3, 4, 5) months
- o Contains [multi source] [multi release rate] nitrogen [for improved root growth]
- o Improved nitrogen consumption
- o Reduced nitrogen overload
- o Reduced nitrogen runoff
- o Facilitates reduced irrigation
- o Improved drought tolerance
- o Equivalent [or improved] efficacy at lower NPK levels
- o Controlled release feeding for uniform growth
- o [Provides] [season-long] control of crabgrass
- o [Provides] long-lasting green
- o Provides slow release nutrients to the soil
- o Creates [stronger] [deeper] root system
- o Feeds [your lawns] roots
- o Apply at the beginning of the growing season [to enhance nutrition and enhance crabgrass control]
- o ([Company name] supports the EPA Greenscapes approach to sustainable lawn care landscaping
- o Combines conventional [*synthetic and/or inorganic*] and natural [organic] fertilizers
- o Reliable [long-lasting] green-up
- o 3 month green-up
- o Easy to apply
- o Delivers quick results [without surge growth]
- o Aids in nutrition and disease suppression
- o Contains (XX % ) organic matter (OM)
- o Combines quick results and long-lasting feeding [as nature intended]
- o Adding the right balance of nutrient sources helps create a healthier lawn and soil
- o Using slowly available nutrients greatly reduces run-off into our waterways

### **[Optional state specific language: Florida]**

In the state of Florida, it is recommended that the user of this product follow the Green Industries and Golf Course Best Management Practices found at:  
[http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP\\_Book\\_Final.pdf](http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_Final.pdf) and  
<http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/gfbmp07.pdf>.

### **[Optional state specific language: Virginia (eff. 12-31-13)]**

In Virginia this lawn/turf fertilizer contains phosphorous and is intended only for nonagricultural use on (1) turf during its first growing season, (2) on turf areas being repaired or renovated, and (3) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not intended for the routine maintenance of turf.

### **[Optional state specific language: Maryland (eff. 10-01-13)]**

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.