

DEODORANT **Hy-Kresol** DISINFECTANT

CONTENTS

| | |
|-------------------------|------|
| ACTIVE INGREDIENTS: | |
| Cresylic Acid | 50% |
| Soap | 30% |
| INERT INGREDIENTS | 20% |
| | 100% |

ACCEPTED

27

UNDER THE DEPARTMENT OF AGRICULTURE
 FUNCTIONAL USE OF THE FEDERAL AGENCY
 FOR ECONOMIC DEVELOPMENT
 ED UNDER THE FEDERAL REGISTER

For Disinfection and Decontamination of Poultry Houses, Stock-yard Premises and Vehicles Used in Interstate Transportation of Livestock.

DANGER: Keep Out of Reach of Children. CORROSIVE. Causes eye damage and severe burns. May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Avoid inhaling fumes or spray mist. Avoid contamination of feed or foodstuffs.

FIRST AID: EXTERNAL: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, get prompt medical attention.

INTERNAL: If swallowed, give milk, white of raw eggs, mucilage, or corn starch paste, and follow with an emetic such as a tablespoonful of mustard in a tumbler of warm water.

NOTE TO PHYSICIAN: Measures against circulatory shock, respiratory depression and convulsion may be needed. Watch for signs of pulmonary edema, which may develop rapidly.

Rinse Empty Container Thoroughly with Water and Discard it.

670-46

GENERAL DISINFECTION OF POULTRY HOUSES

It is recommended that poultry barns be treated between each flock change as follows:

After poultry is removed from barn, wash thoroughly ceilings, walls, posts, rafters, air ducts, feeders and waterers with pressure water and Smith Poultry Barn Cleaner.

Remove litter. When surfaces are dry, apply HY-KRESOL solution (2 oz. per gallon water) and saturate all walls, ceilings, rafters, fans, ducts, posts, feeders and waterers being sure all areas have been reached. To disinfect hard to reach areas, retreatment is recommended.

All treated feeders and waterers must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

Allow to absorb or dry before birds enter. After treatment of surfaces: Ventilate buildings, vehicles, and other closed spaces.

When treating dirt floors with fuel oil, HY-KRESOL may be added at the rate of 2 oz. per gallon.

Apply disinfectant only at recommended rate.

E.P.A. Reg. No. 670-46

EST. 670-MN-1

1910
 University Ave.
 St. Paul, Minn.
 55104



Phone:
 646-6131


ACTUAL SIZE (4 1/4" x 5")

DIRECTIONS: FOR DISINFECTANT SOAP: Mix 1 cup Commercial Pine Oil to pail (4 gals.) of hot water, or 2 oz. to 1 gal. water. Scrub well & rinse.
~~FOR FLEAS: Mix 9 tablespoons per gallon. FOR FLEAS: 1 part to 3 parts water. Spray or scrub. FOR DEODORANT: Dilute 1-20 to 1-200 as desired.~~
Mfg by PROTEX WAX CO 235 48 AVENUE Oakland California

U.S.P.A. Reg. No.



NET CONTENTS
ONE GALLON



PHENOL
COEFFICIENT : 7

ACTIVE INGREDIENTS:
PINE OIL 80%
SOAP 17.3%
INERT INGREDIENTS 2.7%

12 point

ACCEPTED
MAR 3 - 1969

10319 1

10419-1


ACTUAL SIZE (4 1/4" x 5")



U.S. PAT. REG. OFF.

NET CONTENTS
ONE GALLON

Commercial



DISINFECTANT

PHENOL
COEFFICIENT: 7

ACTIVE INGREDIENTS
PINE OIL 80%
SOAP 17.5%
INERT INGREDIENTS 2.1%

DIRECTIONS: FOR DISINFECTANT SOAP: Mix 1 cup Commercial Pine Oil to pail (4 gals.) of hot water, or 2 oz. to 1 gal. water. Scrub well & rinse.
~~FOR FLEAS: Mix 3 tablespoons per gallon. FOR FLEAS: 1 part to 3 parts water. Spray or scrub.~~ FOR DEODORANT: Dilute 1-20 to 1-200 as desired.

12 point