

SANIKLEER

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 28 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10428-19

ACTIVE INGREDIENT:

Poly(oxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride) 30.0 %

INERT INGREDIENT 70.0 %

This product contains 2.67 lb. of active ingredient per gallon and weighs
8.9 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANIKLEER

_____ is used to control algae in swimming pools, exterior spas, whirlpools, hot tubs, and domestic hot water tank effectiveness pools, spas, whirlpools, hot tubs, and domestic hot water tanks. Heavy growth of algae should be cleaned prior to using SANIKLEER.

For pools having just visible algae growth add an initial dose of _____ fluid ounces of SANIKLEER _____ per 10,000 gallons of water. To move settled algae debris by cleaning. For treatment of a pool add initially 12.8 to 23.5 fluid ounces of SANIKLEER per 10,000 gallons of water. Subsequent additions of 4.3 fluid ounces of SANIKLEER per 10,000 gallons of water should be made every 7 days after initial treatment for maintenance.

Spas, whirlpools, hot tubs, or decorative fountains having just visible algae growth require an initial dose of 2 to 4 fluid ounces of SANIKLEER per 1000 gallons of water. For treatment of a freshly cleaned hot tub, or decorative fountain, add initially 1 fluid ounce of SANIKLEER per 1,000 gallons of water. Subsequent additions of 0.5 to 0.8 fluid ounces of SANIKLEER should be made every 5 to 7 days after initial treatment for maintenance. SANIKLEER is compatible with all other pool chemicals used to treat pools and spas and is effective at bactericidal levels. SANIKLEER can be used in place of liquid chlorine chemicals and may reduce the amount of those chemicals required. However, do not mix SANIKLEER with liquid chlorine products.

STORAGE AND DISPOSAL

STORAGE: Keep container closed when not in use. Containers should be placed in a plastic bag and discarded after use. Rinse with clear water. All pesticides should be kept out of reach of children and away from food.

DISPOSAL: Do not reuse empty container. Wrap container in plastic.

Manufactured by:

WESMAR CO., INC.

333 NE 89th

SEATTLE, WA 98107

EPA REG. NO. _____

EPA EST. NO. _____

NET CONTENTS _____