

SUPER-MOL FLY CONTROL BLOCKS  
MEDICATED

---

Prevents the breeding of Horn Flies and Face Flies in the manure of treated cattle: Aids in the prevention of Stomach Worms, Hookworms, and Nodular Worms when used as directed: Aids in the prevention of Foot Rot in cattle, soft tissue, lumpy jaw, Nutritional source of Iodine when fed to Cattle at the rate of 4 oz. per head per day.

---

FEEDING DIRECTIONS

Self Feed Super-Mol Fly Control Block, Medicated, to cattle as the sole source of salt in order that the following amounts will be consumed, based on animal body weight:

400 lbs. - 2 oz. per head daily  
600 lbs. - 3 oz. per head daily  
800 lbs. - 4 oz. per head daily  
1000 lbs. - 5 oz. per head daily

Each 2 oz. supplies 1/2 gram of phenothiazine which is the daily dosage for each 100 lbs. of body weight for preventing the breeding of horn and face flies in the manure of treated cattle. The same level aids in the prevention of stomach worms, hookworms, and Nodular worms in cattle.

Offer free choice to sheep and goats in order that each animal will receive 2 oz. per head daily to supply one (1) gram of phenothiazine as an aid in the prevention of stomach worms, hookworms, and Nodular worms. DO NOT FEED ADDITIONAL SALT.

Blocks should be fed near the animals water supply or near their loafing area. In order to assure proper horn and face fly breeding control, animals should have these blocks available at all times. Offer these blocks to cattle from early spring until the fly season is past. Phenothiazine is not effective against the adult fly. Therefore, it is suggested that the animals be sprayed once in early summer to kill the adult flies present. Then follow with the SUPER-MOL FLY CONTROL BLOCK program to control the re-population of Horn and Face Flies. Since flies migrate from farm to farm, this program will be most effective when carried out on an area wide basis. These blocks should be fed as directed on a year around basis for internal parasite control.

CAUTION: KEEP OUT OF REACH OF CHILDREN

USDA REG. NO.

MANUFACTURED BY  
SUPER-MOL DIVISION  
NATIONAL MOLASSES COMPANY  
TAMPA, FLORIDA 33609

7-55

NET WEIGHT 33 11/32 LBS.

SUPER-MOL FLY CONTROL BLOCKS  
MEDICATED

PESTICIDALLY ACTIVE DRUG INGREDIENT:	
Phenothiazine	2.00%
PESTICIDALLY INERT INGREDIENTS:	98.00%
ACTIVE DRUG INGREDIENT:	
Ethylene Diamine Dihydr iodide	.044%

GUARANTEED ANALYSIS

Crude Protein	Not less than	4.00%
Crude Fat	Not less than	1.00%
Crude Fibre	Not more than	10.50%
Calcium (CA)	Not less than	3.00%
Calcium (CA)	Not more than	4.00%
Phosphorus (P)	Not less than	1.00%
Iodine (I)	Not less than	0.015%
Cobalt (CO)	Not less than	0.010%
Salt (NaCl)	Not less than	19.00%
Salt (NaCl)	Not more than	22.00%
Total Sugars (as invert)	Not less than	23.00%
Vitamin A	Not less than USP Units/LB.	50,000
Vitamin D <sub>3</sub>	Not less than USP Units/LB.	15,000
Total Added Mineral Ingredients	Not more than	52.00%

INGREDIENTS: Dried Cane Molasses, Cane Molasses, Salt, Calcium Carbonate, Cane Bagasse, Alfalfa Meal, Defluorinated Phosphate, Soybean Meal, Cottonseed Meal, Meat and Bone Meal, Vitamin A Palmitate, D-Activated Animal Sterol (source of Vitamin D<sub>3</sub>) Iron Oxide, Iron Carbonate, Iron Sulphate, Magnesium Oxide, Zinc Oxide, Manganous Oxide, Copper Oxide, Cobalt Carbonate, Natural and Artificial Flavors added.

WARNING: DO NOT FEED TO LACTATING DAIRY ANIMALS. Occasional sensitivity to Phenothiazine has been reported in Cattle.

KEEP OUT OF REACH OF CHILDREN

USDA REG. NO.

MANUFACTURED BY  
SUPER-MOL DIVISION  
NATIONAL MOLASSES COMPANY  
TAMPA, FLORIDA 33619

January 13, 1969