

Super-Mol

Fly & Worm Control Blocks - M MEDICATED

PESTICIDALLY ACTIVE DRUG INGREDIENT 2.00 %
Phenothiazine

PESTICIDALLY INERT INGREDIENTS 98.00 %

ACTIVE DRUG INGREDIENT .044
Ethylene Diamine Dihydrochloride

GUARANTEED ANALYSIS

Crude Protein (not less than)	4.00 %
Crude Fat (not less than)	1.00 %
Crude Fiber (not more than)	10.50 %
Calcium (Ca) (not less than)	1.00 %
Calcium (Ca) (not more than)	4.00 %
Phosphorus (P) (not less than)	1.00 %
Iodine (I) (not less than)	1035 %
Cobalt (Co) (not less than)	1010 %
Salt (NaCl) (not less than)	19.00 %
Salt (NaCl) (not more than)	22.00 %
Magnesium	1.00 %
Vitamin A (not less than USP Unit per lb.)	1000
Vitamin D (not less than USP Unit per lb.)	1000
Total Added Mineral Ingredients (not more than)	12.00 %

INGREDIENTS: Dried Cane Molasses, Tank Molasses, Calcium Carbonate, Alfalfa Meal, Cottonseed Meal, Prostate, Soybean Meal, Cottonseed Meal, Meal, Molasses, Vitamin A Palmitate, Activated Animal Protein, Calcium Hydroxide, Magnesium Hydroxide, Zinc Oxide, Magnesium Oxide, Sodium Chloride, Dicalcium Phosphate, Anhydrous Ferric Sulfate.

WARNING —

Do not treat ruminants with this product if they are ill.

FOR RUMINANTS ONLY

33-11/32 LBS. NET WEIGHT

Three Blocks Equal

100 lbs. each

Prevents the breeding of Horn Flies, and Face Flies on the heads of treated cattle. Also controls the following: Stable Fly, Worms, Hairworms, Pinworms, Oestertad, Trichostrongylus, Axed, Hookworms, (Roundworms), pinworms, and Nematode Worms. (Oesophageal stomach parasites when self-medicated). Aids in the prevention of Fogg Rot in cattle with loose

limp jaw. Nutritional supplement when fed at least at the rate of 4 oz. per head per day.

FEEDING DIRECTIONS

Self Feed Fly and Worm Control Blocks M Medicated to cattle as described above at a rate of 4 oz. per head per day. The following amount will be fed per head based on animal body weight:

400 lbs.	2 oz. per head daily
600 lbs.	3 oz. per head daily
800 lbs.	4 oz. per head daily
1000 lbs.	5 oz. per head daily

Each 1/2 oz. Supplies 1/2 cent of phenothiazine which is the daily dosage for 100 lbs. of body weight for preventing the breeding of horn and face flies in the manure of treated cattle. The same level aids in the prevention of stomach worms, hairworms, pinworms, Oestertad, Trichostrongylus, Axed, Hookworms, (Roundworms), pinworms, and Nematode Worms (parasitic stomach parasites) in cattle.

DO NOT FEED TO NON-RUMINANTS

Blocks M will be fed to ruminants with a supply of water. Do not feed to birds, swine, dogs, cats, horses, and other non-ruminants. The following directions have been established for the use of Super M Fly and Worm Control Blocks M with the Fly and Worm Control Block M. It is suggested that the product be stored in a cool, dry place. The product is present in the law with the Fly and Worm Control Block M program. The product is present in the law with the Fly and Worm Control Block M program. The product is present in the law with the Fly and Worm Control Block M program. The product is present in the law with the Fly and Worm Control Block M program.

CAUTION: KEEP OUT OF REACH OF CHILDREN

1974

TAMPA, FLORIDA 33604

Manufactured by

Super-Mol Division

TAMPA, FLORIDA 33604

ACCEPTED WITH COMMENTS