



**ACTIVE INGREDIENTS**  
 Phenothiazine (Biologically Assayed) \_\_\_\_\_ 1.000%  
 OTHER INGREDIENTS\* \_\_\_\_\_ 98.700%  
 Total \_\_\_\_\_ 100.000%

\*Active only to ingredients which are neither insecticides nor drugs.

**OTHER ACTIVE INGREDIENTS**  
 Ethylene Diamine Hydrochloride \_\_\_\_\_ 0.000%

**INGREDIENTS**  
 Cottonseed meal, dehydrated alfalfa meal, yeast and bone meal, corn meal, eggs, milk, zinc acetate, dried molasses, dicalcium phosphate, magnesium carbonate, calcium chloride, calcium sulfate, calcium carbonate, zinc sulfate, magnesium sulfate, magnesium oxide, silicon hexachloride, sulfur, ammonium phosphate, sodium phosphate, dried yeast, thymol, sodium acetate, yeast culture, Vitamin B12 supplement, zinc, sodium chloride, calcium phosphate, riboflavin supplement, zinc oil, inactive dried yeast (source of Vitamin B2), and Vitamin A palmitate.

Manufactured & Guaranteed by V. M. S., INC.  
 Montgomery, Alabama 36101

**GUARANTEED ANALYSIS:**

Crude Protein, not less than \_\_\_\_\_ 18.00%  
 (This includes not more than 0.00% equivalent protein from non-protein nitrogen)  
 Crude Fat, not less than \_\_\_\_\_ 1.00%  
 Crude Fiber, not more than \_\_\_\_\_ 8.00%  
 Calcium (Ca), not more than \_\_\_\_\_ 8.50%  
 Calcium (Ca), not less than \_\_\_\_\_ 4.00%  
 Phosphorus (P), not less than \_\_\_\_\_ 1.50%  
 Iodine (I), not less than \_\_\_\_\_ 0.025%  
 Salt (NaCl), not more than \_\_\_\_\_ 21.00%  
 Salt (NaCl), not less than \_\_\_\_\_ 10.00%  
 Magnesium (Mg), not less than \_\_\_\_\_ 1.25%  
 Manganese (Mn), not less than \_\_\_\_\_ 0.012%  
 Cobalt (Co), not less than \_\_\_\_\_ 0.002%  
 Ash, not more than \_\_\_\_\_ 40.00%  
 Vitamin B2, not less than 10,000 U.S.P. Units per lb.  
 Vitamin A, not less than 40,000 U.S.P. Units per lb.

**FEEDING DIRECTIONS**

Self feed continuously as the sole source of salt so as to provide 3/4 ounce per 100 pounds of body weight per head daily. Do not offer any other mineral or salt mixtures. The above daily intake levels will furnish the proper medication to support drug claim. **CAUTION:** Do not feed this medicated block to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle.

**CAUTION!**

**KEEP OUT OF REACH OF CHILDREN**

Note: For maximum protection against horn flies and face flies, it is recommended that cattle be sprayed once when first starting on a continuous KATTLE-KING Bar-Fly Program. KATTLE-KING Bar-Fly prevents the breeding of horn flies and face flies in the manure, but does not affect any existing adult fly problem. Be sure and supply ample blocks at all times—at least one block for every 10 to 12 head. Since face flies migrate from farm to farm, it is essential that the larvicide program be carried out on an area basis to assist in the reduction of the adult face fly population.