

345-210546
17

SEP - 6 1988

Lee Johnson, Ph.D.
V.M.S., Inc.
1080 Wilbanks Street
Montgomery, AL 36195

Dear Dr. Johnson:

Subject: Amendment - Change in Inert Ingredients
and Revised Labeling
V.M.S. Pest-A-Side Block W/Rabon
EPA Registration No. 10461-11
Your Application Dated December 11, 1987
Your Letter Dated March 17, 1988

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

1. Change ACTIVE INGREDIENTS to the singular.
2. Change INGREDIENTS to FEED INGREDIENTS. See our letter of October 19, 1984, paragraph 2.b.
3. ~~Delete "AND GUARANTEED."~~
4. Delete "for 15 minutes" in the If In Eyes section.
5. ~~Delete "SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS."~~
6. Delete "GENERAL CLASSIFICATION."
7. Revise the Storage and Disposal block. See our letter of October 29, 1984, paragraph 2.c.

YJ

05591:I:Newkirk:LR-7:KENCO:6/14/88:6/23/88:rw:vo:ek:rw
R:58653:Newkirk:LR-7:KENCO:6/29/88:8/11/88:rw

CONCURRENCES

SYMBOL	SURNAME	DATE					

Please provide the following product chemistry information.

- 1. Provide the upper and lower certified limits based on the pure active ingredients rather than the technical or concentrate. Note that the lower limit of the active must be the same as the label claim.
- 2. Based on the current Confidential Statement of Formula (CSF) dated December 11, 1987, your product will not meet the label claim for the active ingredient. Please revise the label or the CSF so the information agrees.
- 3. ~~Provide the EPA Registration Number for the source of the active ingredient.~~

~~The printed labels submitted with your letter of March 17, 1988 are not acceptable and are returned without comment or accepted stamp.~~

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
 Product Manager (15)
 Insecticide-Rodenticide Branch
 Registration Division (TS-767C)

Enclosures



PEST-A-SIDE BLOCK WITH RABON®



To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)	0.30%
diethyl dimethyl phosphite	32.70%
TOTAL	33.00%

*RABON® is a trade name - Contains 1.5% of insecticide per pound.
**Data only in liquid form which is not applicable.

EPA REG. NO. 10461-11
EPA EST. NO. 10461-AI-1
RABON® is a registered trademark of Shell Oil Company

GUARANTEED ANALYSIS:

Crude Protein, minimum	8.00%
Crude Fat, minimum	0.50%
Crude Fiber, maximum	7.50%
Calcium (Ca), minimum	3.50%
Calcium (Ca), maximum	7.50%
Phosphorus (P), minimum	2.00%
Salt (NaCl), minimum	12.00%
Salt (NaCl), maximum	18.00%
Magnesium (Mg), minimum	1.00%
Sulfur (S), minimum	0.15%
Cobalt (Co), minimum	0.0017%
Copper (Cu), minimum	0.0007%
Iodine (I), minimum	0.0007%
Manganese (Mn), minimum	0.00%
Selenium (Se)	0.0020%
Zinc (Zn), minimum	0.047%
Vitamin A, minimum	20,000 I.U. Per lb.
Vitamin D-3, minimum	5,000 LU Per lb.

INGREDIENTS:

Cottonseed meal, dehydrated alfalfa meal, dicalcium phosphate, monocalcium phosphate, salt, calcium carbonate, magnesium limestone, magnesium silca, magnesium sulfate, potassium sulfate, cobalt carbonate, cupric oxide, calcium iodide, ferrous carbonate, manganese oxide, sodium selenite, zinc oxide, vitamin A supplement, D-activated animal sternal (source of vitamin D-3) yeast culture, nicotinic choline chloride, riboflavin supplement, calcium borogluconate, bacillus thuringiensis extract and case molasses.

MANUFACTURED AND GUARANTEED BY
V.M.S., INC.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 06 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

10461-11

**SEE BELOW FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

Keep out of reach of children. Handle if contained. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If inhaled, breathe, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to any body of water. Do not contaminate water by cleaning of equipment or disposal of tanks.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

DIRECTIONS FOR USE:

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 20-25 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg. BAMBON® and Ivermectin per hundred pounds of body weight. Animals should not be starved for animals before initiation of the U.S.S. Post-A-Side Block with BAMBON® program but OBTAIN all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form.

Place near watering and feeding areas. Protect from rain.

If consumption of U.S.S. Post-A-Side Block with BAMBON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or rations. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding U.S.S. Post-A-Side Block with BAMBON® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall.

U.S.S. Post-A-Side Block with BAMBON® may be fed to slaughter and to lactating dairy cows.

U.S.S. Post-A-Side Block with BAMBON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplement fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or slugs on the premises.

In order to achieve optimum fly control, U.S.S. Post-A-Side Block with BAMBON® should be used in conjunction with other good management and sanitation practices.

CAUTION

Follow feeding directions. When this product is consumed as directed, it will provide 0.3 PPM of Selenium (Se).

For specific recommendations on a complete supplement program, contact U.S.S., Inc., Department of Nutrition.

(83-488)

8/68

TAKE TIME



**OBSERVE LABEL
DIRECTIONS**