

(BEST AVAILABLE COPY)

50 LBS. NET WEIGHT




MADE IN MONTGOMERY, ALABAMA
MANUFACTURED BY
V.M.S. INC.

ENPROAL[®] FLY CONTROL SUPPLEMENT WITH RABON[®] ORAL LARVICIDE

TAKE CARE

OBSERVE LABEL
DIRECTIONS



To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT By Weight

2 Chloro-1,1,4
5 trichloroethoxyethyl dimethyl phosphate Minimum—0.33%* Maximum—0.33%
INERT INGREDIENTS Minimum—99.7%** Maximum—99.67%

TOTAL 100.00%

* RABON[®] Insecticide—Contains 1.36 g RABON[®] per pound. RABON[®] is the registered trademark of Shell Oil Company.
** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS:

Crude Protein, minimum	7.50%
Crude Fat, minimum	0.20%
Crude Fiber, maximum	3.00%
Calcium (Ca), maximum	2.75%
Calcium (Ca), minimum	1.75%
Phosphorus (P), minimum	1.00%
Salt (NaCl), maximum	6.00%
Salt (NaCl), minimum	5.00%
Magnesium (Mg), minimum	3.00%
Potassium (K), minimum	2.40%
Sulfur (S), minimum	0.70%
Cobalt (Co), minimum	0.001%
Copper (Cu), minimum	0.03%
Iodine (I), minimum	0.005%
Iron (Fe), minimum	0.1%
Manganese (Mn), minimum	0.12%
Selenium (Se), minimum	0.00028%
Zinc (Zn), minimum	0.21%
Fluorine (F), maximum	0.01%
Vitamin A, minimum	52,000 U.S.P. units per lb.
Vitamin D-3, minimum	10,000 U.S.P. units per lb.
Vitamin E, minimum	25 I.U. per lb.
Total Sugars (as invert)	30.00%

INGREDIENTS:

Cane molasses, cottonseed meal, dried whey, magnesium oxide, salt, defluorinated phosphate, magnesium sulfate, potassium sulfate, zinc sulfate, zinc oxide, iron sulfate, iron carbonate, manganese sulfate, manganese oxide, copper sulfate, sodium selenite, calcium carbonate, calcium iodate, cobalt sulfate, cobalt carbonate, vitamin A supplement, D-activated animal stanol (source of Vitamin D-3) and vitamin E supplement.

10-15-86

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARD TO HUMANS

Caution: This product is toxic to humans. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling. Do not eat, drink, or smoke while using this product.

ENVIRONMENTAL PRECAUTIONS:
This product is toxic to fish and aquatic life. Do not contaminate water bodies or streams with wastes.

STORAGE AND DISPOSAL:
Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store in a dry place in original container.

HAZARDOUS DISPOSAL: Wastes resulting from the use of this product may be disposed of on land or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container into a hole or a hole in the ground. Do not reuse. Do not burn, bury, or dispose of in any other manner. If burned, stay out of smoke.

DIRECTIONS FOR USE:

Use as a fly control supplement for the protection of a manner in the following manner:

Provide one block per 5 head of cattle. Add one additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg. RABON[®] Oral Larvicide per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] program, but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] may be fed to slaughter and to lactating dairy cows.

ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only on cattle manure, but in other existing vegetable matter or silage on the premises.

In order to achieve optimum fly control, ENPROAL[®] FLY CONTROL SUPPLEMENT with RABON[®] should be used in conjunction with other good management and sanitation practices.

Manufactured and Guaranteed By:

V.M.S., INC.

Montgomery, AL 36195 U.S.A.

AT-R0L

EPA Reg. No. 10461-12
EPA EST. No. 10461-AL-1

2/87

10461-12

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: REGISTRATION REREГИSTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	EPA REGISTRATION NO. 10461-12	DATE OF ISSUANCE OCT 15 1986
	TERM OF ISSUANCE Until Reregistered	
	NAME OF PESTICIDE PRODUCT Chloro Control Solution with water	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

160 / 169354
18 / 11

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.


A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Products are hereby registered in accordance with FIFRA section 3(c)(5) that you:

1. Submit and file all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 10461-12."
 - b. Please provide upper and lower limits for both active and inert ingredients.
 - c. Uppercase first letter in the active ingredient Chloro.
 - d. Change "Keep out of lakes, streams or ponds" to "Do not apply directly to water."

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL 	DATE 10/15/86
---	------------------

e. Revise Storage and Disposal Block:

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

PESTICIDE Wastes resulting from the use of this
DISPOSAL: product may be disposed of on-site or at
an approved waste disposal facility.

CONTAINER Open dumping is prohibited. Completely
DISPOSAL: empty bag into mixing equipment. Then
dispose of empty bag in an incinerator
or landfill approved for pesticide
containers, or, if allowed by State and
local authorities, by burning. If burned,
stay out of smoke.

f. Delete "GENERAL CLASSIFICATION":

Any reference to "general use" classification must be removed from Enproal Fly Control Supplement with Rabon labeling. "General Use" as a classification is no longer deemed valid by the Agency, even when products are not subject to restricted use as in the case of your product. If a product is classified for restricted use, this classification notifies individuals that the chemical may be applied only by or under the direct supervision of a certified applicator. The absence of a restricted use classification clearly implies that the aforementioned restriction does not apply to a particular pesticide, or its uses. A second classification for general use is thereby unnecessary and should not appear on your product's labeling.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

25/50 LBS

NET WEIGHT

ACCEPTED
COMMENTS
EPA Label ID: 2/22.7 KG

OCT 15 1986

ENPROAL
FLY CONTROL SUPPLEMENT
WITH RABON® ORAL LARVICIDE

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
amended, for the pesticide
registered under EPA Reg. No.
10-11-12

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	By Weight
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.30% *
INERT INGREDIENTS.....	99.70% **
TOTAL.....	100.00 %

- * FABON® Insecticide - Contains 1.36 g RABON® per pound. RABON® is the registered trademark of Shell Oil Company.
- ** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein, minimum.....	7.50	%
Crude Fat, minimum.....	0.20	%
Crude Fiber, maximum.....	3.00	%
Calcium (Ca), maximum.....	2.75	%
Calcium (Ca), minimum.....	1.75	%
Phosphorus (P), minimum.....	1.00	%
Salt (NaCl), maximum.....	6.00	%
Salt (NaCl), minimum.....	5.00	%
Magnesium (Mg), minimum.....	3.00	%
Potassium (K), minimum.....	2.40	%
Sulfur (S), minimum.....	0.70	%
Cobalt (Co), minimum.....	0.001	%
Copper (Cu), minimum.....	0.03	%
Iodine (I), minimum.....	0.005	%
Iron (Fe), minimum.....	0.19	%
Manganese (Mn), minimum.....	0.12	%
Selenium (Se), minimum.....	0.00028	%
Zinc (Zn), minimum.....	0.21	%
Fluorine (F), maximum.....	0.01	%
Vitamin A, minimum.....	58,000 U.S.P. units	per lb.
Vitamin D-3, minimum.....	10,000 U.S.P. units	per lb.
Vitamin E, minimum.....	25 I.U.	per lb.
Total Sugars (as invert).....	30.00	%

.....
.....
.....

KEEP OUT OF REACH OF CHILDREN (12 pt. type size)

CAUTION (18 pt. type size)

PRECAUTIONARY STATEMENTS (12 pt. type size)

HAZARDS TO HUMANS (12 pt. type size)

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS (12 pt. type size)

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL (12 pt. type size)

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

DIRECTIONS FOR USE:
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 5-8 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the ENPROAL® FLY CONTROL SUPPLEMENT with RABON® program, but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of ENPROAL® FLY CONTROL SUPPLEMENT with RABON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL® FLY CONTROL SUPPLEMENT with RABON® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

ENPROAL® FLY CONTROL SUPPLEMENT with RABON® may be fed to slaughter and to lactating dairy cows.

ENPROAL® FLY CONTROL SUPPLEMENT with RABON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ENPROAL® FLY CONTROL SUPPLEMENT with RABON® should be used in conjunction with other good management and sanitation practices.


MANUFACTURED BY:
V.M.S., INC.
Montgomery, AL 35195
U.S.A.

AT-ROL


2/86

EPA Reg. No. 10461-_____

EPA EST. No. 10461-AL-1



Reg # 10451-12
PRA 15
11/5

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 10451-12	DATE OF ISSUANCE OCT 15 1986
	TERM OF ISSUANCE Until Reregistered	
	NAME OF PESTICIDE PRODUCT Enproal Fly Control Supplement With Rabor.	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)		
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
V.H.S., Inc. 100 Wilbanks Street P.O. Box 406 Montgomery, AL 36195		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:		
1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.		
2. Make the labeling changes listed below before you release the product for shipment:		
a. Add the phrase "EPA Registration No. 10451-12."		
b. Please provide upper and lower limits for both active and inert ingredients.		
c. Uppercase first letter in the active ingredient Chlore.		
d. Change "Keep out of lakes, streams or ponds" to "Do not apply directly to water."		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL 		DATE 10/15/86

160/169354
18/11

e. Revise Storage and Disposal block:

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

PESTICIDE Wastes resulting from the use of this
DISPOSAL: product may be disposed of on-site or at
an approved waste disposal facility.

CONTAINER Open dumping is prohibited. Completely
DISPOSAL: empty bag into mixing equipment. Then
dispose of empty bag in an incinerator
or landfill approved for pesticide
containers, or, if allowed by State and
local authorities, by burning. If burned,
stay out of smoke.

f. Delete "GENERAL CLASSIFICATION":

Any reference to "general use" classification must be removed from Enproal Fly Control Supplement with Rabon labeling. "General Use" as a classification is no longer deemed valid by the Agency, even when products are not subject to restricted use as in the case of your product. If a product is classified for restricted use, this classification notifies individuals that the chemical may be applied only by or under the direct supervision of a certified applicator. The absence of a restricted use classification clearly implies that the aforementioned restriction does not apply to a particular pesticide, or its uses. A second classification for general use is thereby unnecessary and should not appear on your product's labeling.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

25/50 LBS

NET WEIGHT

ACCEPTED
COMMENTS
EPA Label ID: 222.7 KG

OCT 15 1986

ENPROAL
FLY CONTROL SUPPLEMENT
WITH RABON® ORAL LARVICIDE

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
is extended, for the pesticide
registered under EPA Reg. No.

10/16-12

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	By Weight
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.30% *
INERT INGREDIENTS.....	99.70% **
TOTAL.....	100.00 %

* RABON® Insecticide - Contains 1.36 g RABON® per pound. RABON® is the registered trademark of Shell Oil Company.
 ** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein, minimum	7.50	%
Crude Fat, minimum	0.20	%
Crude Fiber, maximum	3.00	%
Calcium (Ca), maximum	2.75	%
Calcium (Ca), minimum	1.75	%
Phosphorus (P), minimum	1.00	%
Salt (NaCl), maximum	6.00	%
Salt (NaCl), minimum	5.00	%
Magnesium (Mg), minimum	3.00	%
Potassium (K), minimum	2.40	%
Sulfur (S), minimum	0.70	%
Cobalt (Co), minimum	0.001	%
Copper (Cu), minimum	0.03	%
Iodine (I), minimum	0.005	%
Iron (Fe), minimum	0.19	%
Manganese (Mn), minimum	0.12	%
Selenium (Se), minimum	0.00028	%
Zinc (Zn), minimum	0.21	%
Fluorine (F), maximum	0.01	%
Vitamin A, minimum	58,000 U.S.P. units per lb	
Vitamin D-3, minimum	10,000 U.S.P. units per lb	
Vitamin E, minimum	25 I.U. per lb	
Total Sugars (as invert)	30.00	%

KEEP OUT OF REACH OF CHILDREN (12 pt. type size)

CAUTION (18 pt. type size)

PRECAUTIONARY STATEMENTS (12 pt. type size)

HAZARDS TO HUMANS (12 pt. type size)

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS (12 pt. type size)

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL (12 pt. type size)

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

DIRECTIONS FOR USE:
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 5-8 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the ENPROAL® FLY CONTROL SUPPLEMENT with RABON® program, but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of ENPROAL® FLY CONTROL SUPPLEMENT with RABON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL® FLY CONTROL SUPPLEMENT with RABON® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

ENPROAL® FLY CONTROL SUPPLEMENT with RABON® may be fed to slaughter and to lactating dairy cows.

