


(BEST AVAILABLE COPY)

PM 15 REG# 10461-12

1

50 LBS NET WEIGHT 22.7 KG



THE ULTIMATE IN MINERAL-VITAMIN NUTRITION
MANUFACTURING
FOR LIVESTOCK

**ENPROAL®
FLY CONTROL
SUPPLEMENT WITH
RABON® ORAL LARVICIDE**



**TAKE TIME
OBSERVE LABEL
DIRECTIONS**

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	By Weight
2-Chloro-1- <i>n</i> -butyl-3-isobutyl-5-trimethylphosphoryl vinyl dimethyl phosphite	Minimum—0.30% Maximum—0.33%
WERT INGREDIENTS	Minimum—98.7% Maximum—99.67%
TOTAL	100.00%

* RABON® Insecticide—Contains 1.36 g RABON® per pound. RABON® is the registered trademark of Shell Oil Company.
* Values only to ingredients which are not toxic.

GUARANTEED ANALYSIS:

Crude Protein, minimum	7.50%
Crude Fat, minimum	8.70%
Crude Fiber, maximum	1.00%
Calcium (Ca), maximum	2.75%
Calcium (Ca), minimum	1.75%
Phosphorus (P), minimum	1.00%
Salt (NaCl), maximum	6.00%
Salt (NaCl), minimum	5.00%
Magnesium (Mg), minimum	3.00%
Potassium (K), minimum	2.40%
Sulfur (S), minimum	0.70%
Cobalt (Co), minimum	0.01%
Copper (Cu), minimum	0.03%
Iodine (I), minimum	0.03%
Iron (Fe), minimum	0.15%
Manganese (Mn), minimum	0.12%
Selenium (Se), minimum	0.002%
Zinc (Zn), minimum	0.21%
Fluorine (F), maximum	0.05%
Vitamin A, minimum	50,000 U.S.P. units per lb.
Vitamin B-1, minimum	10,000 U.S.P. units per lb.
Vitamin E, minimum	25 I.U. per lb.
Total Sugars (as invert)	2.50%

INGREDIENTS:

Case sulfonates, cottonseed meal, dried sugar, magnesium oxide, salt, defluorinated phosphate, magnesium sulfate, potassium sulfate, zinc sulfate, zinc oxide, iron sulfate, iron phosphate, manganese sulfate, manganese oxide, copper sulfate, sodium selenate, sodium selenite, sodium iodate, sodium iodide, cobalt carbonate, vitamin A supplement, D-calciferol, vitamin D-3 supplement, and vitamin E supplement.

NOT DEVELOPED

In Accordance With

10-15-86

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in a dry place in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved state disposal facility.

CONTAINER DISPOSAL: Open dumping is prohibited. Completely empty bag into mixing equipment. Then dispose of empty bag in an incinerator or landfill approved for pesticide containers, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 5-8 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the ENPROAL® FLY CONTROL SUPPLEMENT with RABON® program, but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and feeding areas. Protect from rain. If consumption of ENPROAL® FLY CONTROL SUPPLEMENT with RABON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or rations. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL® FLY CONTROL SUPPLEMENT with RABON® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

ENPROAL® FLY CONTROL SUPPLEMENT with RABON® may be fed to slaughter and to lactating dairy cows.

ENPROAL® FLY CONTROL SUPPLEMENT with RABON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other animal manure, vegetable matter or silage on the premises.

In order to achieve optimum fly control, ENPROAL® FLY CONTROL SUPPLEMENT with RABON® should be used in conjunction with other good management and sanitation practices.

Manufactured and Guaranteed by
V.M.S., INC.
Montgomery, AL 36195 U.S.A.

AT-804

EPA Reg. No. 10461-12
EPA EST. No. 10461-01-1