

11115

1011-1-12

153

50 LB

NET WEIGHT

22.7 KG

ENPROAL®
 FLY CONTROL BLOCK
 WITH RABON® ORAL LARVICIDE

NOT REVIEWED
 in accordance with PR Notice 10-2.
 on draft labelling dated 2/3/90

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

To prevent the development of house flies and stable flies in the manure of pastured horses.

ACTIVE INGREDIENT By Weight

2-Chloro-1-(2,4,3-trichlorophenyl) vinyl dimethyl phosphate*.....	0.30%
INERT INGREDIENTS**.....	99.70%
TOTAL.....	100.00%

* RABON® Insecticide - Contains 1.35 g RABON® per pound. RABON® is the registered trademark of Shell Oil Company.

** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS:

Crude Protein, min.....	7.50%
Crude Fat, min.....	0.20%
Crude Fiber, max.....	3.00%
Calcium (Ca), max.....	2.50%
Calcium (Ca), min.....	2.00%
Phosphorus (P), min.....	1.00%
Salt (NaCl), max.....	6.00%
Salt (NaCl), min.....	5.00%
Magnesium (Mg), min.....	3.00%
Potassium (K), min.....	2.40%
Sulfur (S), min.....	0.70%
Cobalt (Co), min.....	0.001%
Copper (Cu), min.....	0.01%
Iodine (I), min.....	0.005%
Iron (Fe), min.....	0.28%
Manganese (Mn), min.....	0.19%
Selenium (Se).....	0.0013%
Zinc (Zn), min.....	0.0023%
Fluorine (F), max.....	0.01%
Vitamin A, min.....	58,000 U.S.P. units per lb.
Vitamin D-3, min.....	10,000 U.S.P. units per lb.
Vitamin E, min.....	23 I.U. Per lb.
Total Sugars (as invert).....	30.00%

23

FEED INGREDIENTS:

Cane molasses, cottonseed meal, magnesium oxide, salt, defluorinated phosphate, magnesium sulfate, potassium sulfate, zinc oxide, iron carbonate, manganese oxide, sodium selenite, calcium carbonate, calcium iodate, cobalt carbonate, vitamin A supplement, D-activated animal sterol (source of vitamin D-3) and vitamin E supplement.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL


Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only and in a dry place inaccessible to children and pets.

Container Disposal: Completely empty container. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Provide one block per 5-8 head of cattle or horses. Add an additional block when each block has been one-half consumed. Cattle and horses should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved of minerals before initiation of the ENPROAL® FLY CONTROL BLOCK with RABON® program, but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of ENPROAL® FLY CONTROL BLOCK with RABON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL® FLY CONTROL BLOCK with RABON® early in the Spring before flies begin to appear and continue feeding throughout the Summer and into the Fall.

ENPROAL® FLY CONTROL BLOCK with RABON® may be fed to slaughter and to lactating dairy cows.

ENPROAL® FLY CONTROL BLOCK with RABON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle and prevents the development of house flies and stable flies in the manure of treated horses, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle and horse lots and buildings to control adult house flies and stable flies which can breed not only in manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ENPROAL® FLY CONTROL BLOCK with RABON® should be used in conjunction with other good management and sanitation practices.

CAUTION

Follow feeding directions. When this product is consumed as directed, it will provide 0.3 PPM of Selenium (Se).

For specific recommendations on a complete supplement program, contact V.M.S., Inc. Department of Nutrition.

V.M.S. INC.
Montgomery, AL 36195 U.S.A.

(99-AT-ROL)

EPA Reg. No. 10461-12
EPA EST. No. 10461-AL-1

