

prints

0461-12

132

50 LB NET WEIGHT 22.7 KG

MANUFACTURED AND GUARANTEED BY:
V.M.S., INC.
Montgomery, Alabama 36101-8486



ENPROAL

FLY CONTROL SUPPLEMENT WITH RABON[®] ORAL LARVICIDE

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.
To prevent the development of house flies and stable flies in the manure of pastured horses.

ACTIVE INGREDIENTS	BY WEIGHT
E-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate #	0.30%
INERT INGREDIENTS # #	99.70%
TOTAL	100.00%

Each pound contains 1.35 g Rabon per pound.
Refers only to ingredients which are not Larvicidal.
RABON is a registered trademark of Shell Oil Co.

EPA Reg. No. 10461-12
EPA Est. No. 10461-AL-1

GUARANTEED ANALYSIS:

Crude Protein, min	7.50%
Crude Fat, min	0.20%
Crude Fiber, min	3.00%
Calcium (Ca), max	2.50%
Calcium (Ca), min	2.00%
Phosphorus (P), min	1.00%
Salt (NaCl), max	6.00%
Salt (NaCl), min	5.00%
Magnesium (Mg), min	3.00%
Potassium (K), min	2.40%
Sulfur (S), min	0.70%
Cobalt (Co), min	0.001%
Copper (Cu), min	0.01%
Iodine (I), min	0.005%
Iron (Fe), min	0.28%
Manganese (Mn), min	0.19%
Selenium (Se)	0.0013%
Zinc (Zn), min	0.0023%
Fluorine (F), min	0.01%
Vitamin A, min	50,000 I.U. Per Lb.
Vitamin D-3, min	10,000 I.U. Per Lb.
Vitamin E, min	25 I.U. Per Lb.
Total Sugars (as invert)	30.00%

INGREDIENTS:

Cane molasses, cottonseed meal, magnesium oxide, salt, defluorinated phosphate, magnesium sulfate, potassium sulfate, zinc oxide, iron carbonate, manganous oxide, sodium selenite, calcium carbonate, calcium iodate, cobalt carbonate, vitamin A supplement, D-activated animal sterol (source of vitamin D-3) and vitamin E supplement.

NOT REVIEWED

In accordance with PR Notice 82-2.

Based on draft labelling dated 2/13/90

KEEP OUT OF REACH OF CHILDREN
CAUTION!

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS CAUTION!

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only and in a dry place inaccessible to children and pets.

Container Disposal: Completely empty container. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 5-8 head of cattle or horses. Add an additional block when each block has been one-half consumed. Cattle and horses should consume an average of 1.67 lbs. of the block per 100 lbs. of body weight per month. This will supply the recommended average intake of 70 mg RABON ORAL LARVICIDE per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the ENPROAL FLY CONTROL BLOCK WITH RABON program, but omit all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of ENPROAL FLY CONTROL BLOCK with RABON is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding ENPROAL FLY CONTROL BLOCK with RABON early in the Spring before flies begin to appear and continue feeding throughout the Summer and into the fall.

SEE REVERSE SIDE OF THIS SHEET
FOR ADDITIONAL INFORMATION

ENPROAL FLY CONTROL BLOCK with RABON may be fed to slaughter and to dairy cows.

ENPROAL FLY CONTROL BLOCK with RABON prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle and prevents the development of house flies and stable flies in the manure of treated horses, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle and horse lots and buildings to control adult house flies and stable flies which can breed not only in manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ENPROAL FLY CONTROL BLOCK with RABON should be used in conjunction with other good management and sanitation practices.

CAUTION:

Follow feeding directions. When this product is consumed as directed, it will provide 0.3 PPM of Selenium (Se).

For specific recommendations on a complete supplement program, contact V.M.S., Inc. Department of Nutrition.

TAKE TIME TO



OBSERVE LABEL
DIRECTIONS

(99-AT-ROL)

1/90

NOW APPROVED FOR HORSES



"TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED HORSES."

