

10465-1

6-30-1969

1/2

U. S. DEPARTMENT OF AGRICULTURE  
AGRICULTURAL RESEARCH SERVICE  
PESTICIDES REGULATION DIVISION

REGISTRATION NO.

10465-1

35

DATE OF ISSUANCE

June 30, 1969

NAME OF ECONOMIC POISON

COPPER COUNT FUNGICIDE SPRAY CONCENTRATE

**NOTICE OF REGISTRATION UNDER THE FEDERAL  
INSECTICIDE, FUNGICIDE, AND RODENTICIDE ACT**

NAME AND ADDRESS OF REGISTRANT

Mineral Research & Development Corporation  
Box 911  
Charlotte, North Carolina 28201

Attention: Mr. J. T. Rose

**NOTE:** Changes in labeling or formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Pesticides Regulation Division prior to use of the label in interstate commerce. In any correspondence on this product always refer to the above registration number.

On the basis of the information furnished by the registrant, the above named economic poison is hereby registered under Section 4 of the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this registration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Department. In order to protect the public, the Secretary, on his own motion, may at any time cancel the registration of an economic poison in accordance with Section 4 (c) of the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by a trademark held by others.

The following paragraphs are applicable only when checked:

- ☐ To expedite registration, this notice is being issued on the basis of the draft submitted with the application. Certain defects, given below, have been noted. These corrections must be incorporated when the printed labeling is prepared. Five copies of the printed labeling must be submitted to complete the file on this product.
- ☐ The registration for this product is being issued with the understanding that certain defects in the labeling which are noted below will be corrected as soon as possible. Objection is not raised to the use of the present labeling for a reasonable period of time while fully corrected labeling is being prepared. Five copies of the corrected labeling must be submitted.
- ☐

BSH:iba

☐ Attachment is applicable.

Assistant Director  
for Registration


SIGNATURE

APD

~~XXXXXXXXXXXXXXXXXXXX~~

PESTICIDES REGULATION DIVISION

JTT



### DIRECTIONS FOR USE

Use  $\frac{1}{2}$  to  $\frac{3}{4}$  gallon diluted with water per acre in low or high volume sprays. The use of Fungicides for disease control is based on prevention, the plant surfaces must be covered with the Fungicide to successfully prevent infection.

### CAUTION

Harmful if swallowed. Avoid contact with skin. May cause irritation to the nose and throat. In case of ingestion, call physician immediately.

Toxic to fish. Keep this product out of lakes, ponds, or streams. Before re-use, flush container with water and detergent. Do not contaminate water sources by cleaning equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

### NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Registration No. 10465-1

# COPPER-COUNT

## FUNGICIDE SPRAY CONCENTRATE

### FOR USE ON

**BEANS**, for control of common and bacterial blight. Beginning at disease appearance, spray 4 to 5 times at 10-14 day intervals.

**CARROTS**, for control of Early and Late Blight. When plants are 6" high, 3 to 5 applications at 7 to 10 day intervals.

**CELERY**, for control of Early and Late Blight. On appearance of disease, 4 to 7 applications at weekly intervals.

**CUCURBITS**, Cantaloupes, Melons, Cucumbers, Squash, for control of Powdery Mildew, Downy Mildew, Scab, Alternaria and Angular leaf spots. Apply weekly beginning after vining and continue until mid-harvest.

**GRAPES**, for control of Anthracnose. When first leaves appear, just before blooming, just after blooming and at 2 week intervals thereafter until 2 weeks before harvest. Use with 4 pound hydrated lime.

**PEANUTS**, for control Cercospora Leaf Spot. On appearance of disease, 4 to 8 sprays, 10-14 day intervals.

**PEPPERS**, to control Cercospora Leaf Spot, Bacterial Spot. Apply at 7 to 10 day intervals.

**POTATOES (IRISH)**, to control Early and Late Blight. Starting on or before first appearance of the disease in the area, 4 to 6 sprays, 7 to 10 day intervals.

**STRAWBERRIES**, to control Leaf Spot and Scorch. Spray at 7 to 10 day intervals from time new growth starts until harvest.

**SUGAR BEETS**, to control Cercospora Leaf Spot. Starting on disease appearance, 3 to 6 sprays at 14 day intervals.

**TOMATOES**, to control Early and Late Blight and Bacterial Spot, spray in seed beds at emergence and continue on 4 to 5 day schedules. In field, begin applications immediately after transplanting or emergence and continue at 4 to 5 day intervals. In areas or seasons when disease is less severe, time between applications may be 5 to 7 days. Complete coverage is essential to good control.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Cautions

### NET CONTENTS

### GALLONS

### INFO

Since weather conditions vary, consult your State Agricultural Experiment Station for timing and initial application rates. Use proper fittings, nozzles, and a spray solution should be applied to plastic. Do not use in areas with copper.

### FUNGICIDE SP

ACTIVE INGREDIENT  
Copper, Metallic  
(from Copper Ammonium)  
INERT INGREDIENT

Total

Manufactured By

MINERAL RESEARCH  
AND  
DEVELOPMENT

P. O. BOX 911