



DIRECTIONS FOR USE

Use 1/3 to 3/4 gallon diluted with water per acre in low or high volume sprays. The use of Fungicides for disease control is based on prevention, the plant surfaces must be covered with the Fungicide to successfully prevent infection.

CAUTION

Harmful if swallowed. Avoid contact with skin. May cause irritation to the nose and throat. In case of ingestion, call physician immediately.

Toxic to fish. Keep this product out of lakes, ponds, or streams. Before re-use, flush container with water and detergent. Do not contaminate water sources by cleaning equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

USDA Registration No. 10465-1

COPPER-COUNT

FUNGICIDE SPRAY CONCENTRATE

FOR USE ON

BEANS, for control of common and bacterial blight. Beginning at disease appearance, spray 4 to 5 times at 10-14 day intervals.

CARROTS, for control of Early and Late Blight. When plants are 6" high, 3 to 5 applications at 7 to 10 day intervals.

CELERY, for control of Early and Late Blight. On appearance of disease, 4 to 7 applications at weekly intervals.

CUCURBITS, Cantaloupes, Melons, Cucumbers, Squash, for control of Powdery Mildew, Downy Mildew, Scab, Alternaria and Angular leaf spots. Apply weekly beginning after vining and continue until mid-harvest.

GRAPES, for control of Anthracnose. When first leaves appear, just before blooming, just after blooming and at 2 week intervals thereafter until 2 weeks before harvest. Use with 4 pound hydrated lime.

PEANUTS, for control Cercospora Leaf Spot. On appearance of disease, 4 to 8 sprays, 10-14 day intervals.

PEPPERS, to control Cercospora Leaf Spot, Bacterial Spot. Apply at 7 to 10 day intervals.

POTATOES (IRISH), to control Early and Late Blight. Starting on or before first appearance of the disease in the area, 4 to 6 sprays, 7 to 10 day intervals.

STRAWBERRIES, to control Leaf Spot and Scorch. Spray at 7 to 10 day intervals from time new growth starts until harvest.

SUGAR BEETS, to control Cercospora Leaf Spot. Starting on disease appearance, 3 to 6 sprays at 4 day intervals.

TOMATOES, to control Early and Late Blight and Bacterial Spot, spray in seed beds at emergence and continue on 4 to 5 day schedules. In field, begin applications immediately after transplanting or emergence and continue at 4 to 5 day intervals. In areas or seasons when disease is less severe, time between applications may be 5 to 7 days. Complete coverage is essential to good control.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Cautions

Best Available Copy

NET CONTENTS

GALLONS

INFORMATION

Since weather conditions and disease incidence vary, consult your State Agricultural Extension Service for timing and initial applications. Spray Equipment—fittings, nozzles, and all metal parts in contact with spray solution should be steel, iron, aluminum or plastic. Do not use in contact with brass, bronze, copper.

FUNGICIDE SPRAY CONCENTRATE

ACTIVE INGREDIENTS:

Copper, Metallic	8%
(from Copper Ammonium Carbonate)	92%
INERT INGREDIENTS	

Total	100%
-------	------

Manufactured By

MINERAL RESEARCH

AND

DEVELOPMENT CORP.

P. O. BOX 911

CHARLOTTE, N. C. 28201

10465-1

11-3-1969

11/1