

ALPHA 520

Twin Chain Quaternary Ammonium Compound Concentrate

Water Treatment Microbiocide for Building and Industrial Cooling Towers

Active Ingredients

Didecyl dimethyl ammonium chloride	50%
Isopropyl alcohol	20%

Inert Ingredients

EPA Registration No. **P. 9**

Net Weight	U	CT	400 lbs.
------------	---	----	----------

Net Volume 55 qals.

MANUFACTURED BY

UNITED CHEMICAL CORPORATION

OF NEW MEXICO

601 North Lugo P. O. Box 1499
HOBBS, NEW MEXICO 88240

DANGER

KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

USE DIRECTIONS

1. Initially use 6 fluid ounces per 1000 gallons of water to be treated (20 ppm active quaternary).

Should the above dosage not give satisfactory results, use 9 fluid ounces per 1000 gallons of water.

Repeat the initial dose every seven days or increase the frequency if needed.

2. When the above treatment level is successful, use 2 to 3 fluid ounces per 1000 gallons of water to maximize efficiency. Repeat weekly as needed.

Should shoe develop again, go back to initial dosage.

flooding river waters that are inherently low in algal growth and bacteria content may be adequately controlled by the lower range of these dosages slug fed every seven days.

This product is toxic to fish. Improper use or effluent should not be discharged where it will drain into lakes, streams, ponds, or public water.

Should I ever be properly informed of
learning is required

Do not use a negative claim. Reformulate as a reconditioning or disease proof with negative tone, and disease proof. Do not restate negative statement dependent clause.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.