




17-17-17 SYSTEMIC

fertilizer plus insecticide

FOR USE ON BEANS, POTATOES, CORN AND SUGAR BEETS TO CONTROL THE INSECTS SHOWN UNDER DIRECTIONS.

Contains 0.4% THIMET®
Soil Insecticide



Statement of Ingredients

Active Insecticide Ingredients:
 Phorate (O, O-diethyl S-(ethylthio) methyl) phosphorodithioate 0.40%
Insecticidally Inert Ingredients 99.60%

Fertilizer Ingredients

Total Nitrogen (Nitrogen (N)) 17%
 Available Phos. Acid (P₂O₅) 17%
 Water Sol. Potash (K₂O) 17%

DANGER!
Keep Out of Reach of Children



Not for use or storage in or around the home. Poisoning may occur as a result of exposure. May be fatal if swallowed or inhaled. Rapidly absorbed through the skin. Do not breathe dust. Do not get in eyes or on skin. Wash all contaminated clothing with soap and hot water before reuse. Wash thoroughly after handling, and before eating or smoking. Do not contaminate feed or food. Do not reuse empty container. Burn empty bags and stay away from smoke or fumes. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Antidote.— Atropine.

Directions

Thimet is an effective soil and systemic insecticide for early season insect control, which has been combined with fertilizer for efficient and uniform application. Application should be made at planting time, in the row to the side of the seed. Do not apply directly to the seed.

Recommendations for Use on Crops

Crop	Insects	Rate-lbs./acre	Application
<u>POTATOES</u>	Aphids, leafhoppers, leafminers, psyllids, wireworms, flea beetle larve and reduction of flea beetle adults	500 to 750	Granules may be banded on each side of the row at planting time. Use 500 lbs. in sandy or light soils, 750 lbs. in heavy silt, clay loam or muck soils (38" row spacing). Do not apply more than once (1) per season.
<u>CORN</u> At planting time	Corn rootworms	126 to 250	At planting time, in the furrow 1/2" to 1" above seeds on each side of the seed row. Do not place in direct contact with seed. Use 250 lb. rate in areas having high rootworm populations.
<u>SUGAR BEETS</u>	Sugar beet root maggot, leafhoppers, mites, aphids, pea leafminers	250	Apply in 7" band over the row at planting time. Do not place in contact with seed. Make no application later than 30 days before harvest. Do not feed beet tops or silage to dairy cattle.
<u>BEANS</u>	Leafhoppers, aphids, lygus bugs, thrips, mites and seedcorn maggots	250 to 370	Distribute the granules evenly in the row to the side of the seed at planting time. Apply 15 to 20 lbs. per 1000 feet of row. Do not place in direct contact with the seed. Do not feed foliage within 60 days of treatment.

©Trademark of American Cyanamid Company

U.S.D.A. REG. No. 10501-4



FARM BUREAU CO-OPERATIVE
 Antigo, Wisconsin 54409