

RX CATTLE "BUTT" BAG IS A MODERN, NON-MECHANICAL DUST APPLICATOR WHICH CONTAINS 1% CO-RAL © INSECTICIDE. IT IS DESIGNED FOR USE BY BOTH BEEF AND DAIRY CATTLE FOR CONTROL OF HORN FLIES AND AS AN AID IN REDUCING FACE FLY POPULATIONS AS DIRECTED BELOW

# DIRECTIONS

Horn Fly Control:

Suspend CATTLE "BUTT" BAG in areas frequented by cattle or in galeways or lanes through which the animals pass daily for water, feed or mineral. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, CATTLE "BUTT" BAG may be suspended in the exit through which cows leave the milking barn. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle.

Horn Fly Control and as an Aid in Reducing Face Fly Populations:

Suspend CATTLE "BUTT" BAG in the same manner as described under "Horn Fly Control" except as follows: (1) The bottom of the bags should be 18 to 24 inches from the ground to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily.

# INSTALLATION OF CATTLE "BUTT" BAG:


Suspend the bags by inserting rope, chain, or wire through the grommets on the side of the bag and attaching to an overhead structure or by hooking the grommets into hooks fixed to a stationary overhead 2x4 at the heights recommended for horn fly or face fly control. In areas of heavy or frequent rainfall the CATTLE "BUTT" BAG should be protected from the weather.

©CO-RAL is a registered trademark of Farbenfabriken Bayer A.G. Chemagro Corporation Licensee U.S.D.A. REG. NO. 10504-1



CAUTION:
KEEP OUT OF THE
REACH OF CHILDREN

H. ROSE ROSE ONLY



# Do NOT hang "BUTT" BAG over feed, mineral or water troughs. CAUTIONS

Harmful if inhaled or absorbed through the skin. Do not get in eyes, on skin or clothing. Do not breathe dust. Wash thoroughly with soap and water after handling. Do not contaminate feed or food. Keep out of reach of children. If poisoning occurs obtain prompt medical aid. Prolonged exposure will result in choline-sterase depression.

NOTE TO PHYSICIAN — Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.


## CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.


| FOR CONTROL OF H                           |   | _ | - | <br> | D IN | I RE  | DUCII | NG F | ACE FLY         |
|--|---|---|---|------|------|-------|-------|------|-----------------|
| ACTIVE INGREDIENTS  Q Q-Diethyl Q- (3-chlo | 3 |   |   |      | zonv | ran-7 | 7-v1) |      | BY WT.          |
| phosphorothicate+.                         |   |   |   |      |      |       |       |      | . 1.0%          |
| ENERT INGREDIENTS.                         |   |   |   |      |      |       |       | •    | . 99.0%         |
|  |   |   |   |      |      |       |       |      | <b>—</b> 100.0% |

• U. S. PATENT NO. 2.748.146 CANADIAN PATENT NO. 523.206









EX CATTLE "BUTT" BAG IS A MODERN, NON-MECHANICAL DUST APPLI-CATOR WHICH CONTAINS 1% **CO-RAL ® INS**ECTICIDE. IT IS DESIGNED FOR USE BY BOTH BEEF AND DAIRY CATTLE FOR CONTROL OF HORN FLIES AND AS AN AID IN REDUCING FACE FLY FOPULATIONS AS DIRECT ED BELOW

# DIRECTIONS

Hom Fly Control:

Suspend CATTLE "BUTT" BAG in areas frequented by cattle or in galeways. or lanes through which the animals pass daily for water, feed or mineral. Bags may also be suspended in loafing sheds or in front of protected milleral feeders. For lactating dairy cows CATTLE "BUTT" BAG may be suspended in the exit through which cows leave the milking larm. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the tapline of the carle.

Hern Fly Centrel and as an Aid in Reducing Face Fly Populations:

Suspend CATTLE "BUTT" BAG in the same manner as described under "Hom Fly Control" except as follows: (1) The hottom of the hags should be 18 24 inches from the ground to insure that the faces of the cattle will be dusted. 2) The hags rust be located so the animals will be forced to use them daily.

# INSTALLATION OF CATTLE "BUTT" BAG:

Suspend the kags ky inserting rope, chain, or wire through the grommets on the inde of the hag and attaching to an overhead structure or by hooking the prominets into hilloks fixed to a stationary overhead 2x4 at the heights recomrended for hore fly or face fly control. In areas of heavy or frequent rainfall the CATTLE "BUTT" BAG should be protected from the weather.

Property and the first to be the Fight problems Payer A.G. Chemistry Company of the same o

U.S.D.A. REG. NO 10504-1

MANUFACTURED FOR RX VETERINARY PRODUCTS FRESNO, CALIFORNIA 93706



KEEP OUT OF THE REACH OF CHILDREN

CAUTION:

Install as many "BUTT" BAGS as necessary in lanes or other openings to accure proper dusting of the animals. After the bags are in place, pull the reploord to allow the insecticide to fall into the bottom of the bag. When dust bags are grovided on a free-choice basis, allow one 25 pound "BUTT" BAG per 25 to 50 animals.

RESTRICTIONS:

#### Do NOT hang "BUTT" BAG over feed, mineral or water troughs. CAUTIONS

Harmful if inhaled or absorbed through the skin. Do not get in eyes, on skin or lothing. Do not breathe dust. Wash thoroughly with soap and water after handling. Do not contaminate feed or food. Keep out of reach of children. If poisoning occurs obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

NOTE TO PHYSICIAN — Atropine sulfate is antidotal. 2-PAM is also antidotal. and may be administered in conjunction with atropine.

## CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer mixed no warranties, expressed or implied concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

TOR CONTROL OF HORN FLY AND AS AN AID IN REDUCING FACE FLY POPULATIONS ON BEEF AND DAIRY CATTLE. **ACTIVE INGREDIENTS** BY WT  $\bigcirc$  Q-Diethyl Q- (3-chloro-4-methyl-2-exo-(2H)-1-benzonyran-7-y) The sphorochicate but NERT INGREDIENTS. . 99.0%

TOTAL

100.0°

TO SPATIANT NO 2, 48446 CANADIAN PATENT NO 5/3/296

1% CO-RAL® NET CONTENTS 25 LBS.

MANUFACTURED FOR RX VETERINARY PRODUCTS FRESNO, CALIFORNIA 93706