

Front Panel

SSAX

READY TO USE SPRAY DIRECT FROM BOTTLE

A SELECTIVE SYSTEMIC WEED KILLER

Kills Tough Broad leaf and Bushy Weeds and Grasses

ACTIVE INGREDIENTS

Monosodium Acid Methanearsonate.....0.57%

INERT INGREDIENTS.....99.43%

Total arsenic (as elemental) in water soluble form is .55%. *Product contains .15 pounds of Monosodium acid methanearsonate per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional
Precautionary Statements.

NET CONTENTS: 1 QUART (32 OUNCES)
SHAKE WELL BEFORE USING

Manufactured by
GENERAL CONTROL COMPANY, INC.
TUCSON, ARIZONA

EPA Est. No. 10583-AZ-1 EPA No. 10533-EN

Accepted
for sale
MAR 20 1990
EPA Reg. No.
10583-20

Left Side Panel

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Wear protective clothing when handling or applying this product including long pants, long-sleeved shirt, and impermeable gloves and boots. Keep children and domestic animals off treated areas until this product is washed into the soil. Hose down with plenty of water the treated plants one day after spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person. Call a Doctor.

IF ON SKIN: Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water for fifteen minutes. Call a Doctor.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. This product is toxic to fish and other cold blooded animals. Do not contaminate streams, ponds, lakes or other water by the cleaning of equipment or the disposal of container or wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Avoid storing near feed, seed, or food products.

PRODUCT DISPOSAL: Securely wrap container in several layers of paper and discard in trash.

CONTAINER DISPOSAL: Do NOT REUSE container. Rinse thoroughly before discarding in trash receptacle.

BEST AVAILABLE COPY

Right Side Panel

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PLEASE READ DIRECTIONS COMPLETELY BEFORE USING.

This product is designed primarily for spot control of perennial (year round) weeds in growing Bermudagrass lawns as well as in ground cover such as Lippia and Rosemary. This product is a selective systemic: it can kill top growth and the roots of perennial (year round) weeds such as; Nut Grass, Perennial Rye, Wild Carrot, Oxalis, Crab Grass, and Poa Annua (Blue Grass) in one or more applications without killing Bermudagrass Lawns. It is also designed to kill tough woody perennial such as Salt Bush, Desert Broom, and Tumble Weed, as well as Ragweed and White Horse Nettle.

HOW TO APPLY: Spray directly on the foliage (leaves) of the weed until wet. It is not necessary to soak. When sprayed on the green foliage of a weed, this herbicide will move to the root system and destroy it. Much foliar (leafy) area is needed for it to be absorbed, therefore allow the weed to grow at least 3-5 inches before spraying.

Spray the entire weed until wet. The next day hose down with plenty of water all treated plants. Three to five days later weeds will show stress and start to dry and brown out. Annual (seasonal) weeds and grasses are now destroyed. Perennials, though, will require re-treatment. Re-spray any part of the plant that remains green after five days and again when green reappears in about three weeks. Good control can be established within one growing season, but complete eradication may take two or more seasons. When using this material on a tough woody perennial it will be necessary to spray the weed more than once. After the first application, when any growth shows through dead patches, reapply until the weed has fully absorbed the herbicide into its root system and is dead. When Spot spraying in Bermudagrass lawn and ground cover, slight brown spotting may appear, especially in hot weather. This is temporary, though, and will disappear with new growth. This material will cause brown spots in St. Augustine lawns and in Dichondra. Test a small area before using.

COVERAGE: Approx. 250 sq. ft.

BEST AVAILABLE COPY

NOTICE: Neither Manufacturer nor Seller makes any warranty, express or implied, concerning the use, effects of, or results from use of this product, other than as indicated on this label. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions contained on this label.