

DIRECTIONS

FOR THE CONTROL OF GARDEN INSECTS - For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles, aphids, exposed thrips and spider mites. Hold container at least 18" from foliage, spray lightly in one-second bursts spraying both top and underside of foliage.

This spray may be used on such plants as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuchsia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snap dragons, stocks, Wandering Jew and zinnias.

As a fungicide, spray before infection occurs for the prevention of powdery mildew and black spot on roses. Spray at five to seven day intervals while danger of fungus infestation persists.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing spray or contact with eyes or skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs.

Do not spray closer than 18 inches from plants. Do not use near fishbowls or aquaria.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerate.

NOTICE: Seller makes no warranty, express or implied, concerning the use of this product other than stated on the label. Buyer assumes all risk of use, storage and/or handling of this material when such use, storage and/or handling is contrary to label instructions.

MANUFACTURED FOR
LOS ANGELES SEED CO., INC.

LOS ANGELES, CALIFORNIA 90021

EPA Reg. No. 10631-3

Cal. Reg. No. 06222-CA-1

Angel City

**A SPECIAL BLEND OF INSECTICIDES,
FUNGICIDES, AND MITICIDES**

ROSE Spray

**CONTROLS MAJOR ROSE PESTS AND
DISEASES SUCH AS APHIDS, POWDERY
MILDEW, BLACK SPOT AND SIMILAR
FLOWER GARDEN PESTS**

ACTIVE INGREDIENTS:

Pyrethrins	0.025%
Piperonyl Butoxide, Technical*	0.256%
Rotenone	0.128%
Other Cube Extractives	0.237%
Folpet; N-(Trichloromethylthiophthalimide)	0.700%
Carbaryl; 1-naphthyl N-methylcarbamate	1.000%
Petroleum Distillate	0.025%

INERT INGREDIENTS: 97.629%

*Equivalent to 0.205% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.051% of related compounds.

NET WEIGHT 14 OZ.