

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces that will not contact food products in homes, schools, rest rooms and other public places, mop or sponge surface to be treated with a solution containing 1 ounce of ALPHA BAC in 2 gallons of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection. This disinfectant solution will effectively destroy the gram positive microorganism, Staphylococcus aureus and the gram negative microorganism, Salmonella choleraesuis.

HOSPITAL DISINFECTION

To disinfect walls, floors, furnishing and other hard surfaces in hospitals, nursing homes, laboratories and other medical environments, mop or sponge surface with a solution containing 1 ounce of ALPHA BAC per gallon of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection.

Surgical or medical instruments must be thoroughly scrubbed to remove soil and then disinfected by immersion in a solution containing 1 ounce of ALPHA BAC per gallon of water for 10 minutes. One tablespoon of sodium nitrite should be added to each gallon of disinfectant solution to inhibit corrosion of metal instruments.

This disinfectant solution will effectively destroy the gram positive microorganism Staphylococcus aureus, and the gram negative microorganism Salmonella choleraesuis, and Pseudomonas aeruginosa.

BARBER AND BEAUTY PARLOR EQUIPMENT DISINFECTION

Thoroughly clean brushes, combs, razors and shears and rinse free of soil before disinfection. Soak instruments in a solution containing 1 ounce of ALPHA BAC per gallon of water for 10 minutes. Use 1 tablespoon of sodium nitrite per gallon of solution to prevent corrosion of metal instruments.

POULTRY EQUIPMENT DISINFECTION

Poultry brooders, watering founts and feeding equipment may be disinfected after thorough cleaning by sponging or brushing with a solution containing 1 ounce of ALPHA BAC in 2 gallons of water. Small utensils should be immersed in this solution. NOTE: All equipment that will contact food or drinking water must be rinsed with water of drinking quality after disinfection and before reuse.

Hatching eggs may be disinfected by dipping in a solution containing 1 ounce of ALPHA BAC per 2 gallons of water. Do not use this product on eggs intended for human consumption.

KENNELS AND ANIMAL QUARTERS DISINFECTION

To disinfect and deodorize animal quarters mop or sponge a solution containing 1 ounce of ALPHA BAC in 2 gallons of water to cleaned floors and walls. Feeding equipment disinfected with this solution should be rinsed with water of drinking quality before reuse.



FUNGICIDE
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 10634-19

USE DIRECTIONS FOR CLEANING

Lightly soiled areas – surfaces to be cleaned should be mopped or brushed with a solution contain 1 ounce of Alpha-Bac 200 per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas – mop or brush soiled surface with a solution containing 2-3 ounces of Alpha-B 200 per gallon of water. Rinse cleaned surface with fresh water.

GENERAL-PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of Alpha-Bac 200 per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of Alpha-Bac 200 per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas where fungicidal action is required, mop area with a solution containing 2 ounces in 3 gallons of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of Alpha-Bac 200 per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce of Alpha-Bac 200 per gallon of water. If surface is heavily soiled, use Alpha-Bac 200 at a rate of 2-3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing ½ ounce of Alpha-Bac 200 per gallon of water. No rinse required after sanitization.