

10634-13

PM-17

12/20/89

131

CAUTION

Harmful if swallowed. Avoid contamination of feed or feedstuffs. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation. Cover water, drinking fountains, milk utensils, and feed before use.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of spray equipment or disposal of wastes.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME.

WARRANTY:

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS

ON ANIMALS: To protect animals from horn flies, house flies, mosquitoes, and gnats, apply a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.

CAUTION
KEEP OUT OF REACH OF CHILDREN
See side panel for additional precautions

**ALPHA CHEM:
READY TO USE
ANIMAL SPRAY**

A Ready-To-Use PYRENONE Insecticide
KILLS AND REPELS

Kills Houseflies, Stable Flies, Horse Flies, Mosquitoes, Gnats, and Lice.
Acceptable for use on Dairy and Meat Animals and in Dairy Barns, Milk Rooms, and Caines.

ACTIVE INGREDIENTS:

Pyrethrins 0.10%
*Piperonyl Butoxide, Tech. 1.00%

****INERT INGREDIENTS 98.90%**

****Contains Petroleum Distillates**

*Equivalent to 0.8% (butylcarbitol)(6-propylpiperonyl) ether and 0.2% of related compounds.

PYRENONE is a registered trademark of Fairfield American Corp.

NET CONTENTS: 1 U.S. GALLON E.P.A. REG. NO. 10634-13
E.P.A. EST. NO. 10634-MA-1

MANUFACTURED BY
ALPHA CHEMICAL SERVICES, INC. — STOUGHTON, MASS.

To control blood-sucking lice, apply the spray lightly to infested areas, using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two or three weeks as required.

BARN, MILK ROOMS, DAIRY PLANTS: To control flies, mosquitoes, and gnats, apply as a fog or fine mist, directing spray toward the ceiling and upper corners, until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES

Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL:

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL:

Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

BEST AVAILABLE COPY

Not Reviewed. Registrant claims to be in accordance with reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.