

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink promptly a large quantity of water. Do not induce vomiting. Avoid alcohol. Get medical attention.

If In Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN

Produce mucosal damage may contraindicate the use of gastric lavage. Measure against convulsions. Convulsory shock and respiratory depression may not be needed.

DIRECTIONS FOR USE


It is a violation of a federal law to use this product in a manner inconsistent with its labeling.

CLEANING - MILKING MACHINES, UTENSILS, ETC.

Use 1 ounce of ICS in 5 gallons of warm water. (Provides 25 PPM available iodine). Brush all surfaces thoroughly. Rinse with clear water and drain.

BULK TANKS

Use 1 ounce of ICS in 2 1/2 gallons of warm water. (Provides 50 PPM available iodine). Brush all surfaces thoroughly. Rinse with clear water and drain.



DANGER
KEEP OUT OF REACH OF CHILDREN
See side panel for additional precautions

AN IODINE CLEANER-SANITIZER FOR USE ON DAIRY FARMS

ICS

Active Ingredients:
*Iodine.....1.75%
Phosphoric Acid(100%).....16.0%
Inert Ingredients.....82.25%

*From Alpha-(p-nonylphenyl)-omega-hydroxypoly (oxyethylene)-iodine complex
Net Contents: 1 U.S. Gallon
EPA Reg. No. 10634-15
EPA Est. 10634AA-1

ALPHA CHEMICAL SERVICES, INC.
STOUGHTON, MASSACHUSETTS 02172

SANITIZING

To sanitize dairy and food processing equipment, prepare a 12 1/2 ppm solution by adding 1/2 oz

ICS to 5 gallons of water. Thoroughly clean equipment with a suitable detergent and rinse with potable water to remove all soil and milk residue. Immerse or thoroughly wet equipment with the 12 1/2 PPM ICS solution for at least one minute. Remove and allow to air dry. Do not rinse with potable water before reuse. Prepare a fresh solution daily or more frequently as soil is apparent.

UDDER WASHING

Use 1/4 oz. of ICS in 2 1/2 gallons of warm water. (Provides 25 PPM available iodine). Wash each cow's udder with a separate paper towel saturated in this solution. Change solution often to maintain clarity and warmth.

NOTE

The yellowish-brown amber color of ICS is its indicator of germicidal activity. Maintain this color in all cleaning and sanitizing phases. When the color fades, prepare a fresh solution.

Do not use this product as a test kit. ICS qualifies under the federal regulation 171.1010 (b)(3) eliminating the need for a final potable water rinse of use-dilutions providing up to 25 PPM of available iodine.

STORAGE AND DISPOSAL

STORAGE: Store in original container. Avoid temperatures between 32°F to above 130°F. Use container before reuse when labeling. Precaution labels of replacement parts and gaskets of 100 parts are recommended when reusing product. Do not use a container previously used for other product. In case of spill or leak, neutralize spill with Sodium Thiosulfate or Sodium Sulfite. Flush with water. DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

DISPOSAL: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility. Consult Federal State or Public Contractors. Treat spill or equipment. Treat spill or equipment of replacement, or structure and dispose of in an sanitary landfill. If a container of 5 gallons or more and not completely full, empty it before use of it.

- 1. Scrape and gl
- 2. Wash compa
- 3. Rinse
- 4. Sanitize 5 galls Immer two m specifi code.
- 5. Place rack of